

REFERENCES

- ABOLAFIA, J., HOLOVACHOV, O., BOSTRÖM, S. & PEÑA-SANTIAGO, R. (2003). Nematodes of the order Rhabditida from Andalucía Oriental, Spain. The genus *Cervidellus* Thorne, 1937, (Cephalobidae) with additional data on *C. hamatus* Thorne, 1937. *Nematology* 5, 753-776.
- ABOLAFIA, J., LIÉBANAS, G. & PEÑA-SANTIAGO, R. (2002). Nematodes of the order Rhabditida from Andalucía Oriental, Spain. The subgenus *Pseudacrobeles* Steiner, 1938, with a description of a new species. *Journal of Nematode Morphology and Systematics* 4, 137-154.
- ABOLAFIA, J. & PEÑA-SANTIAGO, R. (2001). Rhabditid species (Nematoda, Rhabditida) recorded in Peninsular Spain and Balearic Islands. *Graellsia* 57, 113-131.
- ABOLAFIA, J. & PEÑA-SANTIAGO, R. (2002). Nematodos de orden Rhabditida de Andalucía Oriental. El género *Eucephalobus* Steiner, 1936. *Graellsia* 58, 59-78.
- ABOLAFIA, J. & PEÑA-SANTIAGO, R. (2003a). Nematodes of the order Rhabditida from Andalucía Oriental, Spain. The genus *Acrobeloides* (Cobb, 1924) Thorne, 1937, with description of *A. arenicola* sp. n. and a key to species. *Journal of Nematode Morphology and Systematics* 5, 107-130.
- ABOLAFIA, J. & PEÑA-SANTIAGO, R. (2003b). Nematodes of the order Rhabditida from Andalucía Oriental, Spain. The genus *Chiloplacus* Thorne, 1937, with description of *C. maginensis* sp. n. and a key to species. *Nematology* 5, 243-274.
- ABOLAFIA, J. & PEÑA-SANTIAGO, R. (2003c). Nematodes of the order Rhabditida from Andalucía Oriental, Spain. The genera *Nothacrobeles* Allen & Noffsinger, 1971 and *Zeldia* Thorne, 1925. *Journal of Nematology* 35, 233-243.
- ABOLAFIA, J. & PEÑA-SANTIAGO, R. (2003d). Nematodes of the order Rhabditida from Andalucía Oriental, Spain. The genus *Stegelletina* Andrassy, 1984, with description of two new species. *Russian Journal of Nematology* 11, 37-53.
- ABOLAFIA, J. & PEÑA-SANTIAGO, R. (2005). Nematodes of the order Rhabditida from Andalucía Oriental: *Pseudacrobeles elongatus* (de Man, 1880) comb. n. *Nematology* 7, 917-926.
- ABOLAFIA, J. & PEÑA-SANTIAGO, R. (2006). Nematodes of the order Rhabditida from Andalucía Oriental, Spain. The family Panagrolaimidae, with a compendium of species of *Panagrolaimus* and a key to their identification. *Journal of Nematode Morphology and Systematics* 8, 133-160.
- ABOLAFIA, J. & PEÑA-SANTIAGO, R. (2007). Nematodes of the order Rhabditida from Andalucía Oriental, Spain. The genera *Protorhabditis* (Osche, 1952) Dougherty, 1953 and *Diploscapter* Cobb, 1913 with description of *P. spiculocrestata* sp.n. and a species *Protorhabditis* key. *Journal of Nematology* 39, 263-274.
- ABOLAFIA, J. & PEÑA-SANTIAGO, R. (2009a). Nematodes of the order Rhabditida from Andalucía Oriental, Spain. The genus *Ablechroiulus* Andrassy, 1966 with description of three new species and key to species. *Nematology* 11, 77-94.

- ABOLAFIA, J., PEÑA-SANTIAGO, R. (2009b). Nematodes of the order Rhabditida from Andalucía Oriental, Spain. The genus *Cephalobus* Bastian, 1865 with description of *C. harpagonis* sp. n. and key to species. *Nematology* 11, 485-508.
- ADAM, W. (1930). Ein neuer freilebender Nematoden aus der Erder: *Butlerius filicaudatus* n. sp. *Zoologischer Anzeiger* 91, 139-142.
- AHMAD, I., AHMAD, W. & MAHMOOD, M. (2004). *Peterngus nepenthi* gen. n., sp. n. (Diplogastridae) from Pitcher plants in Singapore. *International Journal of Nematology* 1, 81-84.
- AHMAD, I. & JAIRAJPURI, M. S. (1979). Developmental biology of *Chiloplacus symmetricus*. *Indian Journal of Nematology* 7, 123-139.
- AHMAD, I., SHAH, A.A. & MAHAMOOD, M. (2005). Nematodes of the order Rhabditida from India. Description of a new and a known species of *Diplogastrellus* (Diplogastridae). *Journal of Nematode Morphology and Systematics* 7, 159-168.
- AHMAD, I., SHAH, A.A. & MAHAMOOD, MD. (2007). Nematodes of the Order Rhabditida from India. Description of *Sclerorhabditis tridentatus* gen. n., sp.n. (Rhabditidae). *Nematology* 9, 43-47.
- AHMAD, I., SINGH, G.K., YEON, II-KWEON & CHUNG-DON CHOI (2009). Redescription of *Butlerius butleri* Goodey, 1929 (Nematoda: Diplogasteridae) from South Korea. *Nematology*, 11, 161-169.
- AHMAD, N. & JAIRAJPURI, M. S. (1982). Observations on the development of juveniles and adult of *Parahadronchus shakili* (Jairajpuri, 1969). *Revue de Nematologie* 5, 79-91.
- ALI, S.M., SURYAWANSHI, M.V. & CHISTY, K.Z. (1973). Two new species of *Drilocephalobus* Coomans and Goodey, 1965 (Nematoda: Drilocephalobidae n. fam.) from Marathwada, India, with a revised classification of subfamily Cephalobinae (Paramonov, 1956) Paramonov, 1962. *Nematologica* 19, 308-317.
- ALLEN, M.W. & NOFFSINGER, E.M. (1971). *Nothacrobeles* n. gen., with description of four new species (Nematoda: Cephalobidae). *Journal of Nematology* 3, 145-153.
- ALLEN, M.W. & NOFFSINGER, E.M. (1972). A review of the genus *Zeldia* Thorne, 1937 (Nematoda: Cephalobidae) with description of seven new species. *Proceedings of the Helminthological Society of Washington* 39, 206-223.
- ALLGÉN, C. (1933). Über einige freilebende Nematoden aus dem Niederkongo. *Zoologischer Anzeiger* 103, 312- 320.
- ALLGÉN, C. (1935). Die freilebenden Nematoden des Öresunds. *Capita Zoologica* 6, 5-192.
- ALLGÉN, C. (1947). Weitere Untersuchungen über schwedische Nematoden. I. *Glauxinema filicaudatum* n. gen. n. sp., eine neue Anguillulide von der Westküste Schwedens. *Kungliga Fysiografiska Sällskapets Förhandlingar (Lund)* 17, 73-78.
- ALLGÉN, C. (1952). Über einige freilebende Süßwasser-Nematoden von Mt. Kenya (Brit. Ostafrika). Ein kleiner Beitrag zur Kenntnis tropischer Nematoden. *Ark Für Zoologie* 2, 139- 157.
- ALTHERR, E. (1938). La faune des mines de Bex, avec étude spéciale des Nématodes. *Revue Suisse de Zoologie* 45, 567-720.

- ALTHERR, E. (1952). Les nématodes du Parc national suisse. (Nemaodes libres du sol.) 2^e partie. *Ergebnisse der Wissenschaftlichen Untersuchung des Schweizerischen National Parks* 3, 315-356.
- ALTHERR, E. (1963). Contribution à la connaissance de la faune sables submergés en Lorraine. Nématodes. *Annales de Spéléologie* 18, 53-98.
- ALTHERR, E. (1976). Nématodes des eaux stygorhitrales des Alpes autrichiennes. *Revue Suisse de Zoologie* 83, 779-847.
- AMBROS, V. & FIXSEN, W. (1987). Cell lineage variation among nematodes. *In: Development as an evolutionary process* (Eds. Raff, R.A. & Raff, E.C.) Alan R. Liss, New York, 39-159.
- ANDERSON, R.V. (1965). *Acrobeloides uberrinus* n. sp. with a note on morphologic variation within soil and bacteria-reared populations. *Proceedings of Helminthological Society of Washington* 32, 232-235.
- ANDERSON, R.V. (1969). Comparative morphology and descriptions of three new species of *Teratocephalus* from Canada. *Canadian Journal of Zoology* 47, 829-840.
- ANDERSON, R.V. (1979). An emendation of the genus *Teratorhabditis* (Osche, 1952) Dougherty, 1953 and description of *T. stiannula* sp. n. (Rhabditidae: Nematoda). *Canadian Journal of Zoology* 57, 13-17.
- ANDERSON, R.V. (1983). Description of *Distolabrellus veechi* n. gen., n. sp. (Nematoda: Rhabditidae). *Journal of Nematology* 15, 70-75.
- ANDERSON, R.V. & DARLING, H.M. (1964). Embryology and reproduction of *Ditylenchus destructor* Thorne, with emphasis on gonad development. *Proceedings of the Helminthological Society of Washington* 31, 240-256.
- ANDERSON, R.V. & HOOPER, D.J. (1970). A neotype for *Cephalobus persegnis* Bastian, 1865, redescription of the species and observations on variability in taxonomic characters. *Canadian Journal of Zoology* 48, 457-469.
- ANDERSON, R.V. & HOOPER, D.J. (1971). A neotype for *Eucephalobus striatus* (Bastian, 1865) Thorne, 1937 (Nematoda) and redescription of the species from topotypes and their progeny. *Canadian Journal of Zoology* 49, 451-459.
- ANDRÁSSY, I. (1958). Erd- und süßwassernematoden aus Bulgarien. *Acta Zoologica Academiae Scientiarum Hungaricae* 4, 1-88.
- ANDRÁSSY, I. (1958). Szabadonélő fonálférgék Nematoda libera. *Magyarország állatvilága fauna Hungariae* 3, 1-362.
- ANDRÁSSY, I. (1959). Nematoden aus dem Psammon des Adige- Flusses, I. *Memoire del Museo Civico di Storia Naturale di Verona* 7, 163-181.
- ANDRÁSSY, I. (1962). Neue Nematoden-Arten aus Ungarn. 1. Zehn neue Arten der Unterklasse Secernentea (Phasmidia). *Acta Zoologica Academiae Scientiarum Hungaricae* 8, 1-23.
- ANDRÁSSY, I. (1964). Neue Nematoden- Arten aus Ungarn, III. Fünf neue Arten. *Opuscula Zoologica Budapest* 5, 9-23.

- ANDRÁSSY, I. (1966). Erd- und Süßwasser-Nematoden aus Ghana-Klass Secernentea (Phasmidia). *Annales Universitatis Scientiarum Budapestinensis de Rolando Eötvös Nominatae, Sectio Biologica* 8, 5-24.
- ANDRÁSSY, I. (1967). Die unterfamilie Cephalobinae (Nemaoda. Cephalobidae) und ihre Arten. *Acta Zoologica Hungarica* 13, 1-37.
- ANDRÁSSY, I. (1968). Fauna Paraguayensis. 2. Nematoden aus den Galeriewäldern des Acaray-Flusses. *Opuscula Zoologica Budapest* 8, 167-315.
- ANDRÁSSY, I. (1970). Freilebende Nematoden aus Vietnam. *Opuscula Zoologica Budapest* 10, 5-31.
- ANDRÁSSY, I. (1971). Különös jelenség: fonálférgekben élősködő fonálférgek! (Eine merkwürdige Erscheinung: parasitare Nematoden in Nematoden). *Álattani Közlemenyek* 58, 156-159.
- ANDRÁSSY, I. (1973). *Plectus granulosus* var. *grandepapillatus* Ditlevesen- ein alteres synonym von *Anaplectus submersus* (Hirschmann) (Nematoda: Plectidae). *Opuscula Zoologica Instituti Zoosystematici Unversitatis Budapestinensis* 12, 105-106.
- ANDRÁSSY, I. (1974a). Über vier homonyme Nematodengattungen. *Nematologica* 19, 403-404.
- ANDRÁSSY, I. (1974b). A Nematodák evolúciója és rendszerezése. (The evolution and systematization of Nematoda). *Magyar Tudományos Akadémia Biológiai Osztálya Közlem* 17, 13-58.
- ANDRÁSSY, I. (1976). *Evolution as a basis for the systematization of nematodes*. Budapest. London, Pitman, 288 p.
- ANDRÁSSY, I. (1977). Die Gattungen *Amphidelus* Thorne, 1939. *Paramphidelus* n. gen. and *Etamphidelus* n. gen. (Nematoda: Alaimidae). *Opuscula Zoologica Budapest* 14, 3-43.
- ANDRÁSSY, I. (1978). *Nematoda*. A checklist of the animals inhabiting European inland waters with account of their distribution and ecology. In: (*Limnofauna Europaea*. Ed. Illies, J.), 98-117.
- ANDRÁSSY, I. (1981). Revision of the order Monhysterida (Nematoda) inhabiting soil and inland waters. *Opuscula Zoologica Budapest* XVII- XVIII, 13- 47.
- ANDRÁSSY, I. (1982). Six new species of the suborder Rhabditina (Nematoda: Secernentia). *Revue de Nématologie* 5, 39-50.
- ANDRÁSSY, I. (1983a). A taxonomic review of the suborder Rhabditina (Nematoda- Secernentia) Orstom, Paris, 241 p.
- ANDRÁSSY, I. (1983b). A taxonomic review of the suborder Rhabditina (Nematoda). *Revue de Nématologie* 5, 39-50.
- ANDRÁSSY, I. (1983c). The free-living nematode fauna of the Hortobágy National Park. In: *The fauna of the Hortobágy National Park, Budapest, Hungary*, (Ed. Mahunka, S.). Akadémiai Kiadó, pp. 31-46.
- ANDRÁSSY, I. (1984). *Klasse Nematoda* (Ordungen Monhysterida, Desmoscolecida, Araeolaimida, Chromadorida, Rhabditida). Gustav Fischer Verlag. Stuttgart, pp 509.
- ANDRÁSSY, I. (1985a). The genus *Plectus* Bastian, 1865 and its nearest relatives (Nematoda: Plectidae). *Acta Zoologica Hungarica* 31, 1- 52.

- ANDRÁSSY, I. (1985b). A Dozen new nematode species from Hungary. *Opuscula Zoologica Budapest*. XIX-XX, 3-39.
- ANDRÁSSY, I. (1986). Fifteen new nematode species from the Southern Hemisphere. *Acta Zoologica Academiae Scientiarum Hungaricae* 32, 1-33.
- ANDRÁSSY, I. (1987). The free-living nematode fauna of the Kiskunság National Park. - In: The fauna of the Kiskunság National Park. Budapest. 15-46.
- ANDRÁSSY, I. (1998). Nematodes in sixth continent. *Journal of Nematode Morphology and Systematics* 1, 107-186.
- ANDRÁSSY, I. (2005a). Free-living nematodes of Hungary (Nematoda errantia). In: *Paedozoologica Hungarica* (Eds. Czudi, C. & Mahunka, S) vol.I, 518p.
- ANDRÁSSY, I. (2005b). Free-living nematodes of Hungary (Nematoda errantia). In: *Paedozoologica Hungarica* (Eds. Czudi, C. & Mahunka, S) vol.II, 496p.
- ANDRÁSSY, I. (2005c). Free-living nematodes of Hungary (Nematoda errantia). In: *Paedozoologica Hungarica* (Eds. Czudi, C. & Mahunka, S) vol.III, 608p.
- ANTEBI, A., NORRIS, C.R., HEDGECOCK, E.M. & GARRIGA, G. (1997). Cell and Growth Cone Migrations. In *C. elegans II* (Eds. Riddle, D.L., Blumenthal, T., Meyer, B. & Priess, J.). Chap. 21. pp. 583-609. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
- ARISTOTLE. (384-322 B. C.). 1910 *Historia Animalium*. Translated by D' Arey Wentworth Thompson.
- ARTIGAS, P. (1927). Nematodes de invertebrados. V. *Cruzinema cruzinema* novo genero e nova especie. *Boletim Biológico Laboratório de Parasitologia Faculdade de Medicina de São Paulo* 10, 209-211.
- ARTIGAS, P. (1929). Uma nova especie de nematoideo do genero *Cephalobium* Cobb, 1920. *Boletim Biológico, São Paulo Brazil* 4, 81-84.
- BAJAJ H.K. & DALAL, M.R. (1977). Life cycle of *Verutus mesoangustus* Minagawa Nematoda: Heteroderidae) on *Vetiveria zizanioides* (L.) Nash. (Gramineae). *Fundamental and applied Nematology* 20, 191-196.
- BAKER, A.D. (1962). Checklists of nematode superfamilies Dorylaimoidea, Rhabditoidea, Tylenchoidea, and Aphelenchoidea. Leiden, the Netherlands, Brill, 261 p.
- BALDWIN, J.G., DE LEY, I. T., MUNDO-OCAMPO, M., DE LEY, P., NADLER, S.A. & GEBRE, M. (2002). *Acromoldavicus mojaviensis* n. sp. (Nematoda: Cephalobidae) from the Mojave Desert, California. *Nematology* 3, 343-353.
- BASTIAN, H.C. (1865). Monograph on the Anguillulidae, or free Nematoids, marine, land and freshwater; with descriptions of 100 new species. *Transactions of the Linnaean Society of London* 25, 73-184.
- BEAMS, H.W. & SEKHON, S.S. (1972). Cytodifferentiation during spermiogenesis in *Rhabditis pellio*. *Journal of Ultrastructure Research*.38, 511-527.

- BELOGUROV, O.I. (1977). *Marispelodera stasileonovi* gen. et sp. n., (Nematoda, Rhabditidae) s litorali Komadorskikh ostrovov. [*Marispelodera stasileonovi* gen. et sp. n., Nematoda, Rhabditidae, from the littoral zone of the Commander Islands]. *Zoologicheskii Zhurnal* 56, 813-821 (In Russian).
- BELOGUROV, O.I., MUCHINA, T.I. & CHURIKOVA, N.I. (1977). *Pelodera comandorica* sp. n. (Nematoda, Rhabditidae) from the littoral zone of the commander Islands. *Zoologicheskii Zhurnal* 56, 996-1003.
- BERNARD, E.C. (1979). New species of Bunonematoidea (Rhabditida) from Georgia and Tennessee. *Journal of Nematology* 11, 344-358.
- BERT, W., CLAEYS, M. & BORGONIE, G. (2006). The Comparative Cellular Architecture of the Female Gonoduct Among Tylenchoidea (Nematoda: Tylenchina). *Journal of Nematology* 38, 362-375.
- BERT, W., KARSSSEN, G., VAN DRIESSCHE, R. & GERAERT, E. (2002) The cellular structure of the female reproductive system within the Heteroderinae and Meloidogyninae (Nematoda). *Nematology* 4, 953-964.
- BERT, W., VAN GANSBEKE, R., VANDRIESCHE, R., CLAEYS, M., GERAERT, E. & BORGONIE, G. (2003). Comparative study of the female reproductive system within the Tylenchomorpha (Nematoda: Tylenchina). *Russian Journal of Nematology* 14, 179-184.
- BERT, W., VANGESTEL, S., HOUTHOOFD, W., VAN GANSBEKE, R. & BORGONIE. (2007). The somatic female gonad of Cephalobidae (Nematoda): cellular architecture and associated function. *Nematology* 9, 285-297.
- BIRD, A. F., DE LEY, P. & BIRD, J. (1994) Morphology, oviposition and embryogenesis in an Australian population of *Acrobeloides nanus*. *Journal of Nematology* 25, 607- 615.
- BLAXTER, M. L., DE LEY, P., GAREY, J. R., LIU, L. X., SCHELDEMAN, P., VIERSTRAETE, A., VANFLETEREN, J. R., MACKEY, L. Y., DORRIS, M., FRISSE, L. M., VIDA, J. T. & THOMAS, W. K. (1998). A molecular evolutionary framework for the phylum Nematoda. *Nature* 392, 71-75.
- BLELLOCH, R., ANNA-ARRIOLA, S. S., GAO, D., LI, Y., HODGKIN, J. & KIMBLE, J. (1999). The *gon-1* Gene Is Required for Gonadal Morphogenesis in *Caenorhabditis elegans*. *Developmental Biology* 216, 382-393.
- BLINOVA, S.L. & VOSILITE, B.S. (1976). [*Paramonoviola rhagii* gen. et sp. n. (Rhabditida, Diplogasteridae), a parasite of *Rhagiuminquisitor* L.] (In Russian). *Zoologicheskii Zhurnal* 55, 131-133.
- BLINOVA, S.L. & VOSILITE, B.S. (1978). Ekologomorfologicheskoe izuchenie nematody *Mikoletzkyia sexdentati* n. sp. (Dioplogasteridae) I taksonomicheskij analiz roda *Mikoletzkyia* Weingaertner, 1955. *Trudy Gelmintologicheskoi Laboratorii, Akademia Nauk SSSR (Moskva)* 28, 58-67.
- BLINOVA, S.L. (1982). Entomopathogennye nematody-parazity verditelei lesa Semeistvo Rrabditidy. Izd-vo "Nauka" (Moskva) 132 p (In Russian).
- BORELLUS, P. (1656). 'Observatiouum Micioscopicarum Cenluria,' Hagoe conii.is.384, Paris.

- BORODITSKY, J.M. & SAMOILOFF, M.R. (1973). Effects of growth inhibitors on development of the reproductive system of the free living nematode *Panagrellus redivivus* (Cephalobidae). *Canadian Journal of Zoology* 51, 483-492.
- BOSSINGER, O. & SCHIERENBERG, E. (1992). Cell-cell communication in the embryo of *Caenorhabditis elegans*. *Developmental Biology* 151, 401-409.
- BOSSINGER, O. & SCHIERENBERG, E. (1996). Early embryonic induction in *C. elegans* can be inhibited with polysulfated hydrocarbon dyes. *Developmental Biology* 176, 17-21.
- BOSTRÖM, S. (1984a). A scanning electron microscope study of three species of *Eucephalobus* Steiner, 1936 (Nematoda: Cephalobidae). *Nematologica* 30, 131-139.
- BOSTRÖM, S. (1984b). Description and morphological variability of *Chiloplacus minimus* (Thorne, 1925) Andrassy, 1959 (Nematoda: Cephalobidae). *Nematologica* 30, 151- 160.
- BOSTRÖM, S. (1985a). A new species of *Zeldia* Thorne (Nematoda: Cephalobidae) from Tunisia. *Nematologia Mediterranea* 13, 67-71.
- BOSTRÖM, S. (1985b). *Acrobeles oosiensis* sp. n. (Nematoda: Cephalobidae) from Tunisia. *Nematologia Mediterranea* 13, 123-126.
- BOSTRÖM, S. (1985c). Aspects on the ultrastructure of the cephalic region in two species of Acrobelinae (Nematoda: Cephalobidae). *Acta Zoologica* 66, 211- 216.
- BOSTRÖM, S. (1985d). Description of *Acrobeloides emarginatus* (de Man, 1880) Thorne, 1937 and proposal of *Acrolobus* n. gen. (Nematoda: Cephalobidae). *Revue de Nématologie* 8, 335-340.
- BOSTRÖM, S. (1985e). Descriptions of *Cervidellus serratus* (Thorne, 1925) Thorne, 1937 from Sweden and *C. neftasiensis* n.sp. from Tunisia, with a note on some recently proposed genera (Nematoda: Cephalobidae). *Nematologica* 31, 365- 374.
- BOSTRÖM, S. (1986). Evolution of lip structure in the Cephalobidae (Rhabditida). *Revue de Nématologie* 9, 290.
- BOSTRÖM, S. (1987). A new terrestrial nematode species (Rhabditida: Cephalobidae) from Spitzbergen. *Polar Biology* 7, 375-378.
- BOSTRÖM, S. (1988a). Morphological and Systematic studies of the family Cephalobidae (Nematoda: Rhabtida). Thesis, University of Stokholm, Sweden 34 pp. ISBN 918727207- 5.
- BOSTRÖM, S. (1988b). A scanning electron microscope study of some species of terrestrial nematodes from Spitzbergen. *Nematologica* 33, 366-374.
- BOSTRÖM, S. (1990). Some species of Cephalobidae (Nematoda: Rhabditida) from highland Kenya. *Journal of African Zoology* 104, 127-134.
- BOSTRÖM, S. (1991). Some Cephalobidae (Nematoda: Rhabditida) from sand on the Island of Samos, Greece. *Nematologica* 37, 367-375.
- BOSTRÖM, S. (1992). Some Cephalobidae (Nematoda: Rhabditida) from Crete, Greece. *Fundamental and Applied Nematology* 15, 289- 295.
- BOSTRÖM, S. (1993). Some cephalobids from Ireland and Malaysia (Nematoda: Rhabditida). *Afro Asian journal of Nematology* 3, 128- 134.

- BOSTRÖM, S. (1993). Some cephalobids from Turkey (Nematoda. Rhabditida). *Nematologica Mediterranea*, 21, 295-300.
- BOSTRÖM, S. (1995). Population of *Plectus acuminatus* Bastian, 1865 and *Panagrolaimus magnivulvatus* n. sp. (Nematoda) from nunatakks in Dronning Maud Land, East Antarctica. *Fundamental and Applied Nematology* 18, 25-34.
- BOSTRÖM, S. (1997). *Chiloplectus masleni* sp. nov. and variability in populations of *Plectus acuminatus* Bastian, 1865 (Nematoda: Plectidae) from the naunatak Basen, Vestfgella, Dronning Maud Land, East Antarctica. *Polar Biology* 17, 74-80.
- BOSTRÖM, S. (1998). A scanning electron microscope study of some species of terrestrial nematodes from Spitzbergen. *Nematologica* 33, 366-374.
- BOSTRÖM, S. (2000). A divergent population of *Cervidellus capraeolus* (De Ley, Geraert & Coomans, 1990) Boström & De Ley, 1996 (Nematoda: Cephalobidae) from the Bahamas. *Journal of Nematode Morphology and Systematics* 3, 95-99.
- BOVERI, T. (1887). "Die Bildung der Richtungskörper bei *Ascaris megalcephala* und *Ascaris lumbricoides*." *Jenaische Zeitschrift für Naturwissenschaft* 21, 423-515
- BOVERI, T. (1892). Über die Entstehung des Gegensatzes zwischen den Geschlechtszellen und die somatischen Zellen bei *Ascaris megalcephala*. *Sitzungsberichte der Gesellschaft Morphologische und Physiologische* 8, 114-125.
- BOVIEN, P. (1937). Some type of association between nematodes and insects. *Videnskabelige Meddelelser fra Dansk Naturhistorik Forening* 101, 1-114.
- BRAUCHLE, M., KIONTKE, K., MACMENAMIN, P., FITCH, D.H.A. & PIANO, F. (2009). Evolution of early embryogenesis in rhabditid nematodes. *Developmental Biology* 335, 253-262.
- BRENNER, S. (1974). The genetics of *Caenorhabditis elegans*. *Genetics* 77, 71-94.
- BRUN, J. & CAYROL, J. C. (1970). Le développement embryonnaire du nématode mycophage *Ditylenchus myceliophagus* Goodey, 1958 (Nematoda: Tylenchida). *Nematologica* 16, 523-531.
- BRZESKI, M.W. (1960). *Cephalobus (Heterocephalobus) kaezanowskii* subgen. nov., sp. nov. (Nematoda: Cephalobidae). *Bulletin de l'Académie Polonaise des Sciences. ClassII, Série des Sciences Biologique* 8, 153-165.
- BRZESKI, M.W. (1961). Revision of the genus *Heterocephalobus* Brzeski, 1960 n. grad. (Nematoda, Cephalobidae). *Bulletin de l'Académie Polonaise des Sciences. ClassII, Série des Sciences Biologique* 9, 97-100.
- BRZESKI, M.W. (1962). Three new species of *Acrobeloides* Cobb (Nematoda, Cephalobidae). *Bulletin de l'Académie Polonaise des Sciences. ClassII, Série des Sciences Biologique* 10, 335-339.
- BRZESKI, M.W. (1963). Review of the nematode genus *Anaplectus* De Coninck & Schuurmans Stekhoven (Nematoda, Plectidae). *Bulletin de l'Académie Polonaise des Sciences. ClassII, Série des Sciences Biologique* 11, 35-38.
- BRZESKI, M.W. (1989). *Cruznema velatum* sp. n. and observations on *C. tripartitum* (Linstow) (Nematoda: Rhabditidae). *Annales Zoologici* 43, 71-75.

- BRZESKI, M.W. (1993). *Geomonhystera breviseta* sp. n., *G. mexicana* sp. n. and observation on *G. australis* (Cobb, 1893) Andrassy, 1981 (Nematoda: Monhysterida). *Fundamental and Applied Nematology* 16, 107-114.
- BRZESKI, M.W. (1997). Studies on some species of the genus *Prismatolaimus* de Man, 1880 with description of *P. mulcoomus* sp. n. (Nematoda: Prismatolaimidae). *Annales Zoologici* (Warszawa) 46, 153-166.
- BÜTSCHLI, O. (1873). Beiträge zur Kenntnis der freilebenden Nematoden. *Nov Acta der Kaiserlich-Leopoldinisch-carolinischen Deutschen Akademie der Naturforscher* 36, 3-144.
- BÜTSCHLI, O. (1874). Zur Kenntnis der freilebenden Nematoden, insbesondere der des Kieler Hafens. *Abhandlungen der Senckenbergischen naturforschenden Gesellschaft* 9, 1-56.
- BÜTSCHLI, O. (1875). Vorläufige Mittheilung über Untersuchungen betreffend die ersten Entwicklungsvorgänge im befruchteten Ei von Nematoden und Schnecken. *Zeitschrift für wissenschaftliche Zoologie* 25, 201-213.
- BÜTSCHLI, O. (1876). Untersuchungen über freilebende Nematoden und die Gattung *Chaetonotus*. *Zeitschrift für wissenschaftliche Zoologie* 26, 363-413.
- BYERLY, L., CASSADA, R.C. & RUSSELL, R.L. (1976). The life cycle of the nematode *Caenorhabditis elegans*. *Developmental Biology* 51, 23-33.
- CHIN, D. A. (1977). Embryonic and post-embryonic development of *Cylindrocorpus longistoma* (Nematoda: Cylindrocorporidae). *Nematologica* 23, 62-70.
- CHANG, C., NEWMAN, A. P., & STERNBERG, P. W. (1999). Reciprocal EGF signaling back to the uterus from the induced *C. elegans* vulva coordinates morphogenesis of epithelia. *Current Biology* 9, 237-246.
- CHIN, D. A. & TAYLOR, D.P. (1969a). Sexual attraction and mating patterns in *Cylindrocorpus longistoma* and *C. curzii* (Nematoda: Cylindrocorporidae). *Journal of Nematology* 1, 313-317.
- CHIN, D.A. & TAYLOR, D.P. (1969b). Gametogenesis and chromosome complement in *Cylindrocorpus longistoma* and *C. curzii*. *Nematologica* 15, 525-529.
- CHIN, D.A. & TAYLOR, D.P. (1970). Observations on molting in species of *Cylindrocorpus*. *Nematologica* 16, 1-5.
- CHISHOLM, A. D. (2005). Epidermal morphogenesis (December 01, 2005). *WormBook*, (Ed). The *C. elegans* Research Community, WormBook, doi/10.1895/wormbook.1.35.1, <http://www.wormbook.org>.
- CHITWOOD, B.G. (1930). Studies on some physiological functions and morphological characters of *Rhabditis* (Rhabditidae, Nematodes). *Journal of Morphology and Physiology* 49, 251-274.
- CHITWOOD, B.G. (1933). A revised classification of the Nematoda. *Journal of Parasitology* 20, 131.
- CHITWOOD, B.G. (1937). A revised classification of the Nematoda. *Papers on Helminthology in Commemoration of the 30-Year Jubileum of K.I. Skrjabin and of the All-Union Institute of Helminthology*, Moscow. 69-80.
- CHITWOOD, B.G. (1958). The designation of official names for higher taxa of invertebrates. *Bulletin of Zoological Nomenclature* 15, 660-695.

- CHITWOOD, B.G. & CHITWOOD, M.B. (1950). *Introduction to Nematology*, Monumental Printing Company, Baltimore, revised edition, 213.
- CHITWOOD, B.G. & MCINTOSH, A. (1934). A new variety of *Alloionema* (Nematoda: Diplogasteridae), with a note on the genus. *Proceedings of the Helminthological Society of Washington* 1, 37-38.
- CHIU C.T., BALDWIN, J.G & MUNDO-OCAMPO, M. (2002). *Metacrobeles bradyurus* n. sp. (Nematoda: Cephaloboidea) from Death Valley, California. *Nematology* 4, 645- 652.
- CHONGTI, T. (1982). Studies on the plant-nematodes in South Fujian. II. The species of Rhabditida. *Acta Zoologica Sinica* 28, 157-164.
- CHUANG, S.H. (1962). The embryonic and post-embryonic development of *Rhabditis teres* (A. Schneider). *Nematologica* 7, 317-330.
- CLARK, S.A. (1967). The development and life history of the false root-knot nematode, *Nacobbus serendipiticus*. *Nematologica* 13, 91-101.
- CLAUS, C.F.W. (1862). Über einige im Humuslebende Anguillulinen. *Zeitschrift für Wissenschaftliche Zoologie* 12, 354-359.
- COBB, N.A. (1893). Nematodes, mostly Australian and Fijian. *Macleay New South Wales Miscellaneous Publications (Sydney)* 13, 3-59.
- COBB, N.A. (1898). Extract from MS report on the parasites of stock. *Agricultural Gazette New South Wales* 9, 269-321.
- COBB, N.A. (1914). The North American free-living fresh-water nematodes. *Transactions of American Microscopical Society* 33, 69-134.
- COBB, N.A. (1918). Estimating the nema population of the soil. U.S. Department of Agriculture. *Agricultural Technical Circular of US Department of Agriculture* 1, 48p.
- COBB, N.A. (1924). *Neodiplogaster tropica* n. g. n. sp. *Journal of Parasitology* 11, 105p.
- COBB, N.A. (1929). Observations on the morphology and physiology of nemas; including notes on new species. *Journal of Washington Academy of Science* 19, 283-286.
- CONRADT, B. & XUE, D. (2005). Programmed cell death (October 06, 2005). *WormBook*. (Ed. The C. elegans Research Community, WormBook) doi/10.1895/wormbook.1.32.1, <http://www.wormbook.org>.
- COOMANS, A. (1962). Systematisch-ecologisch onderzoek van de vrijlevende bodemnematoden in België. De vrijlevende nematoden-fauna van weideland, I. *Natuurwetenschappelijk Tijdschrift* 43, 87-132.
- COOMANS, A. (1971). Morphology and nematode systematics. *Journal of Parasitology* 57, 95-99.
- COOMANS, A. & DE WAELE, D. (1979). Species of *Aphanolaimus* (Nematoda, Araeolaimida) from Africa. *Zoologica Scripta* 8, 171-180.
- COOMANS, A. & DE WAELE, D. (1983). Species of *Aphanolaimus* (Nematoda: Araeolaimida) from Belgium. *Hydrobiologia* 101, 165-178.

- COOMANS, A. & GOODEY, J.B. (1965). *Drilocephalobus congoensis* n. g., n. sp. *Nematologica* 11, 116-120.
- COOMANS, A. & HEYNS, J. (1986). *Oncholaimus jessicae* n. sp. (Nematoda: Oncholaimidae) from fresh water in the Transvaal. *South African Journal of Zoology* 21, 197-201.
- COOMANS, A. & JACOBS, L.J. (1983). *Halalaimus algeriensis* n. sp. from the Sahara. *Hydrobiologia* 102, 39-44.
- COOMANS, A. & LIMA, M.B. (1965). Description of *Anatonchus amiciae* n. sp. (Nematoda: Mononchidae) with observations on its juvenile stages and anatomy. *Nematologica* 11, 413-431.
- COOMANS, A. & MULK, M.M. (1980). *Prismatolaimus andrassyanus* nomen novum for *P. andrassyi* apud Mulk & Coomans, 1979, nec *P. andrassyi* Khera & Chaturvedi, 1977. *Nematologica* 26, 283.
- COOMANS, A. & RASKI, D.J. (1988). Two new species of *Prismatolaimus* de Man, 1880 (Nemata: Prismatolaimidae) in Southern Chile. *Journal of Nematology* 20, 288-303.
- COOMANS, A. & RASKI, D.J. (1991). *Aphanonchus intermedius* n. g., n. sp. (Nemata: Araeolaimida). *Nematologica* 37, 8-19.
- COWAN CR & HYMAN AA. (2004). Asymmetric cell division in *C. elegans*: cortical polarity and spindle positioning. *Annual Review of Cell and Developmental Biology* 20, 427-53.
- DASGUPTA, D.R., NAND, S. & SESHADRI, A.R. (1969). Embryology and life history studies on the lance nematode, *Hoplolaimus indicus* Sher. (Abstract) All India Nematology Symposium, New Delhi, August 21-22, p. 31.
- DASGUPTA, D.R., NAND, S. & SESHADRI, A.R. (1970). Culturing, embryology and life history studies on the lance nematode, *Hoplolaimus indicus*. *Nematologica* 16, 235-248.
- DASSONVILLE, A.F. & HEYNS, J. (1984). Freshwater nematodes from South Africa. 7. New and known species collected in Skannerspruit, Pretoria. *Phytophylactica* 16, 15-32.
- DE CONINCK, L.A.P. & SCHUURMANS STEKHOVEN, J.H. (1933). The free-living marine nemas of the Belgian Coast. II, with general remarks on the structure and the systems of nemas. *Memoires Musée Royal d' Histoire Naturelle de Belgique* 58, 1-163.
- DE LEY, P. (1997). The current state of affairs in identification and diagnosis of the genera of the family Cephalobidae (Nematoda: Rhabditida). *Mededelingen Faculteit Landbouwkundige en Toegepaste Biologische Wetenschappen Universiteit Gent* 62, 657-673.
- DE LEY, P. & BLAXTER, M. (2002). Systematic position and phylogeny. In: *The Biology of Nematodes*, (Ed Lee, D.L.). Taylor and Francis, London, New York. pp1-30.
- DE LEY, P., COOMANS, A. & GERAERT, E. (1990a). *Metacrobeles tessellatus* sp. n., second species of a rare genus (Nematoda: Rhabditida). *Nematologica* 35, 25-36.
- DE LEY, P., GERAERT, E. & COOMANS, A. (1990b). Seven cephalobids from Senegal (Nematoda: Rhabditida). *Journal of African Zoology* 104, 287-304.

- DE LEY, P., HERNANDEZ, M.A. & AGUDO, J. (1994). A redescription of *Ypsylonellus similis* (Thorne, 1925) Andrassy, 1984, with descriptions of some related species (Nematoda: Cephalobidae). *Fundamental and Applied Nematology* 17, 1-16.
- DE LEY, P., & SIDDIQI, M.R. (1991). Description of *Pseudacrobeles macrocystis* sp. n., with some new observations on the morphology of Cephalobidae (Nematoda). *Afro-Asian Journal of Nematology* 1, 31-40.
- DE LEY, P., SIDDIQI, M.R. & BOSTRÖM, S. (1993a). A revision of the genus *Pseudacrobeles* Steiner, 1938 (Nematoda: Cephalobidae). Part 1. Subgenus *Pseudacrobeles* grad. n. *Fundamental and Applied Nematology* 16, 219-238.
- DE LEY, P., SIDDIQI, M.R. & BOSTRÖM, S. (1993b). A revision of the genus *Pseudacrobeles* Steiner, 1938 (Nematoda: Cephalobidae). Part 2. Subgenus *Bunobus* subg. n. problematical species, discussion and key. *Fundamental and Applied Nematology* 16, 289-308.
- DE LEY, P., & VANDEBROEK, I. (1992). *Stegelletina laticollaris* n. sp., a second new species of the genus from Senegal (Nematoda: Cephalobidae). *Belgian Journal of Zoology* 122, 139-146.
- DE LEY, P., VAN DE VELDE, M. C., MOUNPORT, D., BAUJARD, P. & COOMANS, A. (1995). Ultrastructure of the stoma in Cephalobidae, Panagrolaimidae and Rhabditidae, with a proposal for a revised stoma terminology in Rhabditida (Nematoda). *Nematologica* 41, 153-182.
- DE MAESENEER, J. & D' HERDE, J. (1963). Méthodes utilisées pour l'étude des anguillules libres du sol. *Revue de Agriculture Bureaux* 16, 441-447.
- DE MAN, J.G. (1876). Onderzoekingen over vrij in de aarde levende Nematoden. *Tijdschrift der Nederlansche Dierkundige Vereeniging* 2, 78-196.
- DE MAN, J.G. (1880). Die einheimischen, frei in der reinen Erde und im süßen Wasser lebenden Nematoden. Vorläufiger Bericht und descriptivsystematischer Theil. *Tijdschrift der Nederlansche Dierkundige Vereeniging* 5, 1-104.
- DE MAN, J.G. (1884). Die frei in der reinen Erde und im süßen Wasser lebenden Nematoden der niederländischen Fauna. *Eine systematisch-faunistische Monographie*, Leiden, 1-206.
- DE MAN, J.G. (1895). Description of three species of Anguillulidae observed in diseased pseudobulbs of tropical orchids. *Proceedings and Transactions of the Liverpool Biological Society* 9, 76-94.
- DE MAN, J.G. (1912). *Odontopharynx longicaudata* n. g., n. sp., eine neue Form von Anguilluliden. *Zoologische Jahrbücher (Systematik)* 33, 637-642.
- DE MAN, J.G. (1920). Description d'une espèce nouvelle du genre *Diplogaster* Max Schultze: *Diplogaster consobrinus*, nov. sp. *Annales de la Société Royale Zoologique et Malacologique de Belgique* 51, 47-54.
- DE MAN, J.G. (1927). Das Männchen der *Rhabditis monhystera* Bütschli. *Zoologischer Anzeiger* 70, 51-57.
- DELATTRE, M. & FELIX, M.A. (1999). Connection of vulval and uterine epithelia in *Caenorhabditis elegans*. *Biology of the Cell* 91, 573-583.
- DELATTRE, M. & FELIX, M.A. (2001). Polymorphism and evolution of vulval precursor cell lineages within two nematode genera, *Caenorhabditis* and *Oscheius*. *Current Biology* 11, 631-643.

- DEMEURE, V., NETSCHER, C. & QUENEHERVE, P. (1980). Biology of plant parasitic nematode *Scutellonema cavenessi* Sher, 1964, reproduction and development and life cycle. *Revue de Nematologie* 3, 213-225.
- DEPPE, U., SCHIERENBERG, E., COLE, T., KRIEG, C., SCHMITT, D., YODER, B. & VON EHRENSTEIN, G. (1978). Cell lineages of the embryo of the nematode *Caenorhabditis elegans*. *Proceedings of National Academy of Sciences USA*, 75, 376-380.
- DEUBERT, K.H. & ZUCKERMAN, B.M. (1967). Development of *Panagrellus redivivus* (Nematoda: Rhabditidae) in chemically defined media. *Experimental Parasitology* 21, 209-214.
- DIAZ, S.A., HAYDON, D.T. & LINDSTRÖM, J. (2010). Sperm-limited fecundity and polyandry-induced mortality in female nematodes *Caenorhabditis remanei*. *Biological Journal of the Linnean Society* 99, 362-369.
- DOLINSKI, C., BALDWIN, J.G. & THOMAS, W.K. (2001). Comparative survey of early embryogenesis of Secernentea (Nematoda), with phylogenetic implications. *Canadian Journal of Zoology* 79, 82-94.
- DOLLIVER, J.S., HILDEBRANDT, A.C. & RIBER, A.J. (1962). Studies of reproduction of *Aphelenchoides ritzemabosi* (Schwartz) on plant tissues in culture. *Nematologica* 7, 294-300.
- DONCASTER, C.C. & SEYMOUR, M.K. (1973). Egg-transport in the genital tract and egg laying of *Aphelenchoides blastophthorus*. *Nematologica* 19, 379-388.
- DOUGHERTY, E.C. (1953). The genera of the subfamily Rhabditinae Micoletzky, 1922 (Nematoda). *Thaper Commemoration Volume*, 69-72.
- DOUGHERTY, E.C. (1955). The genera and species of the subfamily Rhabditina Micoletzky, 1922 (Nematoda). A nomenclatorial analysis- including an addendum on the composition of the family Rhabditidae Örley, 1880. *Journal of Helminthology* 29, 105-152.
- DOUGHERTY, E.C., HANSEN, E.L., NICHOLAS, W.L., MOLLET, J.A. & YARWOOD, E.A. (1959). Axenic cultivation of *Caenorhabditis briggsae* (Nematoda: Rhabditidae) with supplemented and unsupplemented chemically defined media. *Annals of the New York Academy of Sciences* 77, 176-217.
- DUGGAL, C.L. (1978). Copulatory behaviour of male *Panagrellus redivivus*. *Nematologica* 24, 257-268.
- DUJARDIN, F. (1845). *Histoire naturelle des helminthes ou vers intestinaux*. Paris. 654 p.
- EBSARY, B.A. (1986a). Description of *Butlerius canadensis* n. sp. (Nematoda: Diplogasteridae) with an emendation of the genus and a proposal for *Parabutlerius* n. gen. *Canadian Journal of Zoology* 64, 1782-1785.
- EBSARY, B.A. (1986b). *Mononchoides andersoni* n. sp. and two new species of *Koerneria* (Nematoda: Diplogasteridae). *Canadian Journal of Zoology* 64, 2012-2020.
- EHRENSTEIN, G. VON & SCHIERENBERG, E. (1980). Cell lineages and development of *Caenorhabditis elegans* and other nematodes. In '*Nematodes as Biological Models*' vol. I Behavioural and Developmental models. (Ed. B. M. Zuckerman) Academic Press, New York and London, pp 1-17.

- EIZINGER, A., JUNGBLUT, B. & SOMMER, R.J. (1999). Evolutionary change in the functional specificity of genes. *Trends in Genetics* 15, 197-202.
- EIZINGER, A. & SOMMER, R.J. (1997). The homeotic gene *lin-39* and the evolution of nematode epidermal cell fates. *Science* 278, 353–540.
- EMMONS, S.W. (2005). Male Development (November 10, 2005). *WormBook*. (Ed. The *C. elegans* Research Community), WormBook, doi/10.1895/wormbook.1.33.1,
- EROSHENKO, A.S. (2002). A new species of the genus *Cuticularia* (Nematoda: Rhabditida) from island of the Siam Bay. *Zoologicheskii Zhurnal* 81, 628-631.
- EVANS, A.A.F. & FISHER, J.M. (1969). Development and structure of populations of *Ditylenchus myceliophagus* as affected by temperature. *Nematologica* 15, 395-402.
- EYUALEM, A., TRAUNSPURGER, W. & ANDRÁSSY, I. (2006). *Freshwater nematodes: Ecology and Taxonomy* CABI Publishing, Wallingford, Oxfordshire OX108DE, UK, 752p.
- FARKAS, K. (1973). *Teratorhabditis mariannae* n. sp., eine neue Nematodenart aus Champignonkulturen. *Opuscula Zoologica* 11, 61-64.
- FÉLIX, M.A. (2006). *Oscheius tipulae* (August 16, 2006). *WormBook*. (Ed. The *C. elegans* Research Community), WormBook, doi/10.1895/wormbook.1.119.1, <http://www.wormbook.org>.
- FÉLIX, M.A., DE LEY, P., SOMMER, R.J, FRISSE, L.M., NADLER, S.A., THOMAS, W.K., VANFLETEREN, J. R., & STERNBERG, P. W. (2000). Evolution of vulva development in the Cephalobina (nematoda). *Developmental Biology* 221, 68–86.
- FÉLIX, M.A. & STERNBERG, P.W (1996). Symmetry breakage in the development of one-armed gonads in nematodes. *Development* 122, 2129 -2142.
- FÉLIX, M.A. & STERNBERG, P.W. (1997). Two nested gonadal inductions of the vulva in nematodes. *Development* 124, 253–259.
- FILIPJEV, I. (1918). Free-living marine nematodes of the Sevastopol area. *Trudy Osoboï Zoologicheskoi laboratorii i Sebastopol'skoi biologicheskoi stantsii* 4, 1-350.
- FILIPJEV, I. (1921). Free-living marine nematodes of the Sevastopol area. *Trudy Osoboï Zoologicheskoi laboratorii i Sebastopol'skoi biologicheskoi stantsii* 4, 351-614.
- FILIPJEV, I. (1934). The classification of the free-living nematodes and their relation to the parasitic nematodes. *Smithsonian Miscellaneous Collections (Washington)* 89, 1-63.
- FILIPJEV, I. & SCHUURMANS STEKHOVEN, J.H. (1941). *A manual of Agricultural Helminthology*. Leiden, 878 p.
- FITCH, D.H.A. (2000). Evolution of “Rhabditidae” and the male tail. *Journal of Nematology* 32, 235-244.
- FLEGG, J.J.M. (1966). Once yearly reproduction in *Xiphinema vuittenzi*. *Nature, London*. 212, 741.
- FLEGG, J.J.M. (1968). Embryogenic studies of some *Xiphinema* and *Longidorus* species. *Nematologica*, 14, 137-145.

- FRECKMAN, D.W. (1988). Bacteriovores nematodes and organic matter decomposition. *Agriculture, Ecosystems and Environment* 24, 195-217.
- FUCHS, A.G. (1915). Die Naturgeschichte der Nematoden und einiger anderer Parasiten. 1. Des *Ips typographus* L.2. Des *Hylobius abietis* L. *Zoologische Jahrbücher (Systematik)* 38, 109-222.
- FUCHS, A.G. (1930). Neue an Borken-und Rüsselkäfer gebundene Nematoden, halbparasitische und Wohnungseinmieter. *Zoologische Jahrbücher (Systematik)* 59, 505-646.
- FUCHS, A.G. (1931). Einige neue *Rhabditis*-Arten. *Zoologische Jahrbücher (Systematik)* 62, 119-148.
- FÜCHS, G. (1937). Neue parasitische und halbparasitische Nematoden bei Borkenkäfern und einige andere Nematoden. 1. Teil. *Zoologische Jahrbücher (Systematik)* 70, 291-380.
- FUCHS, A.G. (1938). Neue parasiten und halbparasiten bei Borkenkäfern und einige andere Nematoden. *Zoologische Jahrbücher (Systematik)* 71, 123-190.
- FÜRST VON LIEVEN, A. & SUDHAUS, W. (2000). Comparative and functional morphology of the buccal cavity of Diplogastrina (Nematoda) and a first outline of phylogeny of this taxon. *Journal of Zoological Systematics and Evolutionary Research* 38, 37-63.
- FÜRST VON LIEVEN, A. (2002a). Functional morphology, origin and phylogenetic implications of the feeding mechanism of *Tylopharynx foetida* (Nematoda: Diplogastrina). *Russian Journal of Nematology* 10, 11-23.
- FÜRST VON LIEVEN, A. (2002b). The sister group of the Diplogastrina (Nematoda). *Russian Journal of Nematology* 10, 127-137.
- FÜRST VON LIEVEN, A. (2003). The genus *Oigolaimella* Paramonov, 1952 (Nematoda: Diplogastridae) and a description of *Oigolaimella kruegeri* n. sp. and *O. ninae* n. sp. *Nematology* 5, 583-600.
- FÜRST VON LIEVEN, A. (2005). The embryonic moult in diplogastrids. (Nematoda) – homology of developmental stages and heterochrony as a prerequisite for morphological diversity. *Zoologischer Anzeiger* 244, 79-91.
- FÜRST VON LIEVEN, A. (2008a). *Koerneria sudhausi* n. sp. (Nematoda: Diplogastridae); a hermaphroditic diplogastrid with an egg shell formed by zygote and uterine components. *Nematology*, 1, 27-45.
- FÜRST VON LIEVEN, A. (2008b). Functional morphology of spicules and the role of the male gonoduct secretion in the diplogastrid nematode, *Diplogastriana schneideri*. *Journal of Nematode Morphology and Systematics* 10, 177-185.
- GADEA, E. (1972). Some aspects on phylogeny of Nematoda. *Proceedings of International Symposium (11th) on Nematology* (European Society of Nematologists), Abstracts, 23.
- GADEA, E. (1973). Sobre la filogenia interna des los nematodos. *Publicaciones del Instituto de Biología Aplicada, (Barcelona)* 54, 87- 92.
- GAGARIN, V.G. (1977). [Three new species of nematodes from the order Rhabditida from areas of sewage discharge]. (In Russian). *Zoologicheskii Zhurnal* 56, 1245-1248.
- GAGARIN, V.G. (1979). Two new species of free-living fresh-water nematodes. (In Russian). *Zoologicheskii Zhurnal* 58, 596-598.

- GAGARIN, V.G. (1981) Freshwater Nematodes of the European Part. Nauka, Leningrad, Russia, 249 pp.
- GAGARIN, V.G. (1983). [New species of the genus *Mononchoides* (Nematoda, Diplogasterinae)]. (In Russian). *Zoologicheskii Zhurnal* 62, 25-31.
- GAGARIN, V.G. (1987). Deux nouvelles espèces de la famille Monhysteridae (Nematoda). Two species of family Monhysteridae (Nematoda). *Zoologičeskii Zhurnal* 66, 454-556.
- GAGARIN, V.G. (1993). Free-living Nematodes from the Fresh Waters of Russia and Adjacent Countries: Order Monhysterida, Areaolaimida, Chromadorida, Enoplida, Mononchida. Gidrometeoizdat, St. Petersburg, Russia, 352 pp.
- GAGARIN, V.G. (1995). Morphology and taxonomy of the genus *Fictor* Paramonov, 1952 (Rhabditida: Diplogasterida). *Russian Journal of Nematology* 3, 63-64.
- GAGARIN, V.G. (1997). New species of freshwater nematodes of the order Monhysterida from the Novaya Zemlya archipelago (Nematoda). *Zoosystematica Rossica* 6, 21-30.
- GAGARIN, V.G. (1998a). A review of the genus *Mononchoides* Rahm, 1928, with description of two new species from European Russia (Nematoda: Diplogasterida). *Zoosystematica Rossica* 7, 239-241.
- GAGARIN, V.G. (1998b). *Glaxinemella striata* gen. et sp. n. from Europe (Nematoda: Diplogasterida). *Zoosystematica Rossica* 7, 239-241.
- GAGARIN, V.G. (1998c). [Description of *Fictor tumidus* sp. n. (Nematoda, Diplogasterida) from rotten vegetables.] (In Russian). *Zoologicheskii Zhurnal* 77, 1191-1193.
- GAGARIN, V.G. (2000a). [Nematode fauna of manure and compost in Yaroslavl oblast (Russia).] (In Russian.) *Zoologicheskii Zhurnal* 79, 1260-1274.
- GAGARIN, V.G. (2000b). Fauna of free-living nematodes from littoral of Kurilskoe Lake (Russia, Kamchatka Peninsula) and description of *Rhitis eximius* sp. n. (Nematoda, Rhabditida). *Biology of Inland Water* 3.
- GAGARIN, V.G. (2001a). [New species of free-living nematodes from Biwa Lake and inflowing stream (Honshu Island, Japan)]. (In Russian). *Zoologicheskii Zhurnal* 80, 12-25.
- GAGARIN, V.G. (2001b). Free-living Nematodes of the Fresh Waters of Russia and Adjacent Countries Fauna and the Pathways of its Formation, Ecology, Taxonomy, Phylogeny. Nauka Publisher, Moscow, 170 p.
- GAGARIN, V.G. (2002a). A review of the genus *Diplogasteritus* Paramonov, 1952 with description of a new species from Central Russia (Nematoda, Diplogasterida: Diplogasteridae). *Zoosystematica Rossica* 11, 57-73.
- GAGARIN, V.G. (2002b). Description of three new species of *Geomonhystera* (Nematoda: Monhysterida). *Zoosystematica Rossica* 11, 75-84.
- GAGARIN, V.G. (2003). Some new and poorly known species of Tylenchidae and Monhysteridae from Siberia (Nematoda). *Zoosystematica Rossica* 12, 1-6.
- GAGARIN, V.G. (2009). Free-Living Nematodes from a small river in Central Russia. *Inland Water Biology* 2, 25-29.

- GAGARIN, V.G. & NGUYEN VU THANH (2006). Three new species of free-living nematodes of the family Comesomatidae from Mekong River, Vietnam (Nematoda, Monhysterida). *Zoosystematica Rossica* 15, 1-10.
- GAGARIN, V.G. & KLERMAN, A.K. (2007). New species of predatory chromadorids (Nematoda, Chromadorida) from the Mediterranean Sea. *Zoologičeskii Žhurnal* 86, 778-786.
- GAGARIN, V.G. & KUZMIN, L.L. (1972). New data to the taxonomy of the genus *Prismatolaimus* (Nematoda: Onchulidae). *Zoologicheskii Zhurnal*, 51, 1879-1881 (In Russian).
- GAGARIN, V. G. & LEMSINA, L.V. (1982). A male of *Mononchus aquaticus* and two new species of free-living nematodes from Lake Issyk-kul. *Zoologicheskii Zhurnal* 61, 935-937. (In Russian).
- GAGARIN, V.G. & NGUYEN THI THU (2008a). Free-living from the Red River delta. *Inland water biology* 4, 320-325.
- GAGARIN, V.G., & NGUYEN THI THU (2008b). Two new Monhysterid species (Nematoda, Monhysterida) from the mouth of Red River Delta, Vietnam. *Zoologičeskii Zhurnal* 87, 505-510.
- GAGARIN, V.G., & NGUYEN THI THU (2008c). Three new species of the genus *Daptonema* (Nematoda, Xylidae) from the Red River Delta, Vietnam. *Zoologičeskii Zhurnal* 87, 515-523.
- GAGARIN, V.G. & NGUYEN VU THANH (2003). Three new species of free-living nematodes from Vietnamese water bodies. *Zoologičeskii Zhurnal* 82, 1393-1401 (in Russian).
- GAGARIN, V.G. & NGUYEN VU THANH (2007). *Ingenia communis* sp. n., *Adoncholaimus longispiculatus* sp. n. and *Viscosia sedata* sp. n. (Nematoda: Enoplida) from the Mekong River Delta, Vietnam. *International Journal of Nematology* 17, 205-212.
- GAGARIN, V.G. & NGUYEN VU THANH (2008). Four new species of free-living nematodes of family Axonolaimidae (Nematoda, Araeolaimida) from mangrove of Mekong River Delta, Vietnam. *International Journal of Nematology* 18, 133-143.
- GAGARIN, V.G., NGUYEN VU THANH & NGUYEN THI THU (2003). Three new species of free-living nematode from fresh-water bodies of Vietnam (Nematoda: Araeolaimida). *Zoosystematica Rossica* 12, 7-14.
- GAGARIN, V.G., NGUYEN VU THANH & NGUYEN THI THU (2005). *Daptonema salvum* sp. n. and *D. rigidum* sp. n. (Nematoda: Monhysterida) from Chu River, Vietnam. *International Journal of Nematology* 15, 210-214.
- GERAERT, E. (1973). A comparative study of the structure of the female gonads in plants parasitic Tylenchida (Nematoda). *Annales de la Société Royale Zoologique de Belgique* 102, 171-198.
- GERAERT, E. (1976). The female reproductive system in *Deladenus* and *Hexatylus* with a redefinition of the oviduct in the Tylenchida (Nematoda). *Nematologica* 22, 437-445.
- GERAERT, E. (1978). *Enkele nieuwe inzichten in de morfometrie en in de morfologie van de nematoden*. Ghent University, Belgium, Thesis, 259 pp.
- GERAERT, E. (1981). The female reproduction system in nematode systematics. *Annales de la Société Royale Zoologique de Belgique* 110, 73-86.

- GERAERT, E. (1983). The use of the female reproductive system in nematode systematics. In: Stone, A.R., Platt, H.M. & Khalil, L.F. (Eds). *Concepts in nematode systematics*. London, UK, Academic Press, pp. 73-84.
- GERAERT, E., GROOTAERT, P. & DECRAEMER, W. (1980a). Structure of the female reproductive system in some Dorylaimida and Enoplida (Nematoda). *Nematologica* 26, 255-271.
- GERAERT, E., SUDHAUS, W. & GROOTAERT, P. (1980b). The structure of the female genital apparatus in the order Rhabditida (Nematoda). *Annales de la Société Royale Zoologique de Belgique* 109, 91-108.
- GERBER, K. & GIBLIN-DAVIS, R. M. (1990). *Teratorhabditis palmarum* n. sp. (Nematoda: Rhabditidae): An associate of *Rhynchophorus palmarum* and *R. cruentatus*. *Journal of Nematology* 22, 337-347.
- GERLACH, S.A. & RIEMANN, F. (1973). The Bremerhaven checklist of aquatic nematodes. *Veröffentlichungen Institut für Meeresforschung Bremerhaven, Supplement* 4, 1-404.
- GERLACH, S.A. & RIEMANN, F. (1974). The Bremerhaven checklist of aquatic nematodes. *Veröffentlichungen Institut für Meeresforschung Bremerhaven, Supplement* 4, 405-734.
- GIBLIN, R.M. & KAYA, H.K. (1984). *Aduncospiculum halicti* n. gen., n. sp. (Diplogasterida: Diplogasteroididae), an associate of bees in the genus *Halictus* (Hymenoptera: Halictidae). *Revue de Nématologie* 7, 189-197.
- GIBLIN-DAVIS, R.M., YE, W.M., KANZAKI, N., WILLIAMS, D., MORRIS, K. & THOMAS, W.K. (2006). Stomal ultrastructure, molecular phylogeny, and description of *Parasitodiplogaster laevigata* sp. nov. (Nematoda: Diplogasteridae), a parasite of fig wasps. *Journal of Nematology* 38, 137-149.
- GOETTE, A. (1882). Entwicklungsgeschichte der *Rhabditis nigrovenosa*. Untersuch. z. Entwicklungsgeschichte der Würmer. Leipzig.
- GOLDEN, A.M. (1956). Taxonomy of the spiral nematodes (*Rotylenchus* and *Helicotylenchus*) and the developmental stages and host-parasite relationships of *R. buxophilus* n. sp., attacking boxwood. *Maryland Agricultural Experimental Station Bulletin* A- 85, 28p.
- GOLDEN, A.M. (1971). Classification of the genera and higher categories of the order Tylenchida (Nematoda). In: Zuckerman, B.M, Mai, W.F. & Rohde, R.A. (Eds.) *Plant Parasitic Nematodes. Volume 1. Morphology, Anatomy, Taxonomy and Ecology*. Academic Press Inc., New York and London, 191-232.
- GOLDEN, A.M. (1986). Morphology and identification of cyst nematodes. In: *Cyst Nematodes*. (Eds. Lamberti, F. & Taylor, C.E.) Plenum Press, New York 23-45.
- GOLDSTEIN, B., FRISSE, L.M. & THOMAS, W.K. (1998). Embryonic axis specification in nematodes evolution of the first step in development. *Current Biology* 8, 157-160.
- GOLDSTEIN, P. & TRIANTAPHYLLOU, A.C. (1980). The ultrastructure of sperm development in the plant parasitic nematode, *Meloidogyne hapla*. *Journal of Ultrastructure Research* 71, 143-153.
- GÖNCZY, P. & ROSE, L.S. (2005). Asymmetric cell division and axis formation in the embryo (October 15, 2005). *WormBook*, (Ed. The *C. elegans* Research Community), WormBook, doi/10.1895/wormbook.1.30.1, <http://www.wormbook.org>.

- GONZALEZ, L. (1979). Endotoquia matricida en nematodos en la familia Rhabditidae. *Agronomía Costarricense* 3, 191-192.
- GOODEY, T. (1929). On some new and little-known free-living nematodes. *Journal of Helminthology* 7, 27-62.
- GOODEY, T. (1939). *Cylindrocorpus* nom. nov. for *Cylindrogaster* Goodey, 1927 (Nematoda). *Journal of Helminthology* 17, 149-150.
- GOODEY, T. (1951). *Soil and Fresh water nematodes* – a monograph. London, 390 p.
- GOODEY, T. (1963). *Soil and freshwater nematodes* (Revised by Goodey, J.B.). 2nd edition. London: Methuen, 544p.
- GRASSE, P.P. (1965). *Traité de Zoologie*, Tome IV, Fascicule 2-3, Masson, Paris.
- GREEN, C.D. (1966). Orientation of male *Heterodera rostochiensis* Woll. and *H. schachtii* Schm. to their females. *Annals of Applied Biology* 58, 327-339.
- GREENWALD, I. (1997). Development of the Vulva. In *C. elegans II*. (Eds. Riddle, D.L. Blumenthal, T., Meyer, B. & Priess, J.), ColdSpring Harbor, New York: Cold Spring Harbor Laboratory Press.33, 519-541.
- GROOTAERT, P. (1976). The biology of *Mesodiplogaster theritieri* Maupas (Nematoda: Rhabditida). *Biologisch Jaarboek (Dodonaea)* 44, 191-202.
- GROOTAERT, P. & JAQUES, A. (1979). Description of *Butlerius degrissei* n. sp. (Nematoda: Diplogastridae) with notes on culture. *Nematologica* 25, 203-214.
- GROOTAERT, P. & MAERTENS, D. (1976). Cultivation and life cycle of *Mononchus aquaticus*. *Nematologica* 22, 173-181.
- GUNDY, S.D. VAN (1958). The life history of citrus nematode, *Tylenchulus semipenetrans* Cobb. *Nematologica* 3, 283-294.
- GUNHOLD, P. (1952). Uber die in Kompost lebenden Nematoden. *Bonner Zoologische Beitrage*. 3, 151-156.
- GUPTA, V.V.S.R. & YEATES, G.W. (1997). Soil microfauna as bioindicators of soil health. In *Biological Indicators of Soil Health*. (Eds. Pankhurst, C.; Doube, B.M. & Gupta, V.V.S.R.) New York, NY: CAB International. 201-233.
- GYSELS, H. & VAN DER HAEGEN, W. (1962). Post embryonale ontwikkeling en vervellingen van de vrijlevende Nematode *Panagrellus silusiae* (de Man, 1913), Goodey, 1945. *Natuurwet. Tijdschr.* 44, 3-20.
- HASEGAWA, K., FUTAI, K., MIWA, S. & MIWA, J. (2004). Early embryogenesis of the pinewood nematode *Bursaphelenchus xylophilus*. *Development, Growth and Differentiation* 46, 153-161.
- HECHLER, H.C. (1968). Post embryonic development and reproduction in *Diploscapter coronata* (Nematoda: Rhabditidae). *Proceedings of the Helminthological Society of Washington* 35, 24-30.
- HECHLER, H.C. (1970). Reproduction, chromosome number and postembryonic development of *Panagrellus redivivus* (Nematoda: Cephalobidae). *Journal of Nematology* 2, 355-361.

- HERMAN, M.A. (2006). Hermaphrodite cell-fate specification (January 09, 2006). *WormBook*, ed. The *C. elegans* Research Community, WormBook, doi/10.1895/wormbook.1.39.1, <http://www.wormbook.org>.
- HEYNS, J. (1962a). *Elaphonema mirabile* n. gen., n. sp. (Rhabditida), a remarkable new nematode from South Africa. *Proceedings of the Helminthological Society of Washington* 29, 128-130.
- HEYNS, J. (1962b). *Osstella hamata* n. gen., n. sp., *Zeldia serrata* n. sp., and *Acrobeles thornei* n. sp., three new nematodes from South Africa (Rhabditida: Cephalobidae). *Nematologica* 8, 301-306.
- HEYNS, J. (1968). *Paracrobeles laterellus* n. gen., n. sp. from South Africa (Nematoda: Cephalobidae). *Nematologica* 14, 511-514.
- HEYNS, J. (1969). South African species of the genus *Acrobeles* Von Linstow (Nematoda: Cephalobidae). *Phytophylactica* 1, 57-66.
- HEYNS, J. (1975). First report on a survey of the freshwater nematodes of South Africa. 2nd Symposium of the Nematologica Society Southern Africa, 8-9 Apr. 1975, Sellenbosch. Abstract. *Nematologica Society of Southern Africa*, Newsletter No. 7, 3.
- HEYNS, J. (1976). Preliminary results of a survey of freshwater nematodes in South Africa. *Journal of Limnological Society of Southern Africa* 2, 43-45.
- HEYNS, J. (1977a). Fresh water nematodes from South Africa. I. *Euteratocephalus* Andrassy, 1958. *Nematologica* 23, 112-118.
- HEYNS, J. (1977b). Fresh water nematodes from South Africa. 2. *Oncholaimus deconincki* n. sp. *Revue de Zoologie Africaine* 91, 906-912.
- HEYNS, J. (2002). Checklist of free-living nematodes recorded from freshwater habitats in Southern Africa. *Water SA* 28, 449-456.
- HEYNS, J. & COOMANS, A. (1977). Freshwater nematodes from South Africa 2. *Oncholaimus deconincki* n. sp. *Revue de Zoologie Africaine* 91, 906-912.
- HEYNS, J. & COOMANS, A. (1980). Freshwater nematodes from South Africa. 5. *Chronogaster* Cobb, 1913. *Nematologica* 26, 187-208.
- HEYNS, J. & COOMANS, A. (1983) New and known species of *Chronogaster* Cobb, 1913 (Nematoda: Leptolaimidae). *Nematologica* 29, 245-265.
- HEYNS, J. & COOMANS, A. (1989). Three *Monhystrella* species from inland waters in South West Africa – Namibia (Nematoda: Monhysteridae). *Nematologica* 35, 1-14.
- HIPPOCRATES. (460-375 B. C.). 1849. Works of Hippocrates, Translated by. Adams, F. London, "Aphorisms".
- HIRSCHMANN, H. (1951). Über das Vorkommen zweier Mundhöhlentypen bei *Diplogaster lheritieri* Maupas und *Diplogaster biformis* n. sp. und der Entstehung dieser hermaphroditischen Art aus *Diplogaster lheritieri*. I. Mitteilung. *Zoologische Jahrbücher (Systematik)* 80, 132-170.
- HIRSCHMANN, H. (1952). Die Nematoden der Wassergrenze mittelfränkischer Gewässer. *Zoologische Jahrbücher (Systematik)* 81, 313-364.

- HIRSCHMANN, H. (1956). Comparative morphological studies on the soybean cyst nematode, *Heterodera glycines* and the clover cyst nematode, *H. trifolii* (Nematoda: Heteroderidae). *Proceedings of Helminthological Society of Washington* 23, 140-151.
- HIRSCHMANN, H. (1962). The life cycle of *Ditylenchus triformis* with emphasis on post-embryonic development. *Proceedings of the Helminthological Society of Washington* 29, 30-43.
- HIRSCHMANN, H. & TRIANTAPHYLLOU A.C. (1967). Mode of reproduction and development of the reproductive system in *Helicotylenchus dihystra*. *Nematologica*, 13, 558-574.
- HNATEWYTSCH, B. (1929). Die Fauna der Erzgruben von Schneeberg im Erzgebirge. *Zoologische Jahrbücher (Systematik)*, 56, 173-261.
- HOLOVACHOV, O. & BOSTRÖM, S. (2006a). *Panagrolobus vanmegena*, a new genus and species from the Netherlands and Ukraine, with taxonomic revision and two new species of the genus *Teratolobus* Andrassy, 1968 (Nematoda: Cephalobidae). *Nematology* 8: 377-395.
- HOLOVACHOV, O. & BOSTRÖM, S. (2006b). Description of *Deleyia* gen. n. with a discussion of its phylogenetic relationships to the genera *Daubaylia* Chitwood & Chitwood, 1934 and *Myolaimus* Cobb, 1920 (Nematoda: Rhabditida). *Nematology* 8: 223-233.
- HOLOVACHOV O., BOSTRÖM S., TANDINGAN DE LEY I., NADLER S.A. AND DE LEY P. (2009). Description of *Penjatinema novaezeelandiae* sp. n. (Rhabditida: Cephalobidae) from New Zealand – a second species of a rare genus. *Journal of Nematode Morphology and Systematics* 12, 7-18.
- HOLOVACHOV, O., BOSTRÖM, S. & SUSULOVSKY, A. (2001). Description of *Acroukrainicus sagittiferus* gen. et sp. n. (Nematoda: Cephalobidae) from the area of an old sulphur industry in Ukraine. *Russian Journal of Nematology* 9, 51-56.
- HOLOVACHOV, O., BOSTRÖM, S., MUNDO-OCAMPO, M. & VILLENAVE, C. (2008). Description of two new species of the genus *Chiloplacus* Thorne, 1937 (Rhabditida: Cephalobidae) from Israel and Senegal. *Journal of Nematode Morphology and Systematics* 11, 147-157.
- HOLOVACHOV, O., ESQUIVEL, A. & BONGERS, T. (2003). Free-living nematodes from nature reserves in Costa Rica. 5. Bunonematoidea (Rhabditae). *Nematology* 5, 665-676.
- HOLOVACHOV, O., MUNDO-OCAMPO, M., BÖSTRÖM, S. & BUMBARGER, D. (2007). Three new species of genus *Cervidellus* Thorne, 1937 (Rhabditida: Cephalobidae) from the south-western part of North America. *Journal of Nematode Morphology and Systematics* 10, 49-61.
- HORVITZ, H. R. & STERNBERG, P. W. (1982). Nematode post-embryonic cell lineages. *Journal of Nematology* 14, 240-248.
- HOUTHOOFD, W. & BORGONIE, G. (2007). The embryonic cell lineage of the nematode *Halicephalobus gingivalis* (Nematoda: Cephalobina: Panagrolaimoidea). *Nematology* 9, 573-584.
- HOUTHOOFD, W., JACOBSEN, K., MERTENS, C., VANGESTEL, S., COOMANS, A. & BORGONIE, G. (2003). Embryonic cell lineage of the marine nematode *Pellioiditis marina*. *Developmental Biology* 258, 57-69.

- HOUTHOOFD, W., WILLEMS, M., JACOBSEN, K., COOMANS, A. & BORGONIE, G. (2008). The embryonic lineage of the nematode *Rhabditophanes* sp. *International Journal of Developmental Biology* 52, 963-967.
- HUGOT, J.P., BAUJARD, P. & MORAND, S. (2001). Biodiversity in helminths and nematodes as a field of study: an overview. *Nematology* 3, 199-208.
- HUNG, C. L. & JENKINS, W. R. (1969). Oogenesis and embryology of two plant-parasitic nematodes, *Pratylenchus penetrans* and *P. zaei*. *Journal of Nematology* 1, 352-356.
- HUNT, D.J. (1980). *Butlerius macrospiculum* n. sp. and *Cylindrocorpus walkeri* n. sp. (Nematode: Diplogastroidea) from St. Lucia, West Indies. *Revue de Nématologie*, 3, 155-160.
- HUSENI, T.A., AHMAD, I. & FIRDAUSI, N.S. (1997). Three new species in the superfamily Bunonematoidea (Rhabditida) from India. *Nematologia Mediterranea* 25, 41-48.
- HUSSAIN, A., TAHSEEN, Q., KHAN, R. & JAIRAJPURI, M.S. (2004). Description of two new species of *Myctolaimus* Cobb, 1920 (Nematoda: Cylindrocorporidae) from northern India. *Journal of Nematode Morphology and Systematics* 6, 95-102.
- INGLIS, W.G. (1961). Two new species of free-living marine nematodes from the West coast of Scotland. *Hydrobiologia* 18, 284-294.
- INGLIS, W.G. (1964a). The marine Enoplida (Nematoda): a comparative study of the head. *Bulletin of the British Museum (Natural History) Zoology* 11, 263-376.
- INGLIS, W. G. (1964b). The structure of the nematode cuticle. *Proceedings of the Zoological Society of London* 1, 465-502.
- INGLIS, W.G. (1965). *Nudora omercooperi* sp. nov. (Nematoda: Free-living) from an unusual locality in South Africa. *Hydrobiologia* 25, 176-180.
- INGLIS, W.G. (1967). The relationships of the nematode superfamily Seuratoidea. *Journal of Helminthology* 41, 115-136.
- INGLIS, W.G. (1970). Cyatholaimidae (Nematoda) from the coast of Western Australia. *Records of South Australia Museum*. 16, 1-13.
- INGLIS, W.G. (1983). An outline classification of the phylum Nematoda. *Australian Journal of Zoology* 31, 243-255.
- JACOBS, L.J. (1984). The free-living inland aquatic nematodes of Africa: a review. *Hydrobiologia* 113, 259-291.
- JAIRAJPURI, M.S. (1968). Some studies on the morphology and the biology of *Aphelenchus avenae* Bastian, 1865 (Nematoda: Aphelenchoidea). *Labdev Journal of Science and Technology* 6-B, 35, 27-34.
- JOHNIGK, S. A. & EHLERS, R.U. (1999). Juvenile development and life cycle of *Heterorhabditis bacteriophora* and *H. indica* (Nematoda: Heterorhabditidae). *Nematology* 1, 251-260.
- JONES, T.P. (1966). Sex attraction and copulation in *Pelodera teres*. *Nematologica* 12, 518-522.
- JUNGBLUT, B. & SOMMER, R.J. (1998). The *P. pacificus mab-5* gene is involved in the regulation of ventral epidermal cell fates. *Current Biology* 8, 775-778.

- KANZAKI, N., ABE, F., GIBLIN-DAVIS, R.M., KIONTKE, K., FITCH, D.H.A., HATA, K. & SONE, K. (2008). *Teratorhabditis synpapillata* Sudhaus, 1985 (Rhabditida: Rhabditidae) is an associate of the red palm weevil, *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae). *Nematology* 2, 207-218.
- KANZAKI, N., GIBLIN-DAVIS, R.M., WCISLO, W.T., ZENG, Y., YE, W., CENTER, B.J., ESQUIVEL, A. & THOMAS, W.K. (2010). *Acrostichus megaloptae* n. sp. (Nematoda: Diplogastridae), a phoretic associate of *Megalopta* spp. (Hymenoptera: Halictidae). *Nematology* 12, 453-468.
- KANZAKI, N., GIBLIN-DAVIS, R.M., ZENG, Y., YE, W. & CENTER, B.J. (2009). *Acrostichus rhynchophori* n. sp. (Rhabditida: Diplogastridae): a phoretic associate of *Rhynchophorus cruentatus* Fabricius and *R. palmarum* L. (Coleoptera: Curculionidae) in the Americas. *Nematology* 11, 669-688.
- KANZAKI, N., GIBLIN-DAVIS, R.M., ZENG, Y., YE, W., CENTER, B.J. & THOMAS, W.K. (2010). *Acrostichus puri* n. sp. (Nematoda: Diplogastridae), a phoretic associate of *Augochlora pura mosieri* Cockerell (Hymenoptera: Halictidae). *Nematology* 12, 49-64.
- KANZAKI, N., KOBAYASHI, M., NOZAKI, A. & FUTAE, K. (2006). *Myctolaimus platypi* sp. n. (Nematoda: Cyliandrocorporidae) isolated from the oak borer, *Platypus quercivorus* (Coleoptera: Platypodidae). *Nematology* 8, 717-725.
- KANZAKI, N., MASUYA, H. & KUBONO, T. (2008). Description of *Neodiplogaster crenatae* sp. n. and *N. acaloleptae* sp. n. (Nematoda: Diplogasteridae) from Japan. *Nematology* 10, 545-560.
- KAREGAR, A., DE LEY, P. & GERAERT, E. (1998). The genera *Cervidellus* Thorne, 1937 and *Stegelletina* Andrassy, 1984 from Iran, with description of *S. kheirii* sp. n. *Russian journal of Nematology* 6, 23- 36.
- KHAN, S.H. & COOMANS, A. (1980). Observations on the juvenile stages of *Miconchus studeri* (Nematoda: Mononchina). *Biologisch Jaarboek (Dodonaea)* 48, 111-118.
- KHAN, R., BAJAJ, H.K., SULTANA R. & TAHSEEN, Q. (2008). Description of *Diplogastrellus gracilis* (Bütschli, 1876) Paramonov, 1952, *D. sikorai* sp. n. and *Fictor composticola* sp. n. (Nematoda: Diplogastrina) from India. *Nematology* 10: 153-166.
- KHAN, R. & TAHSEEN, Q. (2005). *Pterygorhabditis saprophila* sp. n. and *Bunonema flexistoma* sp. n. (Nematoda: Bunonematoidea) from India. *International Journal of Nematology* 15, 155-162.
- KHERA, S. (1965). Nematodes from the banks of still and running waters, II. *Tridontus longicaudatus* n. gen., n. sp., subfamily Diplogasterinae Micoletzky, 1922 from India. *Nematologica* 11, 249-254.
- KHERA, S. (1967). *Acrobelinema cornis* n. gen., n. sp., subfamily Acrobelinae, Thorne from rhizosphere of millets from India. *Indian Journal of Nematology* 19, 159-163.
- KHERA, S. (1968). Nematodes from the banks of still and running waters. IV. Description of a new subgenus of *Rhabditis* and a new species from India (Subfamily: Rhabditinae). *Journal of the Zoological Society of India* 20, 38-41.
- KHERA, S. (1969). Nematodes from the banks of still and running waters. VI. Rhabditida from sewer. *Journal of Helminthology* 43, 347-363.

- KHERA, S. (1970). Nematodes from the banks of still and running waters. IX. Two new genera belonging to subfamily Diplogasterinae Micoletzky from India. *Revista Brasileira de Biologia* (Rio de Janeiro) 30, 405-409.
- KHERA, S. (1971). Nematodes from the banks of still and running waters. XI. Subfamily Rhabditinae. *Indian Journal of Nematology* 1, 237-243.
- KHERA, S. (1972). Nematodes from the banks of still and running waters. 12. Order Araeolaimida. *Proceedings of Zoological Society of India* 25, 49-58.
- KHERA, S. (1973). Studies on the embryology and post embryonic development of *Acrobelinema cornis* Khera (Nematoda: Cephalobidae) with special reference to development of gonads. *Records of Zoological Survey of India* 67, 165-189.
- KIKUTA, S., KIUCHI, T., AOKI, F. & NAGATA, M. (2008). Development of an entomopathogenic nematode, *Steinernemacarpocapsae*, in cultured insect cells under axenic conditions. *Nematology* 10, 845-851.
- KIMBLE, J. & HIRSH, D. (1979). The post-embryonic cell lineages of the hermaphrodite and male gonads in *Caenorhabditis elegans*. *Developmental Biology* 70, 396-417.
- KIONTKE, K. (1999). The *Rhabditis* (*Rhabditella*) *octopleura* species complex and descriptions of three new species. *Russian Journal of Nematology* 7, 71-94.
- KIONTKE, K. & FITCH, D.H.A. (2002). The phylogenetic relationships of *Caenorhabditis* and other rhabditids. In: *WormBook*, edited by The *C. elegans* Research Community, Online (URL: http://www.wormbook.org/chapters/www_phylogrhabditids/phylorhab.html).
- KIONTKE, K., GAVIN, N.P., RAYNES, Y., ROHRIG, C., PIANO, F., & FITCH, D.H.A. (2004). *Caenorhabditis* phylogeny predicts convergence of hermaphroditism and extensive intron loss. *Proceedings of National Academy of Sciences, USA* 101, 9003-9008.
- KIONTKE, K. & SUDHAUS, W. (1996). *Diplogaster* (*Diplogastrellus*) *cerea* sp. n. from Saguaro cactus rot and a revision of the subgenus *Diplogastrellus* (Nematoda: Diplogastridae). *Nematologica* 42, 173-197.
- KIONTKE, K. & SUDHAUS, W. (2000). Phasmids in male Rhabditida and other secernentean nematodes. *Journal of Nematode Morphology and Systematics* 3, 1-37.
- KIONTKE, K. & SUDHAUS, W. (2006). Ecology of *Caenorhabditis* species. In: *WormBook*, (Ed. The *C. elegans* Research Community), Online (<http://www.wormbook.org>).
- KIRJANOVA, E.S. (1951). Nematody poehvy kholkovogo polja i theliny v Golodnoj step (Uzbekistan). (In Russian). *Trudy Zoologicheskogo Instituta Akademiyi Nauk SSSR* 9, 625-657.
- KIRJANOVA, E.S. (1958). Antarkticheskie predstaviteli presnovodnykh nematod roda *Plectus* Bastian (Nematodes, Plectidae). (In Russian). *Information Bulletin of Soviet. Antarctic Expedition* 3, 101-103.
- KIRJANOVA, E.S. & KRALL, E. (1971). *Paraziticheskie nematody rastenij i mery borby s nimi. Tom II*. Leningrad, 523 p.
- KOKORDÁK, J. (1969). Beitrag zur Systematik und der Ökologie (der Nematoden) der koprophagen käfer. *Folia Veterinaria (Bratislava)*, 13, 143-158.

- KOLOTUEV, I. & PODBILEWICZ, B. (2004). *Pristionchus pacificus* vulva formation: polarized division, cell migration, cell fusion and evolution of invagination. *Developmental Biology*, 266, 322-333.
- KÖRNER, H. (1954). Die Nematodenfauna des vergehenden Holzes und ihre Beziehungen zu den Insekten. *Zoologische Jahrbücher Systematik* 82, 245-353.
- KORNFELD, K. (1997). Vulval development in *Caenorhabditis elegans*. *Trends Genet.* 13, 55-61.
- KREIS, H.A. (1930). Freilebende terrestrische Nematoden aus der Umgebung von Peking (China). *Zoologischer Anzeiger* 84, 283-294.
- KREIS, H.A. (1932). Beiträge zur Kenntnis pflanzenparasitischer Nematoden. *Zeitschrift für Parasitenkunde* 5, 184-194.
- KÜHNE, R. (1995). Nematoden mit phoretischer Bindung an Dung vergrabende *Geotrupes*-Käfer: ein morphologischer Vergleich von *Diplogaster henrichae* Sachs, 1950, *D. hieschmannae* Sachs, 1950, und *D. labiomorphus* sp. n. (Nematoda: Diplogasteridae). *Nematologica* 41, 7-34.
- LAHL, V., HALAMA, C. & SCHIERENBERG, E. (2003). Comparative and experimental embryogenesis of Plectidae (Nematoda). *Development Genes and Evolution* 213, 18-27.
- LAHL, V., SADLER, B. & SCHIERENBERG, E. (2006). Egg development in parthenogenetic nematodes: variations in meiosis and axis formation. *International Journal of Developmental Biology* 50, 393-398.
- LAM, A. B.O. & WEBSTER, J.N. (1971). Morphology and biology of *Panagrolaimus tipulae* n. sp. (Panagrolaimidae) and *Rhabditis (Rhabditella) tipulae* n. sp. (Rhabditidae), from leatherjacket larvae, *Tipula paludosa* (Diptera: Tipulidae). *Nematologica* 17, 201-212.
- LAMBERTI, F. (1975). Taxonomy of *Longidorus* (Micoletzky) Filipjev and *Paralongidorus* Siddiqi, Hooper & Khan. In : *Nematodes Vectors of Plant Viruses* (Eds. Lamberti, F., Taylor, C.E. & Seinhorst, J.W).. New York, Plenum Publishing Company, 71-79.
- LAMBERTI, F. & BLEVE-ZACHEO, T. (1977). Two new species of *Longidorus* (Nematoda: Longidoridae) from Italy. *Nematologia Mediterranea* 5, 73-83.
- LAMBERTI, F. & BLEVE-ZACHEO, T. (1979). Studies on *Xiphinema americanum* sensu lato with descriptions of fifteen new species (Nematoda: Longidoridae). *Nematologia Mediterranea* 7, 51-106.
- LAZAREVSKAYA, S.L. (1964). *Acrostichus minimus* n. sp. (Diplogasteroididae)-a new nematode of the timberman beetle. *Trudy Gelmintologicheskoi Laboratorii, Akademia Nauk SSSR (Moskva)*, 14, 122-127. (In Russian.).
- LEIDY, J. (1851). Contributions to Helminthology. *Proceedings of the Academy of Natural Sciences of Philadelphia* 5, 205-209, 224-227.
- LEUCKART, K.G.F.R. (1891). Über einen an *Aphodius fimetarius* sich verpuppenden freilebenden Rundwurm, *Rhabditis coarctata* n. sp. *Verhandlungen der Deutschen Zoologischen Gesellschaft*, 54-56.
- LINSTOW, VON O.F.B. (1878). *Compendium der helminthologie*. Hanover, Han'sche Buchhandlung, 382 pp.

- LOOF, P.A.A. (1961). The nematode collection of de Man, J.G. *Mededelingen Laboratory of Phytopathology* 190, 169-254.
- LOOF, P.A.A. (1964). Free-living and plant-parasitic nematodes from Venezuela. *Nematologica* 10, 201-300.
- LOOF, P.A.A. (1971). Free-living and plant-parasitic nematodes from Spitzbergen, collected by Van Rossen, H. *Mededelingen van de Landbouwhogeschool Wageningen* 71, 1-86.
- LOOF, P.A.A. (1973). Freshwater nematodes from Suriname collected by J. Van der Land. *Zoologische Verhandelingen* 129, 1-46.
- LORDELLO, L.G.E. (1951). Endotoquia matricida em *Rhabditis* sp. (Nematoda: Rhabditidae). *Anais da Escola Superior de Agricultura Luiz de Queiroz*. 8, 111-114.
- LORDELLO, L.G.E. & DE OLIVEIRA, A.J. (1963). Nematódeos associados a uma doenea da Palmeira imperial. *Revista Brasileira de Biologia* (Rio de Janeiro) 23, 19-24.
- LORDELLO, L.G.E. & ZAMITH, A.P.L. (1959). Observações sobre o genero “*Butlerius*” de nematódeos de vida livre. *Revista Brasileira de Biologia* (Rio de Janeiro) 19, 177-182.
- LORDELLO, L. G. E. & ZEM, A. C. (1977). Ocorrencia de endotoquia matricida em um nematoide da familia Rhabditidae. *Soc Brasil de Nematol* 2, 41-43.
- LORENZEN, S. (1981). Entwurf eines phylogenetischen Systems der freilebenden Nematoden. *Veröffentlichungen Institut für Meeresforschung In Bremerhaven, Supplement* 7, 1-472.
- LUDWIG, H. (1938). Die Variabilität von *Rhabditis teres* unter veränderten Ernährungsbedingungen. *Zeitschrift für Wissenschaftliche Zoologie* 151, 291-336.
- MAGGENTI, A. (1963). Comparative morphology in nemic phylogeny. In: Dougherty, E.C. (Ed.). *The lower Metazoa, comparative biology and phylogeny*. Berkeley, USA. University of California Press, 273-282.
- MAGGENTI, A. (1970). System analysis and nematode phylogeny. *Journal of Nematology* 2, 7-15.
- MAGNUS, A. (1193-1280). 1495 *De animalibus libri vigintisex novissime impressi*. Venezia.
- MAHAMOOD, MD. & AHMAD, I. (2009). Nematodes of Order Rhabditida from India. Description of *Diplogasteroides uriceus* sp. n. and *Rhabditoides papillatus* sp. n. (Diplogastrina) *Nematology* 11, 541-550.
- MAHAMOOD, MD. & AHMAD, I. (2009). Nematodes of the Order Rhabditida from India. First reports of *Paroigolaimella bernensis* and *Oigolaimella longicauda* (Nematoda: Diplogastrina) from the country. *International Journal of Nematology* 19, 151-158.
- MAHAMOOD, MD., AHMAD, I. & SHAH, A. A. (2006). Description of two new species of *Fictor* Paramonov (Nematoda: Diplogastrina). *International Journal of Nematology* 16, 200-207
- MAHAMOOD, MD., AHMAD, I. & SHAH, A. A. (2007). Nematodes of the order Rhabditida from India. Description of two species of *Mononchoides* (Nematoda: Diplogastrina). *Journal of Nematode Morphology and Systematics* 10, 1-10.

- MAHAMOOD, MD., SHAH, A. A. & AHMAD, I. & (2006). Nematodes of the order Rhabditida from India. Description of *Demaniella keirakensis* (Nematoda: Diplogastridae). *International Journal of Nematology* 16, 53-57.
- MANEGOLD, A. & KIONTKE, K. (2002). The association of two *Diplogasteroides* species (Secernentea: Diplogastrina) and cockchafers (*Melolontha* spp., Scarabaeidae). *Nematology* 3, 603-606.
- MANGO, S.E. (2007). The *C. elegans* pharynx: a model for organogenesis (January 22, 2007). *WormBook*, (Ed. The *C. elegans* Research Community), WormBook, doi/10.1895/wormbook.1.129.1, <http://www.wormbook.org>.
- MARTINI, E. (1903). Über Furchung und Gastrulation bei *Cucullanus elegans*. *Zeitschrift für Wissenschaftliche Zoologie* 74, 510-66.
- MARTINI, E. (1909). Über Subcuticula und Seitenfelder einiger Nematoden. IV. *Zeitschrift für Wissenschaftliche Zoologie* 93, 535-624.
- MASLER, E. P., ZASADA, I. A. & SARDANELLI, S. (2008). Hatching behaviour in *Heterodera glycines* in response to low temperature. *Comparative Parasitology* 75, 76-81.
- MASSEY, C.L. (1960a). Nematode parasites and associates of the California five-spined engraver, *Ipsconfusus* (Les.). *Proceedings of the Helminthological Society of Washington* 27, 14-22.
- MASSEY, C.L. (1960b). A new species of Nematoda, *Cylindrocorpus erectus* associated with *Scolytus multistriatus* Marsh in American elm. *Proceedings of the Helminthological Society of Washington* 27, 42-44.
- MASSEY, C.L. (1962). New species of Diplogasteridae (Nematoda) associated with bark beetles in the United States. *Proceedings of the Helminthological Society of Washington* 29, 67-75.
- MASSEY, C.L. (1966a). The genus *Acrostichus* Rahm, 1928, synonym *Diplogasteritus* Paramanov, 1952 (Nematoda). *Proceedings of the Helminthological Society of Washington* 33, 8-13.
- MASSEY, C.L. (1966b). The genus *Micoletzkyia* (Nematoda) in the United States. *Proceedings of the Helminthological Society of Washington* 33, 13-19.
- MASSEY, C.L. (1966c). The nematode parasites and associates of *Dendroctonus adjunctus* (Coleoptera: Scolytidae) in New Mexico. *Annals of the Entomological Society of America* 59, 424-440.
- MASSEY, C.L. (1967). Nematodes associated with tree-infesting insects: Paurodontidae new family and Misticinae new subfamily with a description of one new genus and four new species. *Canadian Journal of Zoology*, 45, 779-786.
- MASSEY, C.L. (1971). Nematodes associates of several species of *Pissodes* (Coleoptera: Curculionidae) in the United States. *Annals of the Entomological Society of America* 64, 162-169.
- MASSEY, C.L. (1974). Biology and taxonomy of nematode parasites and associates of bark beetles in the United States. *Agricultural Forestry Service (Washington), Agricultural Handbook*, 446, 1-233.
- MAUPAS, E.F. (1899). La mue e t l'enkystement chez les nematodes. *Archives de Zoologie Expérimentale et Générale* 7, 563-628.

- MAUPAS, E.F. (1900). Modes et forme de reproduction des nématodes. *Archives de Zoologie Expérimentale et Générale* 7, 563-628.
- MAUPAS, E.F. (1915). Un nouveaux *Rhabditis* d'Algérie. *Bulletin de la Societe d'Histoire naturelle de l'Afrique du Nord. Alger* 7, 51-52.
- MAUPAS, E.F. (1916). Nouveaux *Rhabditis* d' Algérie. *Comptes Rendus des Séances de la Societe de Biologie et de ses Filiales* (Paris) 79, 607-613.
- MAUPAS, E.F. (1919). Essais d'hybridation chez les nématode. *Bulletin Biologique France et Belgique* 52, 466-498.
- MCGHEE, J. D. (2007). The *C. elegans* intestine (March 27, 2007). *WormBook*. (Ed. The *C. elegans* Research Community). WormBook, doi/10.1895/wormbook.1.133.1, <http://www.wormbook.org>.
- MC LAREN, D.J. (1973). The structure and development of the spermatozoon of *Dipetalonema vitae* (Nematoda: Filarioidea). *Parasitology* 66, 447-463.
- MEYL, A.H. (1957). Freelifving nematodes: Exploration Hydrobiologique du Lac Tanganica (1945-1947). *Institut Royal des Sciences naturelles de Belgique, Bruxelles*, 3, 27-51.
- MEYL, A.H. (1960/1961). Die freilebenden Erd- und Süßwassernematoden (Fadenwürmer). In: *Die Tierwelt Mitteleuropas* (Eds. Brohmer, P., Ehrmann, P. & Ulmer, G.). 1, 1-164.
- MICOLETZKY, H. (1922). Die freilebenden Erd-Nematoden mit besonderer Berücksichtigung der steiermark und der Bukowina, Zugleich mit einer revision sämtlicher nichtmariner, freilebender Nematoden in Form von Genus-Beschreibungen und Bestimmungsschlüsse. *Archiv für Naturgeschichte Abteilung* 87, 1-650.
- MORAND, S. & BARKER, G.M. (1995). *Hugotdiplogaster neozelandia* n. gen., n. sp. (Nematoda: Diplogasteridae), a parasite of the New Zealand endemic slug, *Athoracophorus bitentaculatus* (Quoy & Gaimard, 1832) (Gastropoda: Athoracophoridae). *New Zealand Journal of Zoology* 22, 109-113.
- MULVEY, R.H. (1955). Oogenesis in several free-living and plant parasitic nematode. *Canadian Journal of Zoology* 33, 295-310.
- MULVEY, R.H. (1961). The Mononchidae: a family of predaceous nematodes. II. Genus *Anatonchus* (Enoplida: Mononchidae). *Canadian Journal of Zoology* 39, 807-826.
- MUNDO-OCAMPO, M., BALDWIN, J.G., DORADO-RAMÍREZ, O. & MORALES-RUIZ, M.C. (2003). *Acrobeles zapatai* n. sp. (Rhabditida: Cephalobidae) from the Biosphere Reserve "Sierra de Huautla", (Mexico), with a discussion of the taxonomic limits of the genus. *Journal of Nematode Morphology and Systematics* 5, 21-32.
- MUSCHIOL, D. & TRAUNSPURGER, W. (2007). Life cycle and calculation of the intrinsic rate of natural increase of two bacterivorous nematodes, *Panagrolaimus* sp. and *Poikilolaimus* sp. from chemoautotrophic Movile Cave, Romania. *Nematology* 9, 271-284.
- NANCE, J., LEE, J. Y. & GOLDSTEIN, B. (2005). Gastrulation in *C. elegans* (September 26, 2005). *WormBook*, (Ed. The *C. elegans* Research Community), WormBook, doi/10.1895/wormbook.1.23.1, <http://www.wormbook.org>.
- NADLER, S.A., DE LEY, P., MUNDO-OCAMPO, M., SMYTHE, A.B., STOCK, S.P., BUMBARGER, D., ADAMS, B.J., DE LEY, I.T., HOLOVACHOV, O. & BALDWIN, J.G. (2006). Phylogeny of

- Cephalobina (Nematoda): Molecular evidence for recurrent evolution of probolae and incongruence with traditional classifications. *Molecular Phylogenetics and Evolution* 40, 696-711.
- NEEDHAM, F. (1743). *An account of some new microscopical discoveries*. London, 126 p.
- NEHER, D.A. (2001). Role of nematodes in soil health and their use as indicators. *Journal of Nematology* 33, 161-168.
- NESTEROV, P.I. (1970). *Acromoldavicus* n. gen. I pereopisanie vida *Acrobeloides skrjabini* Nesterov et Lisethkaja, 1965 (Nematoda, Cephalobidae). *RIO Akademii Nauk Moldavskoi SSR* 5, 134-138.
- NEUHAUS, C. (1903). Die postembryonale Entwicklung der *Rhabditis nigrovenosa*. *Jenaische Zeitschrift für Naturwissenschaft*. 37, 653-690.
- NICHOLAS, W.L., DOUGHERTY, E.C. & HANSEN, E.L. (1959). Axenic cultivation of *Caenorhabditis briggsae* (Nematoda: Rhabditidae) with chemically undefined supplements; comparative studies on related species. *Annals of the New Yourk Academy of Sciences*. 77, 218-236.
- NIGON, V., GUERRIER, P. & MONIN, H. (1960). L'Architecture polaire de l'oeuf et mouvements des constituants cellulaires au cour des premières étapes du developpement chez quelque nematodes. *Bulletin Biologique de la france et de la Belgique* 94, 132-201.
- ÖRLEY, L. (1880). Az anguillulidák magánrajza. (Monographie der Anguilluliden). *Természetráji Füzetek* (Budapest) 4, 16-150.
- OSCHE, G. (1952). Systematik und Phylogenie der Gattung *Rhabditis* (Nematoda). *Zoologische Jahrbücher (Systematik)* 81, 190-280.
- PAESLER, F. (1939). Massenvorkommen eines neuen Nematoden *Diplogaster schneideri* n. sp. im braunen Schleimfluß einer Roßkastanie. *Zoologischer Anzeiger* 126, 138-140.
- PAESLER, F. (1946). Beitrag zur Kenntnis der im Dünger Bucht. Kieler Bucht. *Kieler Meeresforschungen* 1, 1-48.
- PAESLER, F. (1962). Nematoden in Reisspreu. Beschreibung einiger Arten und Sukzessionserscheinungen. *Opuscula Zoologica Budapest* 4, 105-120.
- PAETZOLD, D. (1955). Untersuchungen an freilebenden Nematoden der Salzwiese bei Aseleben. *Wissenschaftliche Zeitschrift der Universität Halle, Mathematisch-Naturwissenschaftliche Reihe* 4, 1057-1090.
- PAETZOLD, D. (1958). Bemerkungen Zur "Endotokia matricida" von Lordello 1951 *Wissenschaftliche Beiträge, Martin-Luther-Universität Halle-Wittenberg Mathematisch-Naturwissenschaftliche* 7, 81-84.
- PAI, S. (1927). Lebenszyklus der *Anguillula aceti*. *Ehrbg. Zoologische Anzeiger* 74, 257-270.
- PAI, S. (1928). Die Phasen der Lebenszyklus der *Anguillula aceti*. *Ehrbg. Und. Ihre. Experimentell morphologische Beeinflussung. Zeitschrift für Wissenschaftliche Zoologie* 131, 293-344.
- PARAMONOV, A.A. (1951). K poznaniyu saprozoiynykh nematode v SSSR. *Trudy Zoologiceskogo Instituta Akademii Nauk SSSR (Moskva)* (In Russian). 9, 602-612.

- PARAMONOV, A.A. (1952). Opyt ekologitscheskoi klassifikacii fitonematod. *Trudy Gelmintologicheskoi Laboratorii, Akademia Nauk CCCR* 6, 338-369.
- PARAMONOV, A.A. (1964). Osnovy fitogelmintologii II. [Fundamentals of phytonematology.] (In Russian). *Trudy Gelmintologicheskoi Laboratorii, Akademia Nauk SSSR (Moskva)* 446 p.
- PEARSE, A.S. (1942). *An introduction to Parasitology*. Baltimore, USA: Thomas, C.C. 375p.
- PEREZ, J. (1866). Recherches anatomiques et physiologiques sur l'anguillule terrestre (*Rhabditis terricola* Dujardin). *Annales des Sciences Naturelles Zoologie (Series 5)* 6, 152-307.
- PILLAI, J. K. & TAYLOR, D.P. (1968). *Butlerius micans* n. sp. (Nematoda: Diplogasteridae) from Illinois, with observations on its feeding habits and a key to the species of *Butlerius* Goodey, 1929. *Nematologica* 14, 89-93.
- PILLAI, J. K. & TAYLOR, D.P. (1968). Emendation of the genus *Demaniella* Steiner, 1914 (Nematoda: Diplogasterinae) with observations on the biology of *D. basili* n. sp. *Nematologica*, 14, 285-294.
- PIRES-DA SILVA, A. & SOMMER, R.J. (2004). Conservation of the global sex determination gene tra-1 in distantly related nematodes. *Genes and Development* 18, 1198-1208.
- POINAR, G.O. & HESS, R (1985). Spermatogenesis in the insect-parasitic nematode, *Heterorhabditis bacteriophaga* Poinar (Heterorhabditidae: Rhabditida). *Revue de Nematologie* 8, 357-367.
- POINAR, G.O. (1969). Diplogasterid nematodes (Diplogasteridae: Rhabditida) and their relationship to insect disease. *Journal of Invertebrate Pathology* 13, 447-454.
- POINAR, G.O. (1979). *Parasitodiplogaster sycophilon* gen. n., sp. n. (Diplogaster: Nematoda), a parasite of *Elisabethiella stukenbergi* Grandi (Agaonidae: Hymenoptera) in Rhodesia. *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen (Ser. C)* 82, 375-381.
- POINAR, G.O. (1990). Redescription of *Chroniodiplogaster aerivora* (Cobb) gen. n., comb. n.. (Rhabditida: Diplogasteridae) from Termites. *Journal of the Helminthological Society of Washington* 57, 26-30.
- POINAR, G.O. JR., TRIGGIANI, O. & MERRITT, R.W. (1976). Life history of *Eudiplogaster aphodii* (Rhabditida: Diplogasteridae), a facultative parasite of *Aphodius fimetarius* (Coleoptera: Scarabaudae). *Nematologica* 22, 79-86.
- POINAR, G.O., JACKSON, T.A., BELL, N.L. & WAHID, M.B. (2003). *Cylindrocorpus inevectus* sp. n. associated with the oil palm weevil, *Elaeidobius kamerunicus* (Faust) (Coleoptera: Curculionidae), with a synopsis of the family Cylindrocorporidae and establishment of Longibuccidae n. fam. (Diplogastroidea: Nematoda). *Nematology* 5, 183-190.
- POIRAS, L., BALDWIN, J.G., MUNDO-OCAMPO, M & BUMBARGER, D.J. (2002). *Nothacrobeles borregi* n. sp. (Nematoda: Cephalobidae) from Anza Borrego, California and *N. laticollaris* (De Ley and Vandebroek, 1992) n. comb. *Nematology* 4, 865-874.
- POPOVICI, I. (1972). Studies on the biology and population development *Cephalobus persegnis* (Nematoda: Cephalobidae) in agar culture. *Pedobiologia* 12, 123-127.
- RAHM, G. (1924). Beitrag zur Kenntnis der Moostierwelt der preussischen Rheinlande. I. *Archiv für Naturgeschichte* 90 A, S. 174-176.

- RAHM, G. (1928). Alguns nematodes parasitas e semi-parasitas das plantas cultiveiras do Brasil. *Archivos do Instituto de Biológico de Defesa Agrícola e Animal (Sao Paulo)* 1, 239-251.
- RAHM, G. (1929). Nematodes parasitas e semi-parasitas de diversas plantas cultiveiras do Brasil. *Archivos do Instituto de Biológico de defesa Agrícola e Animal (Sao Paulo)* 2, 67-136.
- RAHM, G. (1937). Freilebenden Nematoden vom Yan-Chia-Ping-Tal (Nord China). *Zoologischer Anzeiger* 119, 87-97.
- RASHID, F. & HEYNS, J. (1990a). *Chiloplacus* and *Macrolaimellus* species from South West Africa/Namibia (Nematoda: Cephalobidae). *Phytophylactica* 22, 189-199.
- RASHID, F. & HEYNS, J. (1990b). Description of *Namibinema scaphovulva* n. gen., n. sp. and *Zeldia punctata* (Thorne, 1925) from Namibia (Nematoda: Cephalobidae). *Phytophylactica* 22, 397-403.
- RASHID, F., GERAERT, E. & SHARMA, R.D. (1984). Morphology, taxonomy and morphometry of some Cephalobidae (Nematoda: Rhabditida) from Brazil, with descriptions of two new genera and four new species. *Nematologica* 30, 251-299.
- RASHID, F., GERAERT, E., COOMANS, A. & SUATMADJI, W. (1989). Cephalobidae from the Krakatau region (Nematoda: Rhabditida). *Nematologica* 34, 125-143.
- RATHORE, S.S. & NAMA, H.S. (1992). On two nematodes *Acrobeloides conoidis* n. sp. and *Chiloplacus jodhpurensis* n. sp. from soil around roots of *Brassica juncea*. *Proceedings of the National Academy of Science, India (Section B, part 4)* 62, 509-513.
- REITER, M. (1928). Zur Systematik und Ökologie der zweigeschlechtlichen Rhabditiden. *Arbeiten des Zoologischen Institutes der Universität Innsbruck* 3, 91-184.
- RIDDLE, D. L., BLUEMENTHAL, T., MEYER, B. J. & PRIESS, J. R. (1997). *C. elegans* II. Cold Spring Harbor Laboratory Press. Cold Spring Harbor. 1222 pp.
- RIVAS, X. & ROMAN, J. (1985). De Sarrollo post-embryonario de una poblacion de *Radopholus similis* de Puerto Rico; *Nematropica* 15, 37-41.
- ROMAN, J. & HIRSCHMANN, H. (1969). Embryogenesis and post-embryogenesis in the species of *Pratylenchus* (Nematoda: Tylenchidae). *Proceedings of Helminthological Society Washington* 36, 164-174.
- ROSSNER, J. (1971). Untersuchungen zur Entwicklung von *Rotylenchulus robustus*. *Nematologica* 17, 255-261.
- ROWSE, T.W. (1969). Embryology of *Aphelenchoides dactylocercus*. (Abstract). *Journal of Nematology* 1, 26.
- ROY, T.K. (1973a). Selection of media for culturing free-living nematodes: Axenic culture of *Chiloplacus lentus* Thorne (Nematoda: Rhabditidae). *Indian Journal of Nematology* 3, 64-70.
- ROY, T.K. (1973b). Intra-uterine egg development in *Chiloplacus lentus*. *Indian Journal of Nematology* 3, 82.
- RUDEL, D., RIEBESELL, M. & SOMMER, R. J. (2005). Gonadogenesis in *Pristionchus pacificus* and organ evolution: development, adult morphology and cell/cell interactions in the hermaphrodite gonad. *Developmental Biology* 277, 200-221.

- RUDOLPHI, C.A. (1809). *Entozoorum synopsis cui accedunt mantissa duplex et indices locupletissimi*. Berlin Berolini 811p.
- RUDOLPHI, C.A. (1819). *Entozoorum synopsis cui accedunt mantesia duplex et indices locupletissimi*. Berolini, 811p.
- RÜHM, W. (1954). Die Nematoden als Kommensalen, Halbparasiten und Parasiten der Insekten. *Deutscher Entomologentag in Hamburg* 1953, 168-184.
- RÜHM, W. (1956a). Die Nematoden der Ipiden. *Parasitologische Schriftenreihe* (Jena). 6, 1-437.
- RÜHM, W. (1956b). Bemerkungen über die Pseudodiplogasteroidinae Körner 1954 (Nematoda). *Zoologischer Anzeiger* 156, 293-299.
- RÜHM, W. (1959a). *Cylindrocorpus* subg. Nov. und *Protocylindrocorpus* subg. Nov., zwei Untergattungen der Gattung *Cylindrocorpus* Goodey, 1939. *Nematologica* 4, 76-82.
- RÜHM, W. (1959b). *Diplogasteroides (Rhabdontolaimus) berwigi* n. sp., ein mit einer Raspelplatte ausgestattete *Diplogaster*- Art (Nematoda). *Zoologischer Anzeiger*, 162, 356-361.
- SACHS, H. (1950). Die Nematodenfauna der Rinderexkreme. Eine ökologisch systematische studie. *Zoologische Jahrbücher (Systematik)* 79, 209-272.
- SAMOILOFF, M.R. (1987). Nematodes as indicators of toxic environmental contaminants. In: *Vistas on Nematology*. (Eds Veech, J.A. and Dickson, D.W.). A Commemoration of the twenty-fifth anniversary of the society of nematologists, Hyattsville, MD: Society of Nematologists, Inc. 433-439.
- SANWAL, K.C. (1959). *Cuticonema vivipara* n. gen., n. sp., a new saprophagous nematode of the subfamily Panagrolaiminae Thorne, 1937, from Canada, with a note on the genus *Brevibucca* Goodey, 1935. *Canadian Journal of Zoology* 37, 223-229.
- SANWAL, K.C. (1960). *Panagrellus dubius* n. sp. (Nematoda: Turbatricinae Goodey, 1943), from frass of beetles *Sternochetus lapathi* with remarks on redescriptions of *Anguillula rediviva*. *Canadian Journal of Zoology* 38, 1041-1046.
- SARR, E. COOMANS, A. & LUC, M. (1987). Development and life cycle of *Neodolichodorus rostrulatus* (Siddiqui, 1976), with observations on the copulatory plug (Nematoda: Tylenchina). *Revue de Nematologie* 10, 87-92.
- SAUER, M.R. & ANNELLS, C.M. (1984). Lip region structure in Acrobelinae (Nematoda: Cephalobidae). *Nematologica* 30, 140-150.
- SAUER, M.R. CHAPMAN, R.N. & BRZESKI, M.W. (1979). External morphology of some Acrobelinae. *Nematologica* 25, 482-486.
- SCHIERENBERG, E. (2000). Early development of nematode embryos: differences and similarities. *Nematology*, 2, 57-64.
- SCHIERENBERG, E. (2001). Three sons of fortune: early embryogenesis, evolution and ecology of nematodes. *BioEssays* 23, 841-847.
- SCHIERENBERG, E. (2005). Unusual cleavage and gastrulation in a freshwater nematode: developmental and phylogenetic implications. *Development Genes and Evolution* 215, 103-108.

- SCHIERENBERG, E. (2006). Embryological variation during nematode development. (Ed. The *C. elegans* Research Community). WormBook, 1-13.
- SCHIERENBERG, E. & LAHL, V. (2004). Embryology and phylogeny of nematodes. In: *Nematology Monographs & Perspectives* (Eds Cook, R.C., & Hunt, D.J.), Vol. 2, (Leiden: Brill), pp. 667-679.
- SCHIERENBERG, E., WIEGNER, O., BOSSINGER, O., SKIBA, F. & KUTZOWITZ, M. (1997). Pattern formation and cell specification in nematode embryos: A theme with considerable variations. *Zoology-Analysis of Complex Systems* 100, 320-327.
- SCHNEIDER, A.F. (1860). Über die Muskeln und Nerven der Nematoden. *Archiv für Anatomie, Physiologie und wissenschaftliche Medizin* 224-242
- SCHNEIDER, A.F. (1866). *Monographie der Nematoden*. Berlin, 357 p.
- SCHNEIDER, W. (1923). Niederrheinische freilebende Nematoden. *Zoologischer Anzeiger* 56, 264-281.
- SCHNEIDER, W. (1937). Freilebende Nematoden der Deutschen Limnologischen Sundaexpedition nach Sumatra, Java und Bali. *Archiv für Hydrobiologie, Supplement* 15, 30-108.
- SCHNEIDER, W. (1938). *Diplogaster pararmatus* n. n. nebst Bemerkungen über einige andere *Diplogaster*-Arten. *Zoologischer Anzeiger* 121, 38-43.
- SCHNEIDER, W. (1943). Freilebende Nematoden aus dem Ohridsee. *Posebna Izanija. Srpska Akademija Nauka odeljenje prirodno-matematickih Nauka (Beograd)* 136, 161-184.
- SCHULTE F., (1989). Description of *Rhabditis (Pelodera) pseudoterres* n. sp. (Rhabditidae: Nematoda) with a redescription of its sibling *R. (P.) teres* (Schneider, 1866). *Revue de Nématologie* 12, 387-394.
- SCHULTE, F. (1989). The association between *Rhabditis necromena* Sudhaus & Schulte, 1989 (Nematoda: Rhabditidae) and native and introduced millipedes in South Australia. *Nematologica* 35, 82-89.
- SCHULTZE, M. (1857). *Diplogaster*, Schultze 1857, in Carus, V. *Icones Zootomicse* 1, explan. pl. viii. Verm (Nema.). 2, 109.
- SCHUURMANS STEKHOVEN, J.H. (1943). Nématodes recueillis dans des grottes et des sources en Belgique. *Bulletin du Musée Royal d'Histoire Naturelle de Belgique* 19, 1-20.
- SCHUURMANS STEKHOVEN, J.H. (1951). Nématodes saprozoaires et libres du Congo belge. *Mémoire de l'Institut royale des Science naturelles de Belgique* 39, 1-79.
- SCHUURMANS STEKHOVEN, J.H. & DE CONINCK, L.A.P. (1933). Morphologisches Fragen zur Systematik der freilebenden Nematoden. *Verhandlungen der Deutschen Zoologischen Gesellschaft* 35, 138-143.
- SCHUURMANS STEKHOVEN, J.H. & TEUNISSEN, R.J.H. (1938). Nématodes libres terrestres. *Exploration du Parc National Albert, Mission G. F. De Witte (1933-1935) Fasc.*, 22, 1-229.
- SCOTT, H.L. & WHITTAKER, F.H. (1970). *Pelodera strongyloides* Schneider 1866: A potential research tool. *Journal of Nematology* 2, 193-203.

- SEINHORST, W. (1959). A rapid method for the transfer of nematodes from fixative to anhydrous glycerin. *Nematologica* 4, 67-69.
- SESHADRI, A.R. (1965). Investigations on the biology and life cycle of *Criconemoides xenoplax* Raski, 1952 (Nematoda: Criconematidae). *Nematologica* 10, 540-562.
- SEYMOUR, M.K. (1973). Motion and the skeleton in small nematodes. *Nematologica* 19, 43-48.
- SHAFQAT, S., JAIRAJPURI, M.S. & BILGRAMI, A.L. (1991). Developmental biology of *Dorylaimus stagnalis* Dujardin, 1845 (Nematoda: Dorylaimida). *Revue de Nematologie* 14, 61-71.
- SHAHINA, F. & DE LEY, P. (1997). Two new species of Cephalobidae from Valle de la Luna, Argentina, and observation on the genera *Acrobeles* and *Nothacrobeles* (Nematoda: Rhabditida). *Fundamental and applied Nematology* 20, 329-347.
- SHEMER, G., KISHORE, R., & PODBILEWICZ, B. (2000). Ring formation drives invagination of the vulva in *Caenorhabditis elegans*: Ras, cell fusion, and cell migration determine structural fates. *Developmental Biology*. 221, 233–248.
- SHEPHERD, A. M. & CLARK, S. A. (1976). Spermatogenesis and the ultrastructure of sperm and of the male reproductive tract of *Aphelenchoides blastophthorus* (Nematoda: Tylenchida, Aphelenchida). *Nematologica* 22, 1-9.
- SHEPHERD, A.M., CLARK, S.A. & KEMPTON, A. (1974). Spermatogenesis and sperm ultrastructure in some cyst nematodes, *Heterodera* spp. *Nematologica* 19, 551-560.
- SHER, S.A. (1961). Revision of the Hoplolaiminae (Nematoda) I. Classification of nominal genera and nominal species. *Nematologica* 6, 155-169.
- SHER, S.A. (1963). Revision of the Hoplolaiminae (Nematoda) II. *Hoplolaimus* Daday, 1905 and *Aorolaimus* n. gen. *Nematologica* 9, 267-295.
- SHER, S.A. (1966). Revision of the Hoplolaiminae (Nematoda) VI. *Helicotylenchus* Steiner, 1945. *Nematologica* 12, 1-56.
- SHER, S.A. (1968). Revision of the genus *Hirschmanniella* Luc and Goodey, 1963 (Nematoda: Tylenchoidea). *Nematologica* 14, 243-275.
- SHER, S.A. (1970). Reclassification of the genus *Chitinotylenchus* (Micoletzky, 1922) and a redescription of *C. paragracilis* (Micoletzky, 1922) (Nematoda: Tylenchoidea). *Journal of Nematology* 2, 236-238.
- SHINYA, R., TAKEUCHI, Y. & FUTAI, K. (2009). A technique for separating the developmental stages of the propagative form of the pine wood nematode, *Bursaphelenchus xylophilus*. *Nematology* 11, 305-307.
- SHOKOOHI, E., ABOLAFIA, J. & ZAD, J. (2007). Nematodes of the order Rhabditida from Tehran province, Iran. The family Panagrolaimidae with description of *Halicephalobus persicus* sp. n. and a key to species of *Halicephalobus* Timm, 1956. *Nematology* 9, 693-711.
- SHOSHIN, A.V. (1989). New nematode species of the families Diplogasteridae and Diplogasteroididae from the Baikal Lake. In: *Tylenchida and Rhabditida (Nematoda)- Parasites of Plants and Insects*. (Ed. Ryss, A.Y..) Leningrad. *Trudy Zoologicheskogo Instituta Akademia Nauk SSSR* 194, 83-95.

- SIDDIQI, M.R. (1993). Nematodes of Rainforests: 2. Five new genera and eight new species of cephalobs. *Afro-Asian Journal of Nematology* 3, 212-225.
- SIDDIQI, M.R. (2002). *Catoralaimellus cornutus* gen. n., sp. n., and two new species of *Macrolaimellus* (Nematoda: Cephalobidae). *International Journal of Nematology* 12, 89-97.
- SIGRIST, C.B. & SOMMER, R.J. (1999). Vulva formation in *Pristionchus pacificus* relies on continuous gonadal induction. *Development Genes and Evolution* 209, 451-459.
- SINGH, R.V. & KHERA, S. (1976). A case of *endotokia matricida* in *Acrobeloides* sp. *Indian Journal of Nematology* 6, 103-104.
- SIVAKUMAR, C.V. & SESHADRI, A.R. (1971). Life cycle of reniform nematode, *Rotylenchulus reniformis* Linford and Oliveira, 1940. *Indian Journal of Nematology* 1, 91-93.
- SKIBA, F. & SCHIERENBERG, E. (1992). Cell lineages, developmental timing, and spatial pattern formation in embryos of free-living soil nematodes. *Developmental Biology* 151, 597-610.
- SKRJABIN, K.I., SHIKHOBALOVA, N.P., SOBOLEV, A.A., PARAMONOV, A.A. & SUDARIKOV, V.E. (1954). Kamallanaty, rabditaty, tilenkhaty, trikhotsefalyaty, dioktofimaty i raspredelenie paraziticheskikh namatod pokhoziaevam. In: *Opredelitel' Paraziticheskikh Nematode* 4, Moskva 1-827.
- SKWARRA, E. (1921). Diagnosen neuer freilnemder Nematoden Ostpreuens. *Zoologischer Anzeiger* 53, 66-74.
- SMART, G. & NGUYEN, K.B. (1994). *Rhabditis (Oscheius) pheropsophi* n. sp. (Rhabditida: Rhabdidea). *Journal of Nematology* 26, 19-24.
- SMYTHE, A.B. & NADLER, S.A. (2006). Molecular phylogeny of *Acrobeloides* and *Cephalobus* (Nematoda: Cephalobidae) reveals paraphyletic taxa and recurrent evolution of simple labial morphology. *Nematology* 8, 819-839.
- SOHLENIUS, B. (1979). A carbon budget for nematodes, rotifers and tardigrades in a Swedish coniferous forest soil. *Holarctic Ecology* 2, 30-40.
- SOMMER, R.J. (2000). Evolution of nematode development. *Current Opinion in Genetics and Development* 10, 443-448.
- SOMMER, R.J. (2005). Evolution of development in nematodes related to *C. elegans* (December 14, 2005), In: *WormBook*. (Ed. The *C. elegans* Research Community). available online at <http://www.wormbook.org>, doi/10.1895/wormbook.1.46.1.
- SOMMER, R.J., CARTA, L.K., KIM, S.Y. & STERNBERG, P.W. (1996). Morphological, genetic, and molecular description of *Pristionchus pacificus* sp. n. (Nematoda: Neodiplogastridae). *Fundamental and Applied Nematology* 19, 511-521.
- SOMMER, R.J., EIZINGER, A., LEE, K.Z., JUNGBLUT, B., BUBECK, A. & SCHLAK, I. (1998). The *Pristionchus* HOX gene *Ppa-lin-39* inhibits programmed cell death to specify the vulva equivalence group and is not required during vulval induction. *Development* 125, 3865-3873.
- SOMERS, J.A., SHOREY, H. H. & GASTON, L. K. (1977). Sex pheromone communication in the nematode *Rhabditis pellio*. *Journal of Chemical Ecology* 3, 467-474.

- SOMMER, R.J. & STERNBERG, P.W. (1994). Changes of induction and competence during the evolution of vulva development in nematodes. *Science* 5168, 114-118.
- SOMMER, R.J. & STERNBERG, P.W. (1995). Evolution of cell lineage and pattern formation in the vulval equivalence groups of rhabditid nematodes. *Developmental Biology* 167, 61-74.
- SOMMER, R.J. & STERNBERG, P.W. (1996). Evolution of nematode vulva fate patterning. *Developmental Biology* 17, 396-407.
- SOÓS, A. (1938). Zwei neue tyrophobionte Nematoden-Arten. *Zoologischer Anzeiger* 124, 281-286.
- SPEMANN, H. (1895). Zur Entwicklung des *Strongylus paradoxus*. *Zoologische Jahrbücher*, 8, 301-317.
- STEINBUCH, J.G. (1799) Das Grasalchen, *Vibrio agrostis*. *Naturforscher* 28, 233–259.
- STEINER, G. (1914). Freilebende nematoden aus der schweiz. 1. Teil einer vorläufigen Mitteilung. *Archiv für Hydrobiologie und Planktonkunde* 9, 259-276.
- STEINER, G. (1920). Freilebende Suisswassernematoden aus peruanischen Hochgebirgsseen. *Revue Suisse de Zoologie* 28, 11-44.
- STEINER, G. (1926). Parasitic nemas on peanuts in South Africa. *Centralbl. Bakteriol.*, 67, 16-24.
- STEINER, G. (1929). *Diplogaster entomophaga* n. sp., a new *Diplogaster* (Diplogasteridae, Nematodes) found on a *Pamphilus stellatus* (CHRIST) (Tenthredinidae, Hymenoptera). *Zoologischer Anzeiger* 80, 143-145.
- STEINER, G. (1932). Some nemic parasites and associates of the mountain pine beetle (*Dendroctonus monticolae*). *Journal of agricultural Research* 45, 437-444.
- STEINER, G. (1934). *Procephalobus mycophilus* n. g., n. sp. (Cephalobina), a nematode living in the sclerotia of the fungus *Balansia claviceps*. *Proceedings of the Helminthological Society of Washington* 1, 56-58.
- STEINER, G. (1936a). Opuscula miscellanea nematologica. III. *Proceedings of the Helminthological Society of Washington* 3, 16-22.
- STEINER, G. (1936b). Opuscula miscellanea nematologica. IV. *Proceedings of the Helminthological Society of Washington* 3, 74-80.
- STEINER, G. (1938). Opuscula miscellanea nematologica VII. (I). Observations on nematodes associated with Irish potatoes grown in South Carolina. *Proceedings of the Helminthological Society of Washington* 5, 35-40.
- STERNBERG, P. W. (2005). Vulval development (June, 25 2005). *WormBook*, ed. The *C. elegans* Research Community, WormBook, doi/10.1895/wormbook.1.6.1, <http://www.wormbook.org>.
- STERNBERG, P.W. & HORVITZ, H.R. (1981). Gonadal cell lineages of the nematode *Panagrellus redivivus* and implications for evolution by the modification of cell lineage. *Developmental Biology* 88, 147-166.
- STOCK, S.P. (1993). *Micoletzkyia vidalae* n. sp. (Nematoda: Diplogasteridae), a facultative parasite of *Diabrotica speciosa* larvae (Coleoptera: Chrysomelidae) from Argentina. *Research and Reviews in Parasitology* 53, 109-112.

- STOCK, S.P., CAICEDO, A.M. & CALATAYUD, P.A. (2005). *Rhabditis (Oscheius) colombiana* n. sp. (Nematoda: Rhabditidae), a necromenic associate of the subterranean burrower bug *Cyrtomenus bergi* (Hemiptera: Cydnidae) from the Cauca Valley, Colombia. *Nematology* 7, 363-373.
- STOCK, S.P., CAMPBELL, J.F. & NADLER, S.A. (2001). Phylogeny of *Steinernema* Travassos, 1927 (Cephalobina: Steinernematidae) inferred from ribosomal DNA sequences and morphological characters. *Journal of Parasitology* 87, 877-889.
- STOCK, S.P. & NADLER, S.A. (2006). Morphology and molecular characterization of *Panagrellus* spp. (Cephalobina: Panagrolaimidae): taxonomic status and phylogenetic relationships. *Nematology* 8, 921-938.
- STROME, S. (2005). Specification of the germ line (July 28, 2005). *WormBook*. (Ed. The *C. elegans* Research Community, WormBook), doi/10.1895/wormbook.1.9.1, <http://www.wormbook.org>.
- STROME, S. & WOOD, W.B. (1983). Generation of asymmetry and segregation of germ-line granules in early *C. elegans* embryos. *Cell* 35, 15-25.
- SUDHAUS, W. (1974a). Zur Systematik, Verbreitung, Ökologie und Biologie neuer und wenig bekannter Rhabditiden (Nematoda). 1. Teil. *Zoologische Jahrbücher (Systematik)* 101, 173-212.
- SUDHAUS, W. (1974b). Zur Systematik, Verbreitung, Ökologie und Biologie neuer und wenig bekannter Rhabditiden (Nematoda). 2. Teil. *Zoologische Jahrbücher (Systematik)* 101, 417-465.
- SUDHAUS, W. (1976a). Vergleichende Untersuchungen zur Phylogenie, Systematik, Ökologie, Biologie und Ethologie der Rhabditidae (Nematoda). *Zoologica* 125, 1-229.
- SUDHAUS, W. (1976b). Nomenklatorische Bemerkungen Über arten und gattungen der unterfamilie Rhabditinae sensu lato (Rhabditidae: Nematoda). *Nematologica* 22, 49-61.
- SUDHAUS, W. (1978). Systematik, Phylogenie und Ökologie der holzbewohnenden Nematoden-Gruppe *Rhabditis (Mesorhabditis)* und das Problem "geschlechtsbezogener" Artdifferenzierung. *Zoologische Jahrbücher (Systematik)* 105, 399-461.
- SUDHAUS, W. (1980). Systematisch-phylogenetische und biologisch-ökologische Untersuchungen an *Rhabditis- (Poikilolaimus-)* Arten als Beitrag zur Rassenbildung und Parallelevolution bei Nematoden. *Zoologische Jahrbücher (Systematik)* 107, 287-343.
- SUDHAUS, W. (1985). Revision der Untergattung *Teratorhabditis* von *Rhabditis* (Nematoda) und Beschreibung einer neuen Art. *Zoologische Jahrbücher (Systematik)* 112, 207-224.
- SUDHAUS, W. (1991). Check list of species of *Rhabditis* sensu lato (Nematoda: Rhabditidae) discovered between 1976 and 1986. *Nematologica* 37, 229-236.
- SUDHAUS, W. (1993). Die mittels symbiontischer Bakterien entomopathogenen Nematodengattungen *Heterorhabditis* und *Steinernema* sind keine Schwestertaxa.. *Verhandlungen der Deutschen Zoologischen Gesellschaft* 86, 146.
- SUDHAUS, W. (1994). *Rhabditis (Oscheius) guentheri* sp. n., an unusual species with reduced posterior ovary, with observations on the *Dolichura* and Insectivora groups (Nematoda: Rhabditidae). *Nematologica* 40, 508-533.
- SUDHAUS, W. & FITCH, D. (2001). Comparative studies on the phylogeny and systematics of the Rhabditidae (Nematoda). *Journal of Nematology* 33, 1-70.

- SUDHAUS, W. & FÜRST VON LIEVEN, A. (2003). A phylogenetic classification of the Diplogastridae (secernentea, Nematoda). *Journal of Nematolode Morphology and Systematics* 6, 43-90.
- SUDHAUS, W. & KIONTKE, K. (1996). Phylogeny of *Rhabditis* subgenus *Caenorhabditis* (Rhabditidae, Nematoda). *Journal of Zoological Systematics and Evolutionary Research* 34, 217-233.
- SUDHAUS, W. & KIONTKE, K. (2007). Comparison of the cryptic nematode species *Caenorhabditis brenneri* sp. n. and *C. remanei* (Nematoda: Rhabditidae) with the stem species pattern of the *Caenorhabditis Elegans* group. *Zootaxa* 1456, 45-62.
- SUDHAUS, W. & REHFELD, K. (1990). *Diplogaster coprophilus* n. sp. and *D. affinis* n. sp. (Nematoda, Rhabditidae) from cow pats and related species, with notes on distribution, ecology and phylogeny. *Revue de Nematologie* 13, 51-65.
- SUDHAUS, W. & SCHULTE, F. (1986). Auflösung des Artenkomplexes *Rhabditis* (*Pelodera*) “*strongyloides*” (Nematoda) und Beschreibung zweier neuer Kryptischer Arten mit Bindung an Nagetiere. *Zoologische Jahrbücher (Systematik)* 113, 409-428.
- SUDHAUS, W. SCHULTE, F. & HOMINICK, W.M. (1987). A further sibling species *Rhabditis* (*Pelodera*) *strongyloides* (Nematoda): *Rhabditis* (*P.*) *cutanea* n. sp. from the skin of wood mice (*Apodemus sylvaticus*). *Revue de Nematologie* 10, 319-326.
- SULSTON, J., ALBERTSON, D.G. & THOMSON, J.N. (1980). The *Caenorhabditis elegans* male: postembryonic development of non-gonadal structures. *Developmental Biology* 78, 542–576.
- SULSTON, J. & HODGKIN, J. (1988). Methods. In *The nematode C. elegans* (Ed. Wood, W.B.). pp v 587-606. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
- SULSTON, J. E., SCHIERENBERG, E., WHITE, J.G. & THOMSON, J.N. (1983). The embryonic cell lineage of the nematode *Caenorhabditis elegans*. *Developmental Biology* 100, 64–119.
- SULSTON, J.E. & WHITE, J.G. (1980). Regulation and cell autonomy during postembryonic development of *Caenorhabditis elegans*. *Developmental Biology* 78, 577-597.
- SURYAWANSHI, M.V. (1971). *Alirhabditis indica* n. gen., n. sp. (Rhabditida: Alirhabditidae n. fam.). *Tawdenema indicum* n. gen., n. sp. and *Syedella aurangabadensis* n. gen., n. sp. (Diplogastridae) from Marathwada, India. *Nematologica* 17, 542-552.
- SWART, A., MEYER, J. A. & HEYNS, J. (1989). Descriptions of one new and two known species of Teratocephalidae (Nematoda) from South Africa. *Phytophylactica* 21, 367-377.
- TABASSUM, K.A. & SHAHINA, F. (2002). *Oscheius maqbooli* n. sp. and observations on three known rhabditid species (Nematoda: Rhabditida) from sugarcane fields of Balochistan, Pakistan. *Pakistan Journal of Nematology* 20, 1-21.
- TABASSUM, K.A. & SHAHINA, F. (2008). *Oscheius andrassyi* n. sp. (Nematoda: Rhabditidae) with its key and embryonic and post-embryonic development from Jhang, Pakistan. *Pakistan Journal of Nematology* 26, 125-140.
- TABASSUM, K. A., SHAHINA, F., FIROZA, K. & SIDDIQI, M. R. (2005). Description, embryonic and post embryonic development of *Distolabrellus pakistanensis* sp. n. (Rhabditida: Mesorhabditinae), a new insect parasitic nematode from Pakistan. *Pakistan Journal of Nematology* 23, 9-20.

- TAHSEEN, Q., AHMAD, I. BILGRAMI, A. L. & AHMAD, W. (1992). A scanning electron microscope study on *Mononchoides fortidens* (Schuunnans Stekhoven, 1951) Taylor & Hechler, 1966. *Nematologica* 38, 296-303.
- TAHSEEN, Q., AHMAD, I. & JAIRAJPURI, M.S. (1992). Description and developmental biology of *Plectus zelli* n sp (Nematoda: Araeolaimida). *Fundamental and Applied Nematology* 15, 503-510.
- TAHSEEN, Q., AHMAD, I. & JAIRAJPURI, M.S. (1993). SEM observations and developmental biology of *Tobrilus paludicola* Micoletzky, 1925 (Nematoda: Enoplida). *Nematologica* 39, 512-520.
- TAHSEEN, Q., AHMAD, I. & JAIRAJPURI, M.S. (1994). Embryogenesis and gonad development in *Chromadorita gracilis* and effect of Vitamin E on its development. *Afro-Asian Journal of . Nematology*. 4, 79-83.
- TAHSEEN, Q., AHMAD, I. & JAIRAJPURI, M.S. (1995). Two new species of Plectidae from India (Nematoda: Araeolaimida). *Fundamental and Applied Nematology* 18, 471-477.
- TAHSEEN, Q., AHMAD, I. & JAIRAJPURI, M.S. (1999). Observations on three species of the sub family Acrobelinae (Nematoda: Cephalobidae) from India. *Nematology* 1, 527-537.
- TAHSEEN, Q., HUSSAIN, A. & KHAN, R. (2007). Descriptions of three new and a known species of *Prismatolaimus* de Man 1880 (Nematoda: Enoplida) from freshwater habitats in India. *Journal of Nematode Morphology and Systematics* 9, 109-125.
- TAHSEEN, Q., HUSSAIN, A., SULTANA, R. & KHAN, R. (2009). Descriptions of the Indian species of *Poikilolaimus* Fuchs, 1930 (Nematoda: Rhabditidae) with *P. jodhpurensis* (Khera, 1969) Sudhaus, 1980 revisited and a brief discussion on relationship with other congeners. *Journal of Nematode Morphology and Systematics* 12, 27-40.
- TAHSEEN, Q., HUSSAIN, A., TOMAR, V., SHAH, A.A. & JAIRAJPURI, M.S. (2004). Description of *Metarhabditis andrassyana* gen. n., sp. n. (Nematoda: Rhabditidae) from India. *International Journal of Nematology* 14, 163-168.
- TAHSEEN, Q. & JAIRAJPURI, M.S. (1988). Description and developmental biology of *Teratorhabditis andrassyi* n. sp. (Nematoda: Rhabditida). *Revue de Nematologie* 11, 333-342.
- TAHSEEN, Q., JAIRAJPURI, M.S. & AHMAD, I. (1990). The life cycle of *Mononchoides fortidens* (Nematoda: Diplogastroidea) with emphasis on gonad development. *Nematologica* 36, 440-447.
- TAHSEEN, Q., JAIRAJPURI, M.S. & AHMAD, I. (1991a). Observations on the embryonic and post-embryonic development of *Diploscapter orientalis* (Nematoda: Rhabditida). *Revue de Nematologie* 14, 251-260.
- TAHSEEN, Q., JAIRAJPURI, M.S. & AHMAD, I. (1991b). Some observations on the developmental biology of *Cephalobus parvus* (Nematoda: Cephalobidae). *Nematologia Mediterranea* 19, 29-33.
- TAHSEEN, Q., JAIRAJPURI, M.S. & AHMAD, I. (1995). Development of Gonads in Plant and Soil Nematodes – A Review. *Proceedings of Indian National Science Academy* 61, 15-22.
- TAHSEEN, Q. & NISA, S. (2006). Embryology and gonad development in *Oscheius shamimi* sp. n. (Nematoda: Rhabditida). *Nematology* 7, 1-11.

- TAHSEEN, Q. SHAH, A.A., KHAN, R., SULTANA, R., HUSSAIN, A. & AHMAD, I. (2009). Biogeography and variations in allopatric populations of *Distolabrellus veechi* Anderson, 1983 (Nematoda: Rhabditidae). *Nematology* 11, 815-826.
- TAHSEEN, Q., SIDDIQI, M.R. & ROWE, J. (2002). Study on species of *Diploscapter* Cobb (Nematoda: Rhabditida) from India including *D. indicus* sp. n. *International Journal of Nematology* 12, 183-188.
- TAHSEEN, Q., SULTANA, R. & KHAN, R. (2007). Species of *Teratorhabditis* (Osche, 1952) Dougherty, 1953 (Nematoda: Rhabditida) from sewage and manure samples in India including a discussion on relationships within the genus. *Hydrobiologia* 583, 127-140.
- TAYLOR, D.P. (1964). *Butlerius monhystera* (Nematoda, Diplogasterinae), a new species of predaceous nematodes from Illinois. *Proceedings of the Helminthological Society of Washington* 31, 129-132.
- TAYLOR, D.P. & HECHLER, H.C. (1966). *Mononchoides fortidens* n. comb. and the transfer of *Mononchoides obscuridens* to the genus *Diplogaster*. *Nematologica* 12, 345.
- TAYLOR, T.M., BALDWIN, J.G. & MUNDO-OCAMPO, M. (2004). *Paracrobeles mojavicus* sp. n. (Nematoda: Cephalobidae) from the Mojave Desert, California. *Journal of Nematode Morphology and Systematics* 6, 151-160.
- THOMAS, P.R. (1965). Biology of *Acrobeles complexus* cultivated on agar. *Nematologica*, 11: 395-408.
- THORNE, G. (1925). The genus *Acrobeles* von Linstow, 1887. *Transactions of the American Microscopical Society* 44, 171-210.
- THORNE, G. (1937). A revision of the nematode family Cephalobidae Chitwood and Chitwood, 1934. *Proceedings of the Helminthological Society of Washington* 4, 1-16.
- THORNE, G. (1939). A monograph of the nematodes of the superfamily Dorylaimoidea. *Capita Zoologica* 8, 1-261.
- THORNE, G. (1949). On the classification of the Tylenchida, new order (Nematoda, Phasmidia). *Proceedings of the Helminthological Society of Washington* 16, 37-73.
- THORNE, G. (1961). *Principles of Nematology*. New York: McGraw-Hill Company, Inc. 553p.
- THORNE, G. & SWANGER, H.H. (1936). A monograph of the nematode genera *Dorylaimus* Dujardin, *Aporcelaimus* n. g., *Dorylaimoides* n. g., and *Pungentus* n. g. *Capita Zoologica* 6, 1-223.
- TIMM, R.W. (1956). Marine nematodes from the Bay of Bengal.1. Phasmidea. *Journal Bombay Natural History Society* 54, 87-90.
- TIMM, R.W. (1957). *Pterygorhabditis*, a remarkable new genus of soil nematodes. *Nematologica* 2, 68-71.
- TIMM, R.W. (1959). *Cheilorhabditis* and *Odontorhabditis*, two new genera of soil nematodes allied to *Rhabditis*. *Nematologica* 4, 198-204.
- TIMM, R.W. (1960). *Brevibucca punctata* n. sp. and *Macrolaimus natator* n. sp., new soil nematodes from East Pakistan. *Biologia* 6, 252-256.

- TIMM, R.W. (1960). *Cheilorhabditis* and *Odontorhabditis*, two new genera of soil nematodes allied to *Rhabditis*. *Nematologica* 4, 198-204.
- TIMM, R.W. (1961a). *Prodontorhabditis* n. gen. (Rhabditidae, Prodontorhabditinae n. subfam.), a new soil nematode from East Pakistan. *Proceedings of the Helminthological Society of Washington* 28, 115-117.
- TIMM, R.W. (1961b). Five species of *Diplogaster* (Nematoda, Diplogasteridae) from East Pakistan. *Pakistan Journal of Biology and Agricultural Science* 4, 1-6.
- TIMM, R.W. (1963). Marine nematodes of the suborder Monhysterina from the Arabian sea at Karachi. *Proceedings of Helminthological Society Washington* 30, 34-39.
- TIMM, R.W. (1969). *Megodontolaimus* new genus (Nematoda: Chromadoridae) with description of two new species. *Proceedings of the Helminthological Society of Washington* 36, 36-39.
- TIMM, R.W. (1971). Antarctic soil and freshwater nematodes from the McMurdo Sound Region. *Proceedings of the Helminthological Society of Washington* 38, 42-52.
- TRANTAPHYLLOU, A.C. (1963). Oogenesis in the root-knot nematode *Meloidogyne javanica*. *Nematologica* 7, 105-113.
- TRANTAPHYLLOU, A.C. (1971). 'Genetics and Cytology' in Plant Parasitic Nematodes, Vol. II, B. M. Zuckerman, W. F. Mai and R.A. Rhode (eds), Academic Press, New York and London.
- TRANTAPHYLLOU, A.C. (1981). Oogenesis and the chromosome of the parthenogenetic root-knot nematode *Meloidogyne incognita*. *Journal of Nematology* 13, 95-104.
- TRANTAPHYLLOU, A.C. (1985). Gametogenesis and the chromosomes of *Meloidogyne nataliei*: not typical of other root-knot nematodes. *Journal of Nematology* 17, 1-5.
- TRANTAPHYLLOU, A.C. & HIRSCHMANN, H. (1962). Oogenesis and mode of reproduction in the soyabean cyst nematode, *Heterodera glycines*. *Nematologica* 7, 235-241.
- TRANTAPHYLLOU, A.C. & HIRSCHMANN, H. (1964). Reproduction in plant and soil nematodes. *Annual Review of Phytopathology* 2, 57-80.
- TRANTAPHYLLOU, A.C. & HIRSCHMANN, H. (1966). Gametogenesis and reproduction in the wheat nematode, *Anguina tritici*. *Nematologica* 12, 437-442.
- TSALOLIKHIN, S.Y. (1972). [New species of free-living nematodes from the Baikal lake.] (In Russian.) *Zoologicheskii Zhurnal*, 51: 1559-1563.
- TSALOLIKHIN, S.Y. (1980). Free-living nematodes of Lake Baikal *Novosibirsk, USSR; Nauka*, 120 p (in Russian).
- VAN BENEDEN, É. (1883). Recherches sur la maturation de l'oeuf, la fécondation et la division cellulaire. *Archives de Biologie* 4, 265-640.
- VAN BENEDEN, É. & NEYT, A. (1887). Nouvelles recherches sur la fécondation et la division mitotique chez l'Ascaride mégalocephale. *Bulletin et Memoires de l'Academie Royale de Medecine de Belgique* 14, 215-295.
- VAN DEN BERG, E., SWART, A. & HEYNS, J. (1984). *Elaphonema messinae* sp. n. from the northern Transvaal (Nematoda: Rhabditida). *Phytophylactica* 16, 321-326.

- VAN DER LINDE, W.J. (1938). A contribution to the study of nematodes. *Entomology Memoirs, Department of Agriculture and Forestry, Union of South Africa* 2, 1-40.
- VAN GUNDY, S.D. (1958). The life history of the citrus nematode *Tylenchulus semipenetrans* Cobb. *Nematologica* 3, 283-294.
- VANFLETEREN, J.R. (1973). Amino acid requirements of the free-living nematode *Caenorhabditis briggsae*. *Nematologica* 19, 93-99.
- VANGESTEL, S., HOUTHOOFD, W., BERT, W. & BORGONIE, G. (2008). The early embryonic development of the satellite organism *Pristionchus pacificus*: differences and similarities with *Caenorhabditis elegans*. *Nematology* 10, 301-312.
- VAN WEERDT, L.G. (1960). Studies on the biology of *Radopholus similis* (Cobb, 1893) Thorne, 1949. Part III. Embryology and post-embryonic development. *Nematologica* 5, 43-52.
- VERA, C.P., MAGGENTI, A.R. & LOWNSEBURY, B.F. (1984). Post-embryonic development of the red wood nematode, *Rhizonema sequoiae* (Nematoda: Heteroderidae). *Journal of Nematology* 16, 73-83.
- VIEIRA, P., RYSS, A., SANTOS, A.R. & MOTA, M. (2009). On the vulval morphology of some species of *Bursaphelenchus* (Nematoda: Parasitaphelenchinae). *Nematology* 11, 793-798.
- VINCIGUERRA, M.T. (1994). *Metacrolobus festonatus* gen. n. sp. n. and *Scottinema lindsayae* Timm, 1971 (Nematoda: Cephalobidae) from Sub Antarctic and Antarctic regions with proposal of the new subfamily Metacrolobinae. *Fundamental and applied Nematology* 17, 175- 180.
- VÖLK, J. (1950). Die Nematoden der Regenwürmer und aabesuchenden Käfer. *Zoologische Jahrbücher (Systematik)* 79, 1-70.
- VON LINSTOW, O.F.B. (1906). Neue und bekannte Helminthen. *Zoologische Jahrbucher Abteilung für Systematik* 24, 1-20.
- WACEKE, J.W., BUMBARGER, D.J., MUNDO-OCAMPO, M., SUBBOTIN, S.A. & BALDWIN, J.G. (2005). *Zeldia spannata* sp. n. (Nematoda: Cephalobidae) from the Mojave Desert, California. *Journal of Nematode Morphology and Systematics* 8, 57-67.
- WEINGÄRTNER, I. (1955a). Versuch einer Neuordnung der Gattung *Diplogaster* Schulze 1857 (Nematoda). *Zoologische Jahrbücher (Systematik)* 83, 248-317.
- WEINGÄRTNER, I. (1955b). Berichtigung zur Arbeit Versuch einer Neuordnung der Gattung “*Diplogaster* Schulze 1857 (Nematoda)”. *Zoologische Jahrbücher (Systematik)* 83, 638.
- WESSING, A. (1953). Biologische Beobachtungen bei der Zucht der Regenwurm-nematoden *Rhabditis anomala* P. Hertwig. *Zoologische Anzeiger* 151, 28-38.
- WHITE, J. (2000). Worm tales. *International Journal of Developmental Biology* 44, 39-42.
- WIESER, W. (1953). Die Beziehung zwischen Mundhöhlengestalt, Ernährungsweise und Vorkommen bei freilebenden marinen Nematoden. *Archiv Fur Zoologie* 4, 439-484.
- WIESER, W. (1954). Free-living marine nematodes. II. Chromadoroidea. *Acta Universitatis Lundensis, Sweden (N.F. 2)* 50, 1-148.

- WIESER, W. (1956). Free-living marine nematodes. III. Axonolaimoidea and Monhysteroidea. Report Lund University Chile Expedition. 1948-49. *Acta Universitatis Lundensis* (N.F.2) 52, 1-115.
- WOLF, N., HIRSH, D. & MCINTOSH, J.R. (1978). Spermatogenesis in males of the free-living nematode, *Caenorhabditis elegans*. *Journal of Ultrastructure Research* 63, 155-169.
- WOOD, F.H. (1973). Biology of *Aporcelaimellus* sp. (Nematoda: Aporcelaimidae). *Nematologica* 19, 528-537.
- WOOD, W.B. (1988). In "The nematode *Caenorhabditis elegans*" (Ed.. Wood, W.B). *New York, NY: Cold Spring Harbor Laboratory Press*. Pp 215-241.
- WOOMBS, M. & LAYBOURN-PARRY, J. (1984). Growth reproduction and longevity in nematodes from sewage treatment plants. *Oecologia* 64, 168-172.
- WU, J., DE LEY, P. & LIANG, Y. (2001). Description of new species of the genus *Tylopharynx* (Nemata: Diplogasteroidea), *T. clariamphida* sp. n., with a description of *T. foetida* (Bütschli, 1874) Goffart, 1930. *Journal of Nematology* 33, 83-90.
- YEATES, G.W. (1967a). Studies on nematodes from dune sands.2. Araeolaimida. *New Zealand Journal of Sciences* 10, 287-298.
- YEATES, G.W. (1967b). Studies on nematodes from dune sands.3. Oncholaimidae, Ironidae, Alaimidae and Mononchidae. *New Zealand Journal of Science* 10, 299-321.
- YEATES, G.W. (1967c). Studies on nematode from dune sands. 7. Monhysterida and Chromadorida. *New Zealand Journal of Science* 10, 785-801.
- YEATES, G.W. (1969). Three new Rhabditida (Nematoda) from New Zealand dune sands. *Nematologica* 15, 115-128.
- YEATES, G.W. (1970). The diversity of soil nematode faunas. *Pedobiologia* 10, 104-107.
- YEATES, G.W. (1988). Distribution of Plectidae (Nematoda: Araeolaimida) in pasture soils. *New Zealand Journal of Zoology* 15, 191-202.
- YUEN, P.H. (1965). Further observations on *Helicotylenchus vulgaris*, Yuen, *Nematologica* 11, 623-637.
- YUEN, P.H. (1971). Electron microscope studies on *Aphelenchoides blastophthorus* (Nematoda). I. The female genital tube. *Nematologica* 17, 1-12.
- YUKSEL, H.S. (1960). Observations on the life cycle of *Ditylenchus dipsaci* on onion seedlings. *Nematologica* 5, 289-296.
- ZEIDAN, A.B. & GERAERT, E. (1989). Free-living Nematodes from Sudan. *Nematologica* 35, 277-304.
- ZIEGLER, H.E. (1895). Untersuchungen iiber die ersten Entwicklungsvorgange der Nematoden. *Zeitschrift fur Wissenschaftliche Zoologie* 60, 351-410.
- ZOGRAF, J.K., BERT, W. & BORGONIE, G. (2008). The structure of the female reproductive system of nematodes from the genus *Steinernema* (Rhabditida: Steinernematidae). *Nematology* 10, 883-896.

- ZOJA, R. (1896). Untersuchungen über die Entwicklung der *Ascaris megalcephala*. *Archiv für mikroskopische Anatomie* 47, 218-260.
- ZULLINI, A. (1981). Three new species of Diplogastrids (Nematoda). *Bollettino del Museo Civico di Storia Naturale di Verona* 7, 395-404.
- ZULLINI, A. & LOOF, A.A. (1980). Systematic notes on some species of Diplogasteridae (Rhabditida). *Nematologica* 26, 17-26.
- ZÜR STRASSEN, O. (1892). *Bradynema rigidum*. *Sieb Zeitschrift für Wissenschaftliche Zoologie* 54, 655-747.
- ZÜR STRASSEN, O. (1896). Embryonalentwicklung der *Ascaris megalcephala*. *Archiv für Entwicklungsmechanik der Organismen* 3, 27-105.