

## ACKNOWLEDGEMENTS

I am extremely indebted and thankful to Dr. N.C. Chattopadhyay, Reader, University College of Agriculture, Calcutta University for his valuable guidance; Head, University College of Agriculture, Calcutta University for granting laboratory facilities; Dr. N. Samajpati and Miss S. Maitra, Department of Botany for their valuable help in gel scanning; Sri N. Roychoudhuri, Indian Association for Cultivation of Science for his technical help in scanning electron microscopy; Dr. P.K. Mukherjee, Reader, Department of Botany, Calcutta University in interpretation of scanning electron micrographs; Drs. H. Sen and A.K. Bal of Central Soil Salinity Research Institute, Station Canning for their kind cooperation in conducting field experiment and Prof. K.R. Samaddar of Kalyani University for his helpful suggestions.

I am also thankful to Dr. P.R. Dasgupta, Technical Manager, Sutton & Sons for liberal supply of authentic seeds of tomato cultivars; Prof. R.N. Basu and members of the Ph.D. Committee for their valuable advice during the course of the works and Mr. S. Ghosh for typing this thesis.

I shall be failing in my duties if I do not mention the encouragement and constant help provided by Dr. A.K. Das, Mr. A.K. Pathak, Mr. S.K. Pal and my family members and close friends.

*Amit Kumar Chakraborti*  
(Amit K. Chakraborti).