

## REFERENCES

- Ashley, T. 1972. Zygote shrinkage and subsequent development in some Hibiscus hybrids. Planta 108: 303-317.
- Borssum, W.J. Van. 1966. Malesian Malvaceae revised. Blumea 14: 1-251.
- G.
- Chennaveeraiah, M.S. and Subbarao, 1965. Chromosome number and karyomorphology in Hibiscus cannabinus and H. sabdariffa. Karnak. Univ. 9: 78-87.
- Chizaki, Y. 1934. Breeding of a new interspecific hybrid between H. esculentus L. and H. manihot L. Proc. Crop. Sci. Soc. Japan 6: 164-172.
- \*Cleghorn, G.A. 1872. In Roe, Sr. M.J. 1961. A Taxonomic Study of the Indigenous Hawaiian Species of the Genus Hibiscus (Malvaceae) Pacific Sci. 15: 3-32.
- Dasgupta, A. and R.F. Bhatt. 1980. Polygraphic and Karyomorphological studies in the genus Urera Linn. J. Cytol. Genet. 15 (2): 123-127.
- \*Deshpande, R.B. 1938. Studies in Indian fibre plants. No.5. Further studies on the inheritance of certain characters in Hibiscus sabdariffa L. Indian J. Agri. Sci. 8: 229-243 (c.f. Chennaveeraiah and Subbarao, 1965).
- \*Fukusawa, C.A., F.F. Campos and J.M. Capinpin. 1960. Genetic aspects of the interspecific crosses between kenaf (Hibiscus cannabinus L.) and roselle (Hibiscus sabdariffa L.). Phillipine Agri. (Laguna) 44: 223-235.
- Gadwal, V.R., A.B. Joshi and R.D. Iyer. 1968. Interspecific hybrids in Abelmoschus through ovule and embryo culture. Indian J. Genet. Plant Breed. 28: 269-274.
- \*Gaertner, J. 1791. De fructibus et seminibus plantarum 2: 249-253 (c.f. Borssum, 1966).

- Gamble, J.S. 1921. Flora of the Presidency of Madras.  
H. Santapau (ed.) Botanical Survey of India,  
Calcutta.
- Goetz, S.G. 1982. Segregation of pollen exine features in  
Hibiscus section Furcaria interspecific hybrids  
Grana 21: 21-27.
- \*Hochreutiner, B.P.G. 1900. Revision du genre Hibiscus  
Annuaire Cons. Jard. Bot., Geneve 4: 23-191.
- \*-----, 1924. Genres nouveaux et genres discutees  
de la famille des Malvacees. Condollea 2: 79-90.
- Hogenboom, N.G. 1973. A model for incongruity in intimate  
partner relationships. Euphytica 22: 219-233.
- \*-----, 1975. Incompatibility and incongruity: two  
different mechanisms for non-functioning of intimate  
partner relationships. Proceedings of the Royal  
Society of London (B), 183: 361-375.
- Howard, H.W. 1939. The size of seeds in diploid and autote-  
traploid Brassica oleracea (L.). J. Genetics  
38: 325-339.
- \*Hu, S. 1955. Flora of China Fam. 153 (c.f. Borssum, 1966).
- Hutchinson, J. 1967. The genera of flowering plants. Vol.II.  
Oxford Univ. Press.
- Jambhale, N.D. and Y.S. Nerkar. 1982 a. Induced amphidiploidy  
in the cross Abelmoschus esculentus (L.) Moench. x  
A. manihot L. Medik esp. manihot. Genet. Agr. 36:  
19-22.
- and -----, 1982 b. Indirect selection  
criteria for isolation of induced polyploids in the  
Abelmoschus species hybrids. Cytologia 47: 603-607.

- and  
Joshi, A.B., M.W. Hardas. 1956. Allopoloid nature of okra  
Abelmoschus esculentus (L.) Moench. Nature (London)  
178 (4543): 1190.
- Kostoff, D. 1943. Cytogenetics of the genus Nicotiana. State  
Printing House, Sofia, Bulgaria.
- Kuwada, H. 1959. Studies on interspecific and intergeneric  
hybridization in Malvaceae. I. F<sub>1</sub> plants of H. mutabilis  
(L.) x H. moscheutos L. Jap. J. Breed. 9: 12-20.
- , 1960. Studies on the interspecific reciprocal hy-  
brids between Hibiscus coccineus and H. Moscheutos (No.1)  
Tech. Bull. Fac. Agri. Kagawa Univ. 12 (1): 1-8.
- , 1961 a. Studies on the interspecific hybrids bet-  
ween H. roseus and H. coccineus, H. Moscheutos and  
H. mutabilis. Tech. Bull. Fac. Agri. Kagawa Univ. 13  
(1): 1-7.
- , 1961 b. Studies on interspecific crossing bet-  
ween Abelmoschus esculentus (L.) Moench. and A. manihot  
Medic. and the various hybrids and polyploids derived  
from the above two species. Mem. Fac. Agri. Kagawa Univ.  
8: 1-91.
- , 1962 a. The F<sub>1</sub> hybrids obtained from Hibiscus  
mutabilis x H. coccineus. Tech. Bull. Fac. Agri. Kagawa  
Univ. 14 (1): 1-5.
- , 1962 b. Studies on the interspecific hybrids  
between Hibiscus coccinea and H. moschata. Jap. J. Breed.  
12 (2): 69-74.
- , 1963. Paper chromatographic studies on flower pig-  
ments of Hibiscus coccinea, H. moschata and F<sub>1</sub>, F<sub>2</sub> and F<sub>3</sub>  
hybrids obtained from the crosses between them. Jap. J.  
Breed. 13 (1): 48-51.
- , 1964 a. Studies on interspecific and intergeneric  
hybridization in the Malvaceae. VII. The newly artifici-  
ally raised amphidiploid plant named "Ai-Fuyo" (Hibiscus  
muta-moscheutos) obtained from the progeny of H. mutabilis  
x H. Moscheutos. Jap. J. Breed. 14 (1): 27-32.

- Kuwada, H. 1964 b. Studies on the cause and mechanism of poor plant growth of the  $F_1$  hybrids obtained from the crosses between H. coccineus (H. coccinea) and H. Moscheutos (H. moschata). Jap. J. Breed. 14 (2): 55-61.
- , 1966. A new amphidiploid plant named "Abelmoschus tubercular-esculentus" obtained from the progeny of the reciprocal crossing between A. tuberculatus and A. esculentus. Jap. J. Breed. 16 (1): 21-30.
- , 1967. The new amphidiploid plant named "Momiji-Fuyo" (Hibiscus muta-coccineus Kuwada) obtained from the progeny of H. mutabilis x H. coccineus. J. Jap. Soc. Hort. Sci. 36 (2): 89-94.
- , 1974.  $F_1$  hybrids of Abelmoschus tuberculatus x M. Manihot, with reference to the genome relationship. Jap. J. Breed. 24 (5): 1-4.
- , 1975. Interspecific hybridization between H. acetosella and H. cannabinus. SABRAO J. 7(1): 65-70.
- , 1977 a. Interspecific hybridization between Hibiscus acetosella and H. radiatus (Studies on Interspecific and Intergeneric Hybridization in the Malvaceae XVI). Jap. J. Breed. 27 (4): 51-56.
- , 1977 b.  $F_1$  hybrids obtained from Hibiscus cannabinus x H. radiatus, with reference to the genome relationship. Tech. Bull. Fac. Agri. Kagawa Univ. 28 (60): 19-22.
- and T. Mabuchi. 1977. Characteristics of  $F_1$  hybrids obtained from the crosses between H. asper, H. cannabinus and H. sabdariffa. Jap. J. Breed. 27 (3): 267-273.
- \*Linnaeus, C. 1753. Species Plantarum, London (c.f. Dasgupta and Bhatt, 1980).
- Martin, D.W. and M.Y. Menzel. 1971. Identification of the "Gondwanaland" genome found in South American tetraploids of Hibiscus sect. Furcaria. Abstr. Amer. J. Bot. 58: 465.

Martin, D.W. and M.Y. Menzel. 1972. Confirmations of homologous genomes in some African and New World species of Hibiscus sect. Furcaria. Abstr. ASB Bull. 19: 83.

Masters, M.T. 1875. Malvaceae. In J.D. Hooker. The Flora of British India Vol. I. L. Reeve and Co. London.

\*Medicus, F.K. 1787. Ueber einige kunstliche Geschlechter aus der Malven - Familie, dann der Klasse der Monadelphean, etc. Mannheim.

Menzel, M.Y. 1970. The Gondwanaland genomes of Hibiscus sect. Furcaria. Paper presented at the annual meeting of the Society for the study of Evolution Austin, Texas. Document No. ORG-3348 pp. 6.

-----, S.C. Goetz and W.C. Adamson. 1983 a. Some pieces of the African genome puzzle in Hibiscus sect. Furcaria (Malvaceae). Amer. J. Bot. 70 (2): 285-297.

-----, P.A. Fryxell and F.D. Wilson. 1983 b. Relationships among new World species of Hibiscus section Furcaria (Malvaceae). Brittonia 35(3): 204-221.

----- and D.W. Martin. 1970. Genome affinities of four African diploid species of Hibiscus sect. Furcaria. J. Heredity 61: 178-184.

----- and -----, 1971. Chromosome homology in some intercontinental hybrids in Hibiscus sect. Furcaria. Amer. J. Bot. 58: 191-202.

----- and -----, 1980. Evidence for the presence of an intercontinental genome in the Australian hexaploid alliance of Hibiscus sect. Furcaria. Austr. J. Bot. 28: 369-383.

----- and F.D. Wilson. 1961. Chromosomes and crossing behaviour of Hibiscus cannabinus, H. acetosella and H. radiatus. Amer. J. Bot. 48: 651-657.

Menzel, M.Y. and Wilson, F.D. 1963 a. Cytotaxonomy of twelve species of Hibiscus section Furcaria. Amer. J. Bot. 50: 262-271.

----- and -----, 1963 b. An allododecaploid hybrid of Hibiscus diversifolius and some related F<sub>1</sub> hybrids. J. Heredity 54: 55-60.

----- and -----, 1966. Hybrids and genome relations of Hibiscus gaudariffa, H. Meusei, H. radiatus and H. acetosella. Amer. J. Bot. 53: 270-275.

----- and -----, 1969. Genetic relationships in Hibiscus sect. Furcaria. Brittonia 21: 91-125.

\*Moench, C. 1794. Methodus Plantas Horti botanici et agri Marburgensis, e Stamimum situ describendi. 613-621 (c.f. Borssum, 1966).

Nair, P.G. and P. Kurischan. 1976. A spontaneous hybrid between Abelmoschus tuberculatus (Pal et Singh) and A. esculentus (L.) Moench. New Botanist III (1 & 2): 48-53.

\*Nishiyama, I. 1952. Polyploid-studies in the Brassiceae. I. The seed development in reciprocal crosses between diploid and tetraploid Raphanus. Mem. Res. Inst. Food Sci. Kyoto Univ. 3: 1-7. In Demerec, M. 1958 (ed.) Advances in Genetics. Vol. IX. Academic Press Inc. New York.

-----, and Y. Inamori. 1953. Polyploidy studies in the Brassiceae, III. Hybridization between diploid Brassica species ( $2n = 20$ ) and their autotetraploids. Mem. Res. Inst. Food Sci. Kyoto Univ. 5: 1-11. In Demerec, M. 1958 (ed.) Advances in Genetics. Vol. IX. Academic Press Inc. New York.

Pal, B.P., Singh, H.S. and V. Swarup. 1952. Taxonomic relationship and breeding possibilities of species of Abelmoschus related to Okra (Abelmoschus esculentus) Bot. Gaz. 113: 455-464.

Patil, J.S. 1966. The culture in vitro of immature embryos of okra (Abelmoschus esculentus). Indian J. Plant Physiol. 9: 59-65.

Rakshit, S.C. and B.C. Kundu. 1970. Revision of the Indian species of Hibiscus. Bull. Bot. Sur. India 12 (1-4): 151-175.

\*Sansome, E., S. Satina and A.F. Blakeslee. 1942. Disintegration of ovules in tetraploid-diploid and in incompatible crosses in Datura. Bull. Torrey Bot. Club 69: 405-420.

Sanyal, P. 1958 a. Embryological investigations in Hibiscus sabdariffa x cannabinus and their reciprocal crosses. Nature 181: 1352.

-----, 1958 b. Studies on the pollen tube growth in six species of Hibiscus and their crosses in vivo. Cytologia 23: 460-467.

-----, and B.C. Kundu. 1959. Cytological study of Hibiscus radiatus x H. cannabinus hybrid and genomic relationship of the two species. The Nucleus 11: 99-108.

Schumann, K.M. 1890. Malvaceae. In Engler, H.G.A. and K.A.E. Prantle. Die natuerlichen Pflanzenfamilien 3 (6): 30-53.

Sharma, M.S. 1967. Breeding procedures for Hibiscus. ICAR, New Delhi.

Skovsted, A. 1955. Chromosome numbers in the Malvaceae. I. J. Genetics 31(2): 263-296.

-----, 1944. Some hybridization experiments in the tribe Hibisceae. Compt. Rend. Trav. Lab. Carlsberg (Ser. Physiologique) 24 (1): 1-30.

Tachibana, Y. 1958. Studies on Hibiscus (III). An interspecific hybrid of H. mutabilis L. and H. moscheutos L. Engel Gakkai Zasshi J. of the Hort. Soc. 27: 201-206 (c.f. Wise and Menzel, 1971).

- Thakur, M.R. 1976. Inheritance of yellow vein mosaic in a cross of okra species Abelmoschus esculentus x A. manihot subspecies manihot. SABRAO J. 8 (1): 69-73.
- \*Teshima, T. 1933. Genetical and cytological studies in an interspecific hybrid of Hibiscus esculentus and H. manihot. J. Pac. Agric. Hokkaido Univ. 34: 1-156 (c.f. Menzel and Wilson, 1969).
- Thompson, W.P. 1930. Causes of differences in success of reciprocal interspecific crosses. Amer. Nat. 64: 407-421.
- Toxopeus, H.J. 1947. Preliminary account on the results of some species crosses in Hibiscus. Genetica 24: 90-92.
- \*Ustinov, E.I. 1949. A description of the interspecific hybrid of Hibiscus esculentus x H. manihot. Priroda (Nature) Moscow 6: 58-60.
- Wilson, F.D. and P.A. Fryxell. 1970. Meiotic chromosomes of Cienfuegosia species and hybrids and Hampea species (Malvaceae) Bull. Torr. Bot. Club. 97(6): 367-376.
- and M.Y. Menzel. 1967. Interspecific hybrids between kenaf (Hibiscus cannabinus) and roselle (H. sabdariffa). Euphytica 16: 33-44.
- Wise, D.A. 1973. Genome relationship of Hibiscus mutabilis. J. Heredity 64: 285-288.
- and M.Y. Menzel. 1971. Genetic affinities of the North American species of Hibiscus Sect. Trionum. Brittonia 23: 425-437.
- and -----, 1972. Taxonomic relationship of Hibiscus cubensis Rich., H. nivalis St. Hil., and H. lambertianus H.B.K. Taxon 21: 378-379.
-