

# References

---

- Anderson D, Prasad K, and Stewart R. 1995. Changes in isozyme Profiles of catalase, peroxidase and Glutathione reductase during acclimation to chilling in mesocotyls of maize seedlings. *Plant Physiology* 109: 1247 – 1257.
- Arslan, O., Temur, A., and Tozlu, I. (1997) Polyphenol oxidase from *Allium* sp. *J. Agric. Food Chem.*,45. p: 2861– 2863.
- Ashihara, H., Deng, W. W., Mullen, W., and Crozier, A. (2010). Distribution and biosynthesis of flavan-3-ols in *Camellia sinensis* seedlings and expression of genes encoding biosynthetic enzymes. *Phytochemistry*. 71. p: 559–66.
- Aucamp, J. P., Hara, Y., and Apostolides, Z. (2000). Simultaneous analysis of tea catechins, caffeine, gallic acid, theanine and ascorbic acid by micellar electrokinetic capillary chromatography. *J. Chromatogr. A.*, 876. p: 235-242.
- Ayaz, F. A., Demir, O., Torun, H., Kolcuoglu, Y., and Colak, A. (2008). Characterization of polyphenoloxidase (PPO) and total phenolic contents in medlar (*Mespilus germanica* L.) fruit during ripening and over ripening. *Food chemistry*. 106 (29). p: 291–298.
- Balasaravanan, T., Pius, P. K., Raj Kumar, R., Muraleedharan, N. and Shasany, A. K. (2003). Genetic diversity among south Indian tea germplasm (*Camellia sinensis*, *assamica* and *C. assamica* spp. *lasiocalyx*) using AFLP markers. *Plant Science*. 165.p: 365-372.
- Baritoux, O., Amiot, M. J., Richard, H., and Nicolas, J. J. (1991). Enzymatic browning of basil (*Ocimum basilicum* L.) studies on phenolic compounds and polyphenol oxidase. *Sciences des Aliments.*, 11. p: 49–62.
- Barua, D. N. (1965). Classification of the tea plant: species hybrids. *Two and a Bud*. 12, p: 13-27.
- Biswas, A. K., Biswas, A. K., and Sarkar, A. R. (1971). Biological and chemical factors affecting the evaluations of North East India plain teas. II. Statistical evaluation of the biochemical constituents and their effects on briskness,



- quality and cash valuation of black teas. *Journal of the Science and Food Agriculture*. 22.p: 191–198.
- Bokuchanva, M. A., and Skobeleva, N. I. (1980). The biochemistry of tea manufacture. *CRC Crit. Rev. Food Sci. Nutr.*, 12. p: 303-370.
- Bradford, M. (1976). A Rapid and Sensitive Method for the Quantitation of Microgram Quantities of Protein Utilizing the Principle of Protein-Dye Binding. *Analytical Biochemistry*, Vol. 72, pp. 248-254
- Bucheli, C. S., Dry, I. B. and Robinson, S. P. (1996). Isolation of a full-length cDNA encoding polyphenol oxidase from sugarcane, a C4 grass. *Plant Mol. Biol.*, 31 (6). p : 1233- 1238.
- Cano, M, P., Lobo, M. G., Ancos, B., and Galeazzi, M. (1996). Polyphenol oxidase from Spanish hermaphrodite and female papaya fruits (*Carica papaya*). 1996 *J. Agric. Food Chem.*, 44. p: 3075 – 3079.
- Cary, J. W., Lax, A. R., and Flurkey, W. H. (1992). Cloning and characterisation of cDNAs coding for *Vicia faba* polyphenol oxidase. *Plant Molecular Biology*. 20. p: 245 – 253.
- Chance, B and Maehly, A. C. (1955). Methods in Enzymology (Colowick, S. P., and KAPLAN, eds ), Academic Press, New York. Vol II. p: 764.
- Chazarra, S., García-Carmona, F., and Cabanes, J. (2001). Evidence for a tetrameric form of iceberg lettuce (*Lactuca sativa* L.) polyphenol oxidase: Purification and characterization. *J. Agric. Food Chem.*, 49.p: 4870-4875.
- Chen, L., and Yamaguchi, S. (2002). Genetic diversity and phylogeny of tea plant (*Camellia sinensis*) and its related varieties in the section Thea genus *Camellia* determined by randomly amplified polymorphic DNA analysis. *J. Hort. Sci. & Biotech.*, 77(6)p: 729-732.
- Chen, L., and Yamaguchi, S. (2005). RAPD markers for discriminating tea germplasms at the inter-specific level in China. *Plant Breeding*. 124. p: 404-409.
- Chen, L., and Zhou, Z. X. (2005). Variations of main quality components of tea genetic resources [*Camellia sinensis* (L.) O. Kuntze] preserved in the China National Germplasm Tea Repository. *Plant Foods for Human Nutrition*. 60. p: 31-35.



- Chen, L., Gao, Q., Chen, D. and Xu, C. (2005). The use of RAPD markers for detecting genetic diversity, relationship and molecular identification of Chinese elite tea genetic resources *Camellia sinensis* (L.) O. Kuntze preserved in a tea germplasm repository. *Biodiversity and Conservation*, 14. p: 1433-1444.
- Chen, L., Yu, F, L., and Tong, Q. Q. (2000). Discussions on phylogenetic classification and evolution of sect Thea. *J. Tea. Sci.*, 20. p:89-94.
- Chen, Y.L., Jiang, Y. M., Duan, J., Shi, J., Xue, S. & Kakuda, Y. (2010). Variation in catechin contents in relation to quality of ‘Huang Zhi Xiang’ Oolong tea (*Camellia sinensis*) at various growing altitudes and seasons. *Food Chemistry*, 119, 648–652.
- Chevalier, T., Rigal, D., Mbèguiè-A-Mbèguiè. D., Gauillard, F., Richard-Forget, F., and Fils-Lycaon, B. R. (1999). Molecular cloning and characterization of apricot fruit polyphenol oxidase. *Plant Physiol.*, 119. p: 1261-1269.
- Coggon, P., Moss, G. A., & Sanderson, G. W. (1973). Tea catechol oxidase: isolation, purification and kinetic characterisation. *Phytochemistry*, 12, 1947–1955.
- Dev Choudhury, M. N., and Bajaj, K. L. (1986). Seasonal variation of Water soluble sugars in tea shoot and their role in tea quality. *Two and A Bud*, 33. p: 33-38.
- Dev choudhury, M.N. and Goswami, M.R. 1983. A rapid method for the estimation of total polyphenolic matters in tea (*Camellia sinensis* L.). *Two and A Bud*, 30: 59-61.
- Dignum, J., Kerler, J., and Veroorte, R. (2001). D-glucosidase and peroxidase stability in crude enzyme extracts from green beans of *Vanilla planifolia* Andrews. *Phytochemical Analyzes.*, 12. p: 174–179.
- Dincer, B., Colak, A., Aydin, N., Kadioglu, A., and Guner, S. (2002). Characterization of polyphenoloxidase from medlar fruits (*Mespilus germanica* L., Rosaceae). *Food Chem.*, 77. p: 1–7.
- Dogan, M., Arslan, O., and Dogan, S. (2002). Substrate specificity, heat inactivation and inhibition of polyphenol oxidase from different aubergine cultivars. *International Journal of Food Science and Technology*. 37.p: 415–423.



- Dogan, S., and Dogan, M. (2004). Determination of kinetic properties of polyphenol oxidase from *Thymus* (*Thymus longicaulis* subsp. *Chaubardii* var *chaubardii*). *Food Chemistry*. 88.p: 69–77.
- Doyle, J. J., and J. L. Doyle, 1987: A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem. Bull.* 19, 11—15.
- Dry, I. B. and Robinson, S. P. 1994. Molecular cloning and characterization of grape berry polyphenol oxidase. *Plant Mol. Biol.*, 26. p: 495-502.
- Durmishidze, S.V. and Pruidze, G.N. 1980. Phenolases and peroxidases of tea plant and grapevine shoots. *Fiziol. Rast.*, 27: 1064-1071.
- Erat, M., Sakiroglu, H., and Kufrevioglu, O. I. (2006). Purification and characterization of polyphenol oxidase from *Ferula* sp. *Food chemistry*. 95. p: 503–508.
- Eungwanichayapant, P. D., and Popluechai, S. (2009). Accumulation of catechins in tea in relation to accumulation of mRNA from genes involved in catechin biosynthesis. *Plant Physiol Biochem.*, 47. p:94–7.
- Ezaki, B., Katsuhara, M., Kawamura, M., and Matsumoto, H. (2001). Different mechanisms of four aluminum (Al)-resistant transgenes for Al toxicity in *Arabidopsis*. *Plant Physiol.*, 127. p: 918-927
- Fujita S, Saari N, Maegawa M, Tetsuka T, Hayashi N, Tono T. Purification and properties of polyphenol oxidase from cabbage (*Brassica oleracea* L.). *J. Agric. Food Chem* 1995; 43: (113) 1138–1142.
- Gawlik-Dziki, U., Zlotek, U., and Swieca, M. (2008). Characterization of polyphenol oxidase from butter lettuce (*Lactuca sativa* var. *Capitata* L.). *Food Chem.*, 107. p: 129–135.
- Giovana C., Ceni, Eliana, M., Baldissera, Octa'vio, A. C., and Antunes, J., Oliveira, V., Dariva, C., de Oliveira, D. (2008). Oxidases from mate tea leaves (*Ilex paraguariensis*): extraction optimization and stability at low and high temperatures. *Bioprocess Biosyst Eng.*, 31. p: 541–550.
- Golbeck, J. H., and Cammarata, K. V. (1981). Spinach thylakoid polyphenol oxidase. Isolation, activation and properties of the native chloroplast enzyme. *Plant Physiology*. 67. p: 877–884.



- Gomez, K. A., Gomez, A. A. (1976). Statistical procedures for agricultural research with emphasis of rice. International Rice Research Institute, Los Banos, Philippines.
- Gomez, K.A. and Gomez, A.A. 1984. Statistical Procedures for Agricultural Research. John Wiley and Sons, New York. p. 20–30.
- Gomez-Lopez, V. M. (2002). Some biochemical properties of polyphenol oxidase from two varieties of avocado. *Food chemistry*. 77. p: 163–169.
- Gregory, R. P. F., and Bendall, D. S. (1966). Purification and some properties of polyphenol oxidase from tea. *Biochem. J.*, 101. p: 569-581
- Gulati, A. and Ravindranath, S. D. (1996). Seasonal variation in quality of Kangra tea (*Camellia sinensis* (L) O Kuntze) in Himachal Pradesh. *J. Sci. Food Agric.*, 71. p: 231-236.
- Hackett, C. A., Wachira, F. N., Paul, S., Powell, W. and Waugh, R. (2000). Construction of a genetic linkage map for *Camellia sinensis* (tea). *Heredity*. 85. p: 346- 355.
- Halder, J., Tamuli, P., and Bhaduri, A. N. (1998). Isolation and characterization of polyphenol oxidase from Indian tea leaf (*Camellia sinensis*). *Journal of Nutritional Biochemistry*. 9. p:75–80.
- Harbowy, M. E., and Balentine, D. A. (1997). Tea chemistry. *Crit. Rev. Plant Sci.*, 16. p: 415–480.
- Hedge, J.E. and Hofreiter, B.T. 1962. *In Carbohydrate chemistry.*, Academic Press, New York. p. 17.
- Hudson, J.B., Durairaj, J., Muraleedharan, N. (2002). Guidelines on Tea Culture in South India. Allied Publishers Private Limited, Chennai, India.
- ISO 14502–2: 2005. Tea–Method for determination of substances characteristics of green and black tea–Part–2: Content of catechins in green tea–Method using high-performance chromatography.
- Jaccard P (1908) Nouvelles recherches sur la distribution florale. *Bulletin Societe Vaudoise des Siences Naturelles*. 44: 223–270.
- Jain, C. and Takeo, T. 1984. A review. Enzymes of tea and their role in tea making. *J. Food Biochem.*, 8: 243-279.



- Jain, J. C., Ullah, M. R. and Dev choudhury, M. N. (1979). L-Phenylalanine ammonia lyase in tea leaf. *Two and A Bud.*, 26. p: 67-69.
- Jayaganesh, S. 2009. Chemistry of magnesium in nutrition of tea. Ph.D. Thesis submitted to Bharathiar University, Coimbatore, Tamil Nadu, India. p. 127 and 129.
- Jeyaramraja, P. R., Rajkumar, R., and Jayakumar, D. (2003). Soil moisture stress-induced alterations in bioconstituents determining tea quality. *J. Sci. Food Agric.*, 83. p: 1187-1191.
- Jiang, Y. M., Zauberman, G., and Fuchs, Y. (1997). Partial purification and some properties of polyphenol oxidase extracted from litchi fruit pericarp. *Postharvest Biology and Technology*. 10. p: 221–228.
- Jibu Thomas, Raj Kumar, R., and Mandal, A. K. A. (2006). Metabolite profiling and characterization of somaclonal variants in tea (*Camellia spp.*) for identifying productive and quality accession. *Phytochemistry*. 67. p: 1136–1142.
- Jibu Thomas. 2008. '*Bio-markers*' to elucidate the desirable traits for quality and yield in tea (*Camellia spp.*) germplasm. Ph.D. thesis. Bharathiar University, Coimbatore, India. p: 76.
- Joseph Lopez, S., Jibu Thomas, Pius, P. K., Raj Kumar, R., and Muraleedharan, N. (2005). A reliable technique to identify superior quality clones from tea germplasm. *Food Chemistry*. 91. p: 771-778
- Jyotsnabaran, Halder, Prodip, Tamuli, and Bhaduri A. N. (1998). Isolation and characterization of polyphenol oxidase from Indian tea leaf (*Camellia sinensis*). *Nutritional Biochemistry*. 9. p: 75-80.
- Kaundun, S. S., and Park, Y. G. (2002). Genetic structure of six Korean tea populations as revealed by RAPD-PCR markers", *Crop Sci.*, 42. p: 594-601.
- Kaundun, S. S., Zhyvoloup, A., and Park, Y. G (2000). Evaluation of the genetic polymorphism among elite tea (*Camellia sinensis* var. *sinensis*) accessions using RAPD markers. *Euphytica*. 115. p: 7—16.
- Klabunde, T., Eicken, C., Sacchettini, J.C., and Krebs, B. (1998). Crystal structure of a plant catechol oxidase containing a dicopper center. *Nature Structural Biology*. 5 (12). p: 1084 – 1090.



- Kottur. 2010. Influence of pigments and phenolics on the quality and flavour of black teas. Ph.D. thesis. Bharathiar University, Coimbatore, India.
- Kowalski, S.P., Eannetta, N.T., Hirzal, A. T., and Steffens, J.C. (1992). Purification and characterization of polyphenol oxidase from glandular trichomes of *solanum berthaultii*. *Plant physiol.*,100.p: 677-684.
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, 227. 680–685.
- Lai, J. A., Yang, W. C. and Hsiao, J. Y. 2001. An assessment of genetic relationships in cultivated tea clones and native wild tea in Taiwan using RAPD and ISSR markers. *Botanical Bulletin of Academia Sinica.*, 42.p: 93-100.
- Lamkey, K. R., and Lee, M. (1993). Quantitative genetics, molecular markers, and plant improvement. In: B. C. Imrie and J. B. Hacker (ed) *Focused plant improvement : Towards responsible and sustainable agriculture*. Proc. 10th Australian Plant Breeding Conf., Gold Coast, 18-23 April 1993. Organizing committee, Australian Convention and Travel Service: Canberra. P: 104-115.
- Li Guo, Ying Ma, John, S., Sophia Xue. (2009). The purification and characterisation of polyphenol oxidase from green bean (*Phaseolus vulgaris* L.) *Food Chemistry*. 117. p: 143–151.
- Liang, G, L., Zhou, C, Q., Lin, M, J., Chen, J, Y., and Liu, J. S. (1994). Karyotype variation and evolution of sect *Thea* in Guizhou. *Acta. Phytotaxonomica. Sinica.*, 32. p: 308-315.
- Liang, Y. W., Ping, L .P., Yu, S. L., and Hai-Hang, L. (2010). Cloning, microbial expression and structure–activity relationship of polyphenol oxidases from *Camellia sinensis*. *Journal of Biotechnology*. 145. p : 66-72.
- Liang, Y.R., Lu, J.L., Zhang, L.Y., Wu, S. & Wu, Y. (2005). Estimation of tea quality by infusion colour difference analysis. *Journal of the Science of Food and Agriculture*, 85. 286–290.
- Lin, Y. L., Juan, I. M., Chen, Y. L., Liang, Y. C. and Lin, J. K. (1996). Composition of polyphenols in fresh tea leaves and associations of their oxygen-radical-absorbing capacity with antiproliferative actions in fibroblast cells. *J. Agric. Food Chem.*, 44. p: 1387-1394.



- Lineweaver, H., & Burk, D. (1934). The determination of enzyme dissociation constant. *Journal of American Chemical Society*, 56, 658–661.
- Luck, H. (1974). Catalase, In: Bergmeyer (Ed), *Methods in Enzymatic analysis I*. Academic Press, New York. p: 885
- Magoma, G. N., Wachira, F. N., Obanda, M., Imbuga, M., and Agong, S. G. (2000). The use of catechins as biochemical markers in diversity studies of tea (*Camellia sinensis*). *Genetic Resources and Crop Evolution*. 47(2).p: 107–114.
- Marimuthu, S. (2000). Role of enzymes in tea processing. *Final Technical Report submitted to National Tea Research Foundation, Kolkata, India*. p: 3.
- Marimuthu, S., and Gunasekar, M. (1998). Tea manufacture: Role of enzymes. *Project Report submitted to NTRF., Tea Board (India)*. p. 12- 13 and 29.
- Mariya john, K. M., Sasikumar, R., Balasubramanian, M., Saravanan, M., and RajKumar, R. (2003). Influence of light on catechin biosynthesis in tea. *Tea*. 24.p: 80–86.
- Marri, C., Frazzoli, A., Hochkoepler, A., and Poggi, V. (2003). Purification of a polyphenol oxidase isoform from potato (*Solanum tuberosum*) tubers. *Phytochemistry*. 63. p: 745-752.
- Martin-Diana, A. B., Rico, D., Barry-Ryan, C., Mulcahy, J., Frias, J., and Henehan, G. T. M. (2005). Effect of heat shock on browning-related enzymes in minimally processed iceberg lettuce and crude extracts. *Bioscience, Biotechnology and Biochemistry*. 69(9). p: 1677– 1685.
- Mayer, A. M. (2006). Polyphenol oxidases in plants and fungi: Going places? A review. *Phytochemistry*. 67. p: 2318 – 2331.
- Mazzafera, P., and Robinson, S. P. (2000). Characterization of polyphenol oxidase in coffee. *Phytochemistry*. 55. p: 285–296.
- McCord, J.M. and I. Fridovich. 1969. Superoxide dismutase: anenzymic function for erythrocuprein (hemocuprein). *J. Biol. Chem*. 244: 6049- 6055.
- McDowell, I., Feakes, J. and Gay, C. 1991. Phenolic composition of black tea liquors as a means of predicting price and country of origin. *J. Sci. Food Agric.*, 55(4): 627-641.





- Melchinger, A. E. (1993). Use of RFLP markers for analyses of genetic relationships among breeding materials and prediction of hybrid performance. Proc. of the First Int. *Crop Sci. Cong.*
- Molla, M. M. (1992). Effect on polyphenol oxidase activity in extracts of tea leaf prepared by maceration at different temperature. Sri Lankan Journal of Tea Science. 61. p: 15-19
- Mondal, T. K. (2002). Assessment of genetic diversity of tea (*Camellia sinensis* (L.) O Kuntze) by inter-simple sequence repeat polymerase chain reaction. *Euphytica*. 128. p: 307-315.
- Moore, S. and Stein, W.H. 1948. In: *Methods in Enzymol* (Eds. Colowick, S.P. and Kaplan, N.D.), Academic Press, New York, 3: p. 468.
- Muthumani, T., and Senthil Kumar, R. S. (2004). Quality evaluation of CTC black teas (*Camellia sinensis* (L) O. Kuntze) from Anamallais region of south India, *J. Plantn. Crops.*, 32(suppl.). p: 417-421.
- Nagai, T., and Suzuki, N. (2001). Partial purification of polyphenol oxidase from chinese cabbage. *Journal of Agricultural and Food Chemistry*. 49. p: 3922-3926.
- Nei, M .and Li, W. (1979). Mathematical model for studying genetic variation in terms of restriction endonuclease, *Proc Natl Acad Sci USA*, 76 . 5296-5273.
- Nkya, E., Kouno, C., Li, Y.J., Yang, C.P., Hayashi, N., and Fujita, S. (2003). Purification and characterization of polyphenol oxidase from garland chrysanthemum (*Chrysanthemum coronarium* L.). *J. Agric. Food Chem.*, 51. p: 5467–5471.
- Oberly, L.W. and D.R. Spitz. 1985. Nitroblue Tetrazolium. In R.A. Greenwald (ed.), *CRC Handbook of Methods for Oxygen Radical Research*. CRC Press, Boca Raton, pp. 217- 220.
- Okinda Owuor, P., Obanda, M., Hastings, E., Nyirenda, Nicholas, I. K., Mphangwe, Louwrance, P., Wright Zeno Apostolides. (2006). The relationship between some chemical parameters and sensory evaluations for plain black tea (*Camellia sinensis*) produced in Kenya and comparison with similar teas from Malawi and South Africa. *Food Chemistry*. 97. p: 644–653



- Othieno, C.O. (1992). Soils. In: Willson, K.C., Clifford, M.N. (Eds.), *Tea: Cultivation to Consumption*. Chapman and Hall, London. P: 137–170.
- Owuor, P. O., and Obanda, M. (1993). Fermentation temperature and duration effects on black tea quality. *Tea.*, 14. p: 21-30.
- Park, E. Y., and Luh, B. S. (1985). Polyphenol oxidase of Kiwifruit. *Journal of Food Science*. 50. p: 679–684.
- Park, J. S., Kim, J. B., Haha, B. S., Kim, K. H., Ha, S. H., Kim, J. B., and Kim, Y. H. (2004). EST analysis of genes involved in secondary metabolism in *Camellia sinensis* (tea), using suppression subtractive hybridization, *Plant Sci.*, 166. p: 953-961.
- Parry, A. D., Goodsall, C. W., and Safford, D. S. (1996). The involvement of polyphenol oxidase and peroxidase in the oxidation of polyphenols during the manufacture of black tea. *Polyphenolic communications*. 96. (039).
- Paul, S., Wachira, F. N., Powell, W. and Waugh, R. (1997). Diversity and genetic differentiation among populations of Indian and Kenyan tea (*Camellia sinensis* (L.) O. Kuntze) revealed by AFLP markers. *Theor Appl Genet.*, 94. p: 255-263.
- Pruidze, G. N., Mchedlishvili, N. I., Omiadze, N. T., Gulua, L., and Pruidze, k. (2003). Multiple forms of phenol oxidase from Kolkhida tea leaves (*Camellia Sinensis* L.) and Mycelia Sterila IBR35219/2 and their role in tea production. *Sci. Food Res. Intern.*, 36. p: 587-595.
- Punyasiri, P. A. N., Herath, N.L. and Amarakoon, A. M. T. (1990). An improved procedure for the isolation of flavanols from tea leaves. *Sri Lankan J. Tea Sci.*, 59(1). p: 24-28.
- Raj Kumar, R., Manivel, L. and Marimuthu, S. (1993). Evaluation of clonal characteristics in mature tea. *Journal of Plantation Crops*. 21 (suppl). p: 241-247.
- Ramasamy, V., and Raju, K. (1993). Tea fermentation-constraints and advances. In Mulky, M.J. and Sharma, V.S. (Eds.), *Tea culture processing and marketing*, New Delhi Oxford and IBH Publishing Co. Pvt. Ltd. P: 147-157.
- Ramaswamy, S. 1986. Improving tea quality in south India. UPASI Tea Scientific Department Bulletin, 41: 12–24.



- Ravichandran, R. (2002). Carotenoid composition, distribution and degradation to flavour volatiles during black tea manufacture and the effect of carotenoid supplementation on tea quality and aroma. *Food Chem.*, 78.p: 23-28.
- Ravichandran, R., and Parthiban, R. (1998). Changes in enzyme activities (polyphenol oxidase and phenylalanine ammonia lyase) with type of tea leaf and during black tea manufacture and effect of enzyme supplementation of dhool on black tea quality. *Food Chem.*, 62(3). p: 277–281.
- Raymond, J., Rakariyatham, N., and Azanza, J. L. (1993). Purification and some properties of polyphenoloxidase from sunflower seeds. *Phytochemistry*. 34. p: 927-931.
- Roberts, E. A. H. (1962). Economical importance of food substances: Tea fermentation. In Geissmann, T.A. (Ed.), *The chemistry of flavonoids compounds*. Pergamon, Oxford. P: 468-512.
- Roberts, G.R. and Fernando, R.S.S. (1981). Some observations on the correlation of polyphenol content to the quality of tea clones. *Tea Quart.*, 50: 30-34.
- Robertson, A., and Bendall, D.S. (1983). Production and HPLC analysis of black tea theaflavins and thearubigins. *Phytochemistry*. 22. p: 883-887.
- Robinson, S.P., and Dry, I.B. (1992). Broad bean leaf polyphenol oxidase is a 60-kilodalton protein susceptible to proteolytic cleavage. *Plant Physiology*. 99. p: 317– 323.
- Rohlf , FJ. (2002) NTSYS-pc:numerical taxonomy system ver.2.1.Exeter publishing Ltd., Setauket, Newyork.
- Ronald, S. K., and Ronald, S. (1991). Beverages and chocolate. In *Pearson's composition and analysis of foods*, 9<sup>th</sup> ed., Longman Scientific and Technical publishers, England. P: 356 - 362.
- Ruan, J., Wu, X., Ye, Y. and Hardter, R. (1998). Effect of potassium, magnesium and sulphur applied in different forms of fertilisers on free amino acid content in leaves of tea (*Camellia sinensis* O. Kuntze). *J. Sci. Food Agric.*, 76.p: 389-396.
- Ruan, J., Wu, X., Ye, Y., and Hardter, R. (1998). Effect of potassium, magnesium and sulphur applied in different forms of fertilisers on free amino acid content in leaves of tea (*Camellia sinensis* L). *J. Sci. Food Agric.*, 76. p: 389-396.



- Sakiroglu, H., Kufrevioglu, I. O., Kocacaliskan, I., Oktay, M., & Onganer, Y. (1996). Purification and characterization of Dog-rose (*Rosa dumalis Rechst.*) polyphenol oxidase. *Journal of Agricultural and Food Chemistry*, 44, 2982–2986.
- Sanderson, G.W. (1972). The chemistry of tea and tea manufacturing. In: Runeckles, V.C. and Tso, T.C. (Ed). *Recent Advances in Phytochemistry: structural and functional aspects of Phytochemistry*. 5.p: 247- 316.
- Sanderson, G.W., (1964). The chemical composition of fresh tea flush as affected by clone and climate. *Tea Quarterly*. 35.p: 101–109.
- Sanger, F., Nicklen, S., and Coulson, A. R. (1977). DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci., USA*. 74. p: 5463-5467.
- Saravanan, M., Maria John, K. M., Raj Kumar, R., Pius, P. K. and Sasikumar, R. (2005). Genetic diversity of UPASI tea clones (*Camellia sinensis* (L.) O. Kuntze) on the basis of total catechins and their fractions. *Phytochemistry*. 66.p: 561-565.
- Scebba, F., Sebastiani, L., Vitagliano, C. (1998). Changes in activity of antioxidative enzymes in wheat (*Triticum aestivum*) seedling under cold acclimation. *Physiol. Plant*, 104(6). p: 747-752.
- Schwartz B, Olgin A, Klinman P. The role of copper in topa quinone biogenesis and catalysis, as probed by azide inhibition of a copper amine oxidase from yeast. *J. Biochem* 2001; 40: 2954–2963.
- Sealy, J. (1958). A revision of the genus *Camellia*. Royal Horticultural Society, London.
- Senthil Kumar, R. S. (2004). Studies on improving quality of south Indian black teas. Final Technical Report submitted to Tea Board, Kolkata, India. P:1-71.
- Senthil Kumar, R. S. (2004). Studies on improving quality of south Indian black teas. Final technical report submitted to Tea Board, Kolkata, India. p: 1-71.
- Senthil Kumar, R.S., Narayanan Nair Muraleedharan, N., Subramanian Murugesan, S., Kottur, G., Prem Anand, M., and Nishadh,A.(2011). Biochemical quality characteristics of CTC black teas of south India and their relation to organoleptic evaluation. *Food Chemistry*. 129.p: 117–124.



- Shahar, T., Henning, N., Gutfinger, T., Hareven, D. and Lifschitz, E. (1992). The tomato 66.3-kDa polyphenol oxidase gene: molecular identification and developmental expression. *Plant Cell*. 4. p: 135-147.
- Sharma V.S., and Venkataramani. (1974). The tea complex-taxonomy of tea clones, *Proc. Indian Acad. Sci.*, 80B, p: 178-187.
- Siddique, M., Sinha, N. K., and Cash, J. N. (1992). Characterization of polyphenol oxidase from Stanley plums. *Journal of Food Science*. 57. p: 1177–1179.
- Singh, H. P., and Ravindranath, S. D. (1994). Occurrence and Distribution of PPO activity in floral organs of some standard and local cultivars of tea. *J. Sci. Food. Agric.*, 64. p: 117-120.
- Singh, I. D. (1979). Indian tea germplasm and its contribution to the world's tea industry. *Two and a Bud* 26.p: 23-26.
- Singh, K., Kumar, S., Rani, A., Gulati, A., and Ahuja, P. (2009). Phenylalanine ammonia-lyase (PAL) and cinnamate 4-hydroxylase (C4H) and catechins (flavan-3-ols) accumulation in tea. *Funct. Integr. Genomics.*,9. p:125–34.
- Squire, G. R., Obaga, S. M., and Othieno, C. O. (1993). Altitude, temperature and shoot production of tea in Kenyan highlands. *Experimental Agriculture*. 29. p: 107-120.
- Subramanian, N., Venkatesh, P., Ganguli, S., and Sinkar, V. P. (1999). Role of polyphenol oxidase and peroxidase in the generation of black tea theaflavins. *J. Agric. Food Chem.*, 47. p : 2571-2578.
- Sud, R.G. and Badyal, J. (1989). Varietal and seasonal variations in chemical constituents of tea (*Camellia sinensis* (L.) O. Kuntze) in Himachal Pradesh. *Sri Lankan J. Tea Sci.*, 58(1): 73-78.
- Sud, R.G. and Baru, A. (2000). Seasonal variations in theaflavins, thearubigins, total colour and brightness of Kangra orthodox tea (*Camellia sinensis* (L) O Kuntze) in Himachal Pradesh. *J. Sci. Food Agric.*, 80: 1291-1299.
- Sud, R.G. and Baru, A. (2000). Seasonal variations in theaflavins, thearubigins, total colour and brightness of Kangra orthodox tea (*Camellia sinensis* (L) O Kuntze) in Himachal Pradesh. *J. Sci. Food Agric.*, 80. p: 1291-1299.



- Sud, R.G., and Badyal, J. (1989). Varietal and seasonal variations in chemical constituents of tea (*Camellia sinensis* (L.) O. Kuntze) in Himachal Pradesh. *Sri Lankan J. Tea Sci.*, 58(1). p: 73-78.
- Susana Selle's-Marchart., Juan Casado-Vela., and Roque Bru-Martínez. (2006). Isolation of a latent polyphenol oxidase from loquat fruit (*Eriobotrya japonica* Lindl.): Kinetic characterization and comparison with the active form. *Archives of Biochemistry and Biophysics*. 446. p: 175-185
- Swain, T. and Hillis, W.E. ( 1959). The phenolic constituents of *Prunus domestica*. I. The quantitative analysis of phenolic constituents. *J. Sci. Food Agric.*, 10: 63-68.
- Takeo, T. and Baker, J.E. ( 1973). Changes in multiple forms of polyphenol oxidase during maturation of tea leaves. *Phytochem.*, 12: 21-24.
- Takino, Y., Ferretti, A., Flanagan, V., Giancurto, M., Vogel, M. (1965). Structure of Theaflavin, a Polyphenol of Black Tea. *Tetrahedron Lett.* P: 4019–4025.
- Tanaka, J., Yamaguchi, N., and Nakamura, Y. (2001). Pollen parent of tea cultivar Sayamakaori with insect and cold resistance may not exist. *Breed. Res.*, 3. p: 43-48.
- Tanaka, J., Yamaguchi, N., and Nakamura, Y. (2001). Pollen parent of tea cultivar Sayamakaori with insect and cold resistance may not exist. *Breed. Res.*, 3. p: 43-48.
- Tapen, K., and Modal, T. (2008). Tea breeding. *Breeding plantation tea crops. Tropical Species.* P: 545-547.
- Thanaraj, S. N. S., and Seshadri, R. (1990). Influence of polyphenol oxidase activity and polyphenol content of tea shoot on quality of black tea. *J. Sci. Food Agric.*, 51. p: 57-69.
- Thompson J.D., Gibson T.J., Plewniak F., Jeanmougin F., Higgins D.G. (1997). The CLUSTAL\_X windows interface: Flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* 25, 4876–4882.
- Thygesen, P. W., Dry, I. B. and Robinson, S. P. (1995). Polyphenol oxidase in potato. *Plant Physiology*. 109. p: 525-531.



- Tsushida, T., and Takeo, T. (1981). The hydroxylation of p-cumarate to coffeate by the tea enzyme. *Study of Tea*.60. p: 29–33.
- Tu, Y.Y., Xu, X.Q., Xia, H.L., Watanabe, N. (2005). Optimization of theaflavin biosynthesis from tea polyphenols using an immobilized enzyme system and response surface methodology. *Biotechnol. Lett.* 27. p: 269–274.
- Ullah, M. R. (1977). Alternative test for assessment of fermentation. In: *Proc of Tocklai Conference*, 29. p: 75-77.
- Unal, M., and Sener ,A. (2006). Determination of some biochemical properties of polyphenol oxidase from Emir grape (*Vitis vinifera* L. cv. Emir). *J. Sci. Food Agric.*, 86: 2374–2379.
- Van Rossun, M. W .P. C., Alberda, M. and Van Der Plas L. H. W. (1997). Role of
- Wachira, F. N., Ng’etich, W., Omolo, J., and Mamati, G. (2002). Genotype x environment interactions for tea yields. *Euphtica*. 127. p: 289–296.
- Wachira, F. N., Powell, W., and Waugh, R. (1997). An assessment of genetic diversity among *Camellia sinensis* L. (cultivated tea) and its wild relatives based on randomly amplified polymorphic DNA and organelle-specific STS. *Heredity*. 78. p: 603-611.
- Wachira, F. N., Waugh, R., Hackett, C. A., and Powell, W. (1995). Detection of genetic diversity in tea (*Camellia sinensis*) using. RAPD markers. *Genome*. 38. p: 201-210.
- Waliszewski, K. N, Marquez, O., and Pardio, T. (2009). Quantification and characterisation of polyphenol oxidase from vanilla bean. *Food Chem.*, 117. p: 196-203.
- Wesche-Ebeling, P., Montgomery, M.W. (1990). Strawberry polyphenol oxidase: Extraction and partial characterization. *J. Food. Sci.*, 55. p: 1320-1325.
- Xu, N., and Chen, Z. (2002). Green tea, black tea and semi fermented tea. In: *Tea – Bioactivity and Therapeutic potential*. (Ed. Yang su Zhen), Taylor and Francis, London. P: 35-56.



- Yamamoto, T., Juneja, L. J., Chu, D. C., and Kim, M. (1998). Chemistry and application of green tea. New York: CRC, p: 1-158.
- Yang, C. P., Fujita, S., Ashrafuzzaman, M. D., Nakamura, N., and Hayashi, N. (2000). Purification and characterization of polyphenol oxidase from banana (*Musa sapientum* L.) pulp. *Journal of Agriculture and Food Chemistry*. 48. p: 2732–2735.
- Yang, C.S., Lee, M.J., Chen, L., Yang, G.Y. (1997). Polyphenols as inhibitors of carcinogenesis. *Environ. Health Perspect.*, 105.p: 4971–4976.
- Yang, ChangPeng, Fujita, S., Kohno, K., Kusubayashi, A., Ashrafuzzaman, M. D., and Hayashi, N. (2001). Partial purification and characterization of polyphenol oxidase from banana (*Musa sapientum* L.) peel. *Journal of Agricultural and Food Chemistry*. 49. p:1446.
- Yao, L. H., and Nursten, H. E. (1997). The chemical oxidation of individual (-) - epicatechin gallate and (-) - epigallocatechin gallate. *J. Food Fermentation and Industry.*, 23(5). p: 21-25.
- Yao, L., Caffin, N., Arcy, B.D., Jiang, Y., Shi, J., Singanusong, R., Liu, X., Datta, N., Kakuda, Y. and Xu, Y. 2005. Seasonal variations of phenolic compounds in Australia-Grown Tea (*Camellia sinensis*). *J. Agric. Food Chem.*, 53. p: 6477-6483.
- Yelena, G., Sheptovitsky, Brudvig, G.W. (1996). Isolation and characterization of spinach photosystem II membrane-associated catalase and polyphenol oxidase. *Biochem.*, 35.p: 16255-16263.
- Yi-Liang Wu, Li-Ping Pan, Si-Li Yu, and Hai-Hang Li. (2010) Cloning, microbial expression and structure–activity relationship of polyphenol oxidases from *Camellia sinensis*. *Journal of Biotechnology*, 145.p: 66–72.
- Yokozawa, T., Nakagawa, T., Shu, S. and Juneja, L.R. (2001). Protective effects of green tea polyphenols against renal disease. In: Proc. 2001 Int. Conf. on O.Cha (Tea) Culture and Science, Session III Production, Shizuoka, Japan. P: 22-25. 2001.
- Young- Goo, P., Kaundun, S. S., and Zhyvoloup, A. (2002). Use of the bulked genomic DNA-based RAPD methodology to assess the genetic diversity among abandoned Korean tea plantations. *Genetic Resources and Crop Evolution*. p: 1-0 145. p: 66–72.





Zawistowski, J., Biliaderis, C. G., and Murray, E. D. (1988). Isolation and some properties of an acidic fraction of polyphenol oxidase from Jerusalem artichoke (*Helianthus tuberosus* L.). *Journal of Food Biochemistry*. 12. p: 23–35.

Zhao, D., Liu, Z S., and Xi, B. (2001). Cloning and alignment of polyphenol oxidase cDNA of tea plant. *Journal of Tea Science*, 21(2): 94–98.

