

REFERENCES

- Ahmad, M. and Benjakul, S. (2010) Extraction and Characterization of Pepsin-Solubilized Collagen from the Skin of Unicorn Leatherjacket (*Aluterus monocerous*). *Food Chem*, 120: 817-824.
- Alberts, B., Johnson, A., Lewis, J., Raff, M., and Roberts, K. (2002) *Molecular Biology of The Cell*; Garland Science: New York, NY, USA.
- Alder-Nissen, J. (1979). Determination of the degree of hydrolysis of food protein hydrolysates by trinitrobenzene sulphonic acid. *J. Agric. Food Chem.*, 27: 1256-1262.
- AOAC, (2012) Official Methods of Analysis. Association of Official Analytical Chemist. (19th edition)
- Arend, W.P, and Dayer, J.M. (1990) Cytokines and cytokine inhibitors or antagonists in rheumatoid arthritis. *Arthritis Rheum.*, 33: 305–315.
- Arnesen, J.A., and Gildberg, A. (2007) Extraction and characterisation of gelatine from Atlantic salmon (*Salmo salar*) skin. *Bioresource Technology.*, 98(1), 53-57.
- Astusi, O., Kawahito, Y., Prudovsky, I., Tubouchi, Y., and Kimura, M. (2005) Copper chelation with tetrathiomolybdate suppresses adjuvants induced arthritis and inflammation associated cachexia in rats. *Arthr. Res. Thera.*, 7: 1174-82.
- Badii, F., and Howell, N.K. (2006). Fish gelatin: structure, gelling properties and interaction with egg albumen proteins. *Food Hydrocoll.*, 20: 630-640

References

- Bae, I., Osatomi, K., Yoshida, A., Osako K., Yamaguchi A., and Hara K., (2008) Biochemical properties of acid-soluble collagens extracted from the skins of underutilised fishes. *Food Chem.*, 108: 49–54.
- Bailey, A.J, and Light, N.D., (1989). Connective tissue in meat and meat products. London: Elsevier Applied Science.
- Bailey, A.J., Paul, R.G., and Knott, L. (1998). Mechanisms of maturation and aging of collagen. *Mech Aging and Develop* 106: 1-56.
- Balakrishnan, B., Mohanty, M., Umashankar, P.R., and Jayakrishnan, A. (2005) Evaluation of an in situ forming hydrogel wound dressing based on oxidized alginate and gelatin. *Biomat.*, 26: 6335–42.
- Balian, G., and Bowes, J.H., (1977).The structure and properties of collagen. In: The science and technology of gelatin, Ward AG, Courts A (Ed), London, Academic Press, 1–31.
- Barnett, S.E., and Irving, S.J., (1991) Studies of wound healing and the effect of dressings. In: Szycher M, editor. High performance biomaterials.Lancaster: Technonic. 583-620.
- Barrientos, S., Stojadinovic, O., Golinko, M.S., Brem, H., and Tomic-Canic, M. (2008) Growth factors and cytokines in wound healing. *Wound Repair Regen.*, 16: 585-601.
- Bazin, S., and Delaumay, A. (1976), Preparation of acid and citrate soluble collagen. In: The methodology of connective tissue research, Hall DA (Ed), Joynson–Bruvvers, Oxford, 13–18.
- Bello, A.E., Oesser S. (2006) Collagen hydrolysate for the treatment of osteoarthritis and other joint disorders: a review of the literature. *Cur.Med. Res. Opin.*, 22(11): 2221-2232.

References

- Berg, R.A., and Prockop, D.J. (1973). Thermal transition of a non-hydroxylated form of collagen: Evidence for a role for hydroxyproline in stabilizing the triple-helix of collagen. *Biochem Biophys Res Commun.*, 52: 115–120.
- Birk, D.E., and Bruckner, P. (2005). Collagen suprastructures. *Topics in Current Chemistry* 247: 185-205.
- Blagojevic, M., Jinks, C., Jeffery, A., and Jordan, K.P. (2010) Risk factors for onset of osteoarthritis of the knee in older adults: a systematic review and meta-analysis. *Osteoarthritis Cartilagen.*, 18: 24-33
- Bongartz, T., Cantaert, T., and Atkins, S.R., (2007) Citrullination in extra-articular manifestations of rheumatoid arthritis. *Rheumatology (Oxford).*, 46: 70.
- Box, G.E.P., and Wilson, K.B. (1951). On the experimental attainment of optimum conditions. *J. R. Stat. Soc. B Met.*, 13: 1-35.
- Boyce, S.T., Christianson, D.J., and Hansbrough, J.F. (1988a) Structure of a collagen-GAG dermal skin substitute optimized for cultured human epidermal keratinocytes. *Biomed Mater Res.*, 22: 939–957.
- Boyce, S.T., Glafkides, M.C., Foreman, T.J., and Hansbrough, J.F. (1988b) Reduced wound contraction after grafting of full-thickness burns with a collagen and chondroitin-6-sulfate (GAG) dermal skin substitute and coverage with biobrane. *J Burn Care Rehabil.*, 9(4): 364-70.
- Brand, D.D., Latham, K.A., and Rosloniec, E.F. (2007) Collagen-induced arthritis. *Nature Protocols* 2: 1269-1275.
- Brief, A.A., Maurer, S.G., and Di Cesare, P.E. (2001) Use of glucosamine and chondroitin sulfate in the management of osteoarthritis. *J Am Acad Orthop Surg.*, 9: 71-78.

References

- Brodsky, B., and Persikov, A.V., (2005). Molecular structure of the collagen triple helix. *Adv Protein Chem.*, 70: 301-339.
- Broughton, G., Janis, J.E., and Attinger, C.E. (2006). The basic science of wound healing (retraction of Witte M., Barbul A. In: *Surg Clin North Am* 1997; 77:509-528). *Plast Reconstr Surg* 117(7 Suppl):12S-34S.
- Bryant, S.J. and Anseth K.S. (2001) The effects of scaffold thickness on tissue engineered cartilage in photocrosslinked poly(ethylene oxide) hydrogels. *Biomater.*, 22: 619–626.
- Burghagen (1999). Collagen. In H. Belitz and W. Grosch (Eds.), *Food chem.* Berlin, Germany: Springer-Verlag Berlin Heidelberg 2:540-547.
- Cai, X., Zhou, H., Wong, Y., Xie, Y., Liu, Z., Jiang, Z., Bian, Z., Xu, H. and Liu, L. (2007). Suppression of the onset and progression of collagen-induced arthritis in rats by QFGJS, a preparation from an anti-arthritic Chinese herbal formula. *J. Ethnopharmacol.*, 110(1): 39-48.
- Caissie, R., Gingras, M., Champigny, M.F., and Berthod, F. (2006) *In vivo* enhancement of sensory perception recovery in a tissue-engineered skin enriched with laminin. *Biomater.*, 27: 2988-2993.
- Campos, A.C., Groth, A.K., and Branco, A.B., (2008). Assessment and nutritional aspects of wound healing. *Curr Opin Clin Nutr Metab Care* 11: 281-288.
- Chauhan, B. and Gupta, R. (2004). Application of statistical experimental design for optimization of alkaline protease production from *Bacillus sp.* RGR-14. *Process. Biochem.*, 39:2115-2122.
- Chen, L.Y., Remondetto, G.E., and Subirade, M. (2006) Food protein-based materials as nutraceutical delivery systems. *Tr Food Sci Technol.*, 17(5): 272-283.

References

- Chen, P., Marsilio, E., Goldstein, R.H., Yannas, and Spector, M. (2005) Formation of lung alveolar like structures in collagen-glycosaminoglycan scaffolds *in vitro*. *Tissue Eng.*, 11:1436-1448.
- Chen, X.G., Wang, Z., Liu, W.S., and Park, H.J. (2002) The effect of carboxymethylchitosan on proliferation and collagen secretion of normal and keloid skin fibroblasts. *Biomat.*, 23: 4609-4614.
- Chitra, P., Sajith lal, G.B., and Chandrakasan, G. (1998) Influence of Aloe vera on the glucoaminoglycans in the matrix of healing dermal wounds in rats. *J. Ethnopharmacol.* 59:179-186.
- Cho, S.M., Gu, Y.S., and Kim, S.B. (2005) Extraction optimization and physical properties of yellowfin tuna (*Thunnus albacares*) skin gelatine compared to mammalian gelatins. *Food Hydrocoll.*, 19: 221-229.
- Choi, Y.S., Hong, S.R., Lee, Y.M., Song, K.W., Park, M.H., and Nam, Y.S. (1999) Study on gelatin-containing artificial skin: I. Preparation and characteristics of novel gelatin-alginate sponge. *Biomat.*, 20: 409-17.
- Choi, Y.S., Lee, S.B., Hong, S.R., Lee, Y.M., Song, K.W., and Park, M.H. (2001) Studies on gelatin-based sponges. Part III: a comparative study of cross-linked gelatin/alginate, gelatin/hyaluronate and chitosan/ hyaluronate sponges and their application as a wound dressing in full-thickness skin defect of rat. *J Mater Sci Mater Med.*, 12(1):67-73.
- Chvapil, M. (1979). Industrial uses of collagen. In: Fibrous proteins: Scientific, industrial and medical aspects, Parry DAD, Creamer LK (Eds), Academic Press, London, 247– 269.
- Ciarlo, A.S., Paredi, M.E., and Alicia,N. (1997) Isolation of soluble collagen from Hake skin (*Merluccius hubbssi*). *J Aquat Food Prod Tech.*, 6 (1): 65-77.

References

- Cioca, G. (1981) Process for preparing macromolecular biologically active collagen. US Patent 4279812.
- Clark, R.A.F. (1996) Wound repair: overview and general considerations. In: Clark RAF, editor. *The molecular and cellular biology of wound repair*. New York: Plenum Press, 3–50.
- Clemente, A. (2000) Enzymatic protein hydrolysates in human nutrition. *Food Sci. Techn.* 11: 254-262.
- Conti, B., Giunchedi, P., Genta, I., and Conte, U. (2000) The preparation and in vivo evaluation of the wound-healing properties of chitosan microspheres. *Pharm Sci.*, 10: 101-104
- Cooper, G.M., (2000) *The Cell: A molecular approach*. (2nd edition).
- Courtenay, J.S., Dallman, M.J., Dayan, A.D., Martin, A., and Mosedale, B. (1980) Immunisation against heterologous type II collagen induces arthritis in mice. *Nature.*, 283:666-668.
- Dai, L., Ye, Z.Q., and Tang, M.A. (2000) Expression of transforming growth factor β 1, in rheumatoid synovia and its relationship with synovial pathological change, *Chinese J. Rheumatol.*, 4: 357-60.
- Darby, N.J. and Creighton T.E. (1993) *Protein structure* Oxford University Press.
- Davis, S.C., Badiavas, E., Rendon-Pellerano, M.I., and Pardo, R.J. (1999) Histological comparison of postoperative wound care regimens for laser resurfacing in a porcine model. *Dermatol Surg.*, 25:387-393.
- De Almeida Jackix, E., Cúneo, F., Amaya-Farfan, J., de Assunção, J.V., and Quintaes, K.D. (2010) A food supplement of hydrolyzed collagen improves compositional and biodynamic characteristics of vertebrae in ovariectomized rats. *J. Med. Food*, 13:1385-1390.

References

- De Lusto, F., Dasch, J., Keefe, L., and Ellingsworth, J. (1990) Immune responses to allogenic and xenogenic implants of collagen and collagen derivates, *Clin. Orthop. Relat. Res.* 260: 263-279.
- De Rycke, L., Peene, I., and Hoffman, I.E. (2004) Rheumatoid factor and anticitrullinated protein antibodies in rheumatoid arthritis: diagnostic value, associations with radiological progression rate, and extra-articular manifestations. *Ann Rheum Dis.*, 63:1587.
- Dressler, D.P., Barbee, W.K., and Sprenger, R. (1980) The effect of Hydron burn wound dressing on burned rat and rabbit ear wound healing. *J Trauma.*, 20(12):1024-8.
- Drury, J.L., and Mooney, D.J. (2003) Hydrogels for tissue engineering: Scaffold design variables and applications. *Biomater.*, 24: 4337-4351.
- Duan, R., Zhang, J., Du, X., and Konno, K., (2009) Properties of collagen from skin, scale and bone of carp (*Cyprinus carpio*) *Food chem.*, 112: 702-706.
- Edwards, R., and Harding, K.G. (2004) Bacteria and wound healing. *Curr Opin Infect Dis.*, 17: 91-96.
- Emna Soufi Kechaoua, C., Jean-Pascal Bergéb, Pascal Jaouena, and Raja Ben Amarc. (2013) Optimization of common cuttlefish (*Sepia officinalis*) protein hydrolysate using Pepsin by Response Surface Methodology. *J aquat food prod t.*, 24: 270-282.
- Exposito, J.Y., Le Guellec, D., Lu, Q., and Garrone, R. (1991) Short chain collagens in sponges are encoded by a family of closely related genes. *J. Biol. Chem.*, 266: 21923-21928.

References

- Fahmi, A., Morimura, S., Guo, H.C., Shigematsu, T., Kida, K., and Uemurac Y., (2004). Production of angiotensin I converting enzyme inhibitory peptides from sea bream scales. *Process Biochem.*, 39(10): 1195-1200.
- Falguni P., Shilpa V., Mann B. (2010) Production of proteinaceous antifungal substances from *Lactobacillus brevis* NCDC. *Int J Dairy Technol.* 63: 70-76.
- Feldmann, M., Brennan, F.M., Maini, R.N. (1996) Role of cytokines in rheumatoid arthritis. *Annu. Rev. Immunol.*, 14:397-440.
- Felson, D.T., Lawrence, R.C., Dieppe, P.A., Hirsch, R., Helmick, C.G., and Jordan, J.M. (2000) Osteoarthritis: new insights. Part 1: the disease and its risk factors. *Ann Intern Med.*, 133: 635-46.
- Ferreira, I., Pinho, O., Tavares, P., Pereira, A., and Roque, A.C. (2007) Development and validation of an HPLC/UV method for quantification of bioactive peptides in fermented milk. *J Liq Chromatogr Relat Technol.*, 30 (14): 2139-47.
- Foegeding, E.A., Lanier, T.C., and Hultin, H.O. (1996). Collagen. In: Food chemistry, Fennema OR (Ed), 3: 902-906, New York: Marcel Dekker, Inc.
- Foulquier, C. (2007) Peptidyl arginine deiminase type 2 (PAD-2) and PAD-4 but not PAD-1, PAD-, and PAD-6 are expressed in rheumatoid arthritis synovium in close association with tissue inflammation. *Arthritis Rheum.* 56: 3541-3553.
- Franceschi, R.T., Iyer, B.S. (1992) Relationship between collagen synthesis and expression of the osteoblast phenotype in MC3T3-E1 cells. *J Bone Miner Res.*, 7:235-246.
- Francis Suh, J.K., and Matthew, H.W.T. (2000). Application of chitosan-based polysaccharide biomaterials in cartilage tissue engineering: a review. *Biomat.*, 24: 2589-2598.

References

- Friess, W., and Lee, G. (1966) Basic thermo analytical studies of insoluble collagen matrices, *Biomat.*, 17: 2289- 2294.
- Froget, S., Barthelemey, E., Guillot, F., and Soler, C. Courdet, M.C Benbaunan, M., Dosquet, C. (2003) Wound healing mediator production by human dermal fibroblasts grown within a collagen– GAG matrix for skin repair in humans. *Eur. Cytokine Netw.*, 14: 60-64.
- Fujita, H. and M. Yoshikawa, (1999) LKPNM: A prodrug-type ACE inhibitory peptide derived from fish protein. *Immunopharmacology*, 44: 123-127.
- Funk, J.L., Oyarzo, J.N., Frye, J.B., Chen, G., Lantz, R.C., Jolad, S.D., Solyom, A.M., and Timmermann, B.N. (2006) Turmeric extracts containing curcuminoids prevent experimental rheumatoid arthritis. *J Nat Prod.*, 69(3): 351-355.
- Gbogouri, G.A., Linder, M., Fanni, J., and Parmentier, M. (2004) Influence of hydrolysis degree on the functional properties of salmon byproducts hydrolysates. *J Food Sci.*, 69(8): C615-C622.
- Gelse, K., Poschl, E., and Aigner, T. (2003). Collagens-structure, function, and biosynthesis. *Adv Drug Del Rev.*, 55: 1531-1546.
- Gilsenan, P.M., and Ross-Murphy, S.B. (2000). Rheological characterisation of gelatins from mammalian and marine sources. *Food Hydrocoll.*, 14:191-195.
- Gomathi, K.D., Gopinath, M.A., Rafiuddin, R., and Jayakumar, (2003) Quercetin incorporated collagen matrices for dermal wound healing process in rat. *Biomat.*, 24: 2767-2772.
- Gomez-Guillen, M.C., and Montero, P. (2001) Method for the production of gelatin of marine origin and product thus obtained. International Patent PCT/S01/00275.

References

- Gomez-Guillen, M.C., Turnay, J., Fernandez-Díaz, M.D., Olmo, N., Lizarbe, M.A., and Montero, P. (2002) Structural and physical properties of gelatin extracted from different marine species: A comparative study. *Food Hydrocoll.*, 16: 25-34.
- Gosain, A., and DiPietro, L.A. (2004) Aging and wound healing. *World J Surg.*, 28:321-326.
- Greenwald, R.A., (1991) Animal models for evaluation of arthritis drugs. *Experimental and Clinical Pharmacology.*, 13: 75-83.
- Grossman, S., and Bergman, M. (1992) Process for the production of gelatin from fish skins. U.S. Patent 5093474.
- Gudmundsson, M., and Hafsteinsson, H. (1997) Gelatin from cod skins as affected by chemical treatments. *J Food Sci.*, 62, 37-39.
- Guerard, F., Guimas, L., and Binet, A. (2002) Production of tuna waste hydrolysates by a commercial neutral protease preparation. *J Mol Catalysis B.*, 19–20: 489-498.
- Guillermin, F., Beaupied, H., Fabien-Soulé, V., Tome, D., Benhamou, C.L., Roux, C., and Blais, A. (2010) Hydrolyzed collagen improves bone metabolism and biomechanical parameters in ovariectomized mice: An in vitro and in vivo study. *Bone.*, 46(3): 827-834.
- Guo, S. and DiPietro, L.A. (2010) Factors Affecting Wound Healing. *J Dent Res.*, 89(3): 219-229.
- Hanai, K., Takeshita, F., and Honma, K. (2006) Atelocollagen-mediated systemic DDS for nucleic acid medicines. *Ann. N. Y. Acad. Sci.*, 1082: 9-17.
- Hansen, S.L., Voigt, D.W., Wiebelhaus, P., and Paul, C.N. (2001) Using skin replacement products to treat burns and wounds, *Adv. Skin Wound Care.*, 14: 37-44.

References

- Harirforoosh, S., and Jamali, F. (2009) Renal adverse effects of nonsteroidal anti-inflammatory drugs. *Expert Opin Drug Saf.*, 8(6): 669-681.
- Harris, E.D. (1990) Rheumatoid arthritis. Pathophysiology and implications for therapy. *N Eng J Med.*, 322:1277-1289.
- Hasegawa, D., Fujii, R., Yagishita, N., Matsumoto, N., and Aratani, S. (2010) E3 Ubiquitin Ligase Synoviolin Is Involved in Liver Fibrogenesis. *PLoS ONE* 5(10): 13590. doi:10.1371/journal.pone.0013590
- Hassan, F., and Sherief, P. M. (1994). Role and application of fish collagen. *Seafood Export Journal*, 25: 19-24.
- Hayashi, T. and Nagai, Y. (1980) The anomalous behavior of collagen peptides on sodium dodecyl sulfate-polyacrylamide gel electrophoresis is due to low content of hydrophobic aminoacid residues. *J Biochem.* 87(3): 803-808
- Henrotin, Y., Mobasher, A., and Marty, M. (2012) Is there any scientific evidence for the use of glucosamine in the management of human osteoarthritis? *Arthritis Res Ther.*, 14:201-7.
- Hinman, C. and Maibach, H. (1963) Effect of air exposure and occlusion on experimental human skin wounds. *Nature* 26(200): 377-8.
- Hinman, C.D. and Maibach, H., (2000) *Nature.*, 377-378.
- Hinrich, W.L., Lommen, E.J., Wildevuur, C.R., and Feijen, J. (1963) Fabrication and characterization of an asymmetric polyurethane membrane for use as a wound dressing. *J Appl Biomater.*, 3(4):287-303.
- Ho, H.O., Lin, C.W., and Sheu, M.T. (2001) Diffusion characteristics of collagen film. *J. Control. Release* 77: 97-105.

References

- Hoffman, S. (2001) Hydrogels for biomedical applications. *Annals of the New York Academy of Sciences.*, 944:62–73.
- Holzer, D. (1996) Gelatin production, US Patent 5484888.
- Hong, Y. Chen, W.W., Jia, Z.B., and Huang, G.R. (2013) Application of Response Surface Methodology to Optimize Hydrolysis of Collagen from Croaker Scale *Adv. J. Food Sci. Technol.*, 5: 1646-1651.
- Hopkins, S.J. (1990) Cytokines and eicosanoids in rheumatic diseases, *Ann Rheum Dis.*, 49:207-210.
- Horch, R.E. and Stark, G.B. (1998) Comparison of the effect of a collagen dressing and a polyurethane dressing on the healing of split thickness in graft (STSG) donor sites, *Scand. J. Plast. Reconstr. Surg. Hand Surg.* 32: 407-413.
- Huo, H., Li, B.F., Zhao, X., Zhuang, Y.L., Ren, G.Y., Yan, Y.P., Cai, M.Y., Zhang, X.K., and Chen, L. (2009) The effect of pacific cod (*Gadus macrocephalus*) skin gelatin polypeptides on UV radiation-induced skin photoaging in ICR mice. *Food Chem.*, 115(3): 945-950.
- Huo, J.X. and Zhao, Z. (2009) Study on enzymatic hydrolysis of *Gadus morhua* skin collagen and molecular weight distribution of hydrolysates. *Agric. Sci. China.*, 8(6): 723-729.
- Hye Kyung Kim, Myung-Gyou Kim, and Kang-Hyun Leem. (2013) Osteogenic Activity of Collagen Peptide via ERK/MAPK Pathway Mediated Boosting of Collagen Synthesis and Its Therapeutic Efficacy in Osteoporotic Bone by Back-Scattered Electron Imaging and Microarchitecture Analysis, *Molecules.*, 18: 15474-15489.
- Ignateva, N.Y., Danilov, N.A., Averkiev, S.V., Obrezkova, M.V., Lunin, V.V., and Sobol, E.N. (2007) Determination of hydroxyproline in tissues and the evaluation of the collagen content of the tissues, *J. Analytical Chem.* 62: 51–57.

References

- Ishida, Y., Fugita, T., and Asai, K. (1981) New detection and separation methods for amino acids by high performance liquid chromatography. *J Chromatogra.*, 204: 143-148.
- Iwai, K., Hasegawa, T., and Taguchi, Y. (2005) Identification of food derived collagen peptides in human blood after oral ingestion of gelatin hydrolysates. *J Agric Food Chem.*, 53:6531-6536.
- Jamilah, B., and Harvinder, K.G. (2002) Properties of gelatins from skins of fish-black tilapia (*Oreochromis mossambicus*) and red tilapia (*Oreochromis nilotica*). *Food Chem.*, 77: 81–84.
- Je, J.Y., Kim, S.Y., and Kim, S.K. (2005) Preparation and antioxidative activity of hoki frameprotein hydrolysate using ultrafiltration membranes. *Eur Food Res Technol.*, 221:157-62.
- Jeon, O., Song, S.J., Lee, K.J., Park, M.H., Lee, S.H., and Hahn, S.K. (2007) Mechanical properties and degradation behaviors of hyaluronic acid hydrogels cross-linked at various cross-linking densities. *Carbohydrate Polymers*, 70(3):251-257.
- Jerosch, J. (2011) Effects of glucosamine and chondroitin sulfate on cartilage metabolism in OA: Outlook on other nutrient partners especially omega-3 fatty acids. *Int J Rheumatol.*, 20: 969-972.
- Jhon, M.S. and Andrade, J.D. (1973) Water and hydrogels. *Journal of Biomedical Materials Research.*, 7:509-522.
- Jongjareonrak, A., Rawdkuen, S., Chaijan, M., Benjakul, S., Osako, K., and Tanaka, M. (2010) Chemical compositions and characterisation of skin gelatin from farmed giant catfish (*Pangasianodon gigas*). *Food Sci Technol.*, 43: 161-165.
- Jung, W.K., Karawita, R., Heo, S.J., Lee, B.J., Kim, S.K., and Jeon, Y.J. (2006) Recovery of a novel Ca-binding peptide from Alaska Pollack (*Theragra chalcogramma*) backbone by pepsinolytic hydrolysis. *Process Biochem.*, 41(9): 2097-2100.

References

- Karim, A.A., and Bhat, R. (2009) Fish gelatin: properties, challenges, and prospects as an alternative to mammalian gelatins. *Food Hydrocoll.* 23: 563-576.
- Katarzyna, Dybka, and Piotr walczak. (2009), Collagen Hydrolysate as a new diet supplement. *Food Chemistry and Biotechnology*, 73: 83-84.
- Keira, S.M., Ferreira, L.M., Gragnani, A., Duarte, I.S., and Barbosa, J. (2004) Experimental model for collagen estimation in cell culture. *Acta Cir Bras.*, 19:17-22.
- Keiser, H.W., Stark, G.B., Koop, J., Balcerkiewicz, A., Spilker, G., and Kreysel, H.W. (1994) Cultured autologous keratinocytes in fibrin glue suspension, exclusively and combined with STS allograft, *Burns.*, 20: 23-29.
- Kim, H.J., Choi, E.Y., Oh, J.S., Lee, H.C., Park, S.S., and Cho, C.S. (2000) Possibility of wound dressing using poly(l-leucine)/poly(ethylene glycol)/ poly(l-leucine) triblock copolymer. *Biomat.*, 21(2): 131-41.
- Kim, J.S., and Park, J.W. (2004) Characterization of acid-soluble collagen from pacific whiting surimi processing byproducts. *J. Food Sci* 69(8): 637-642.
- Kim, S.K., and Mendis, E. (2006) Bioactive compounds from marine processing byproducts –a review. *Food Res Int.*, 39: 383–93.
- Kim, S.K., and Wijesekara, I. (2010) Development and biological activities of -derived bioactive peptides: A review. *J Funct Foods.*, 2:1–9
- Kim, S.K., Kim, Y.T., Byun, H.G., Nam, K.S., Joo, D.S., Shahidi, F. (2001) Isolation and characterization of antioxidative peptides from gelatin hydrolysate of *Alaska pollack* skin. *J. Agric. Food Chem.* 49(4):1984-1989

References

- Kim, U.J., Park, J.C., Li, J., Jin, H. J., Valluzzi, R., and Kaplan, D. L. (2004) Structure and properties of silk hydrogels. *Biomacromolecules.*, 5:786-792.
- Kimura, S., Miyauchi, Y., and Uchida, (1991) Scale and bone type I collagens of carp (*Cyprinus carpio*) *Physiol.*, 99B (2): 473-476.
- King, S.R., Hickerson, W.L., and Proctor, K.G. (1991) Beneficial actions of exogenous hyaluronic acid on wound healing. *Surgery.*, 109: 76-84.
- Kinloch, A., Lundberg, K., and Wait, R. (2008) Synovial fluid is a site of citrullination of autoantigens in inflammatory arthritis. *Arthritis Rheum.*, 58: 2287.
- Kinne, R.W., Brauer, R., and Stuhlmuller, B. (2000) Macrophages in rheumatoid arthritis, *Arthritis Res.*, 2: 189-202.
- Kisiday, J., Jin, M., Kurz, B., Hung, H., Semino, C., and Zhang, S. (2002) Self-assembling peptide hydrogel fosters chondrocyte extracellular matrix production and cell division: implications for cartilage tissue repair. *Proc Natl Acad Sci U S A.*, 99(15): 9996-10001.
- Kittiphattanabawon, P., Benjakul, S., Visessanguan, W., Kishimura, H., and Shahidi, F. (2010). Isolation and characterisation of collagen from the skin of brown banded bamboo shark (*Chiloscyllium punctatum*). *Food Chem.*, 119: 1519-1526.
- Kittiphattanabawon, P., Benjakul, S., Visessanguan, W., Nagai, T., and Tanaka, M. (2005) Characterisation of acid-soluble collagen from skin and bone of bigeye snapper (*Priacanthus tayenus*). *Food Chem.*, 89(3): 363-372.
- Knapp, T.R., Luck, E., and Daniels, J.R. (1977) Behaviour of solubilised collagen as a bioimplant. *J Surg Res* 23: 96-105.

References

- Korhonen, H., Pihlanto-Leppala, A., and Tupasela, T. (1998). Impact of processing on bioactive proteins and peptides. *Trends in Food Science and Technology*, 9: 307-319.
- Kristinsson, H.G., and Rasco, B.A. (2000) Fish protein hydrolysates: production, biochemical, and functional properties. *Crit Rev Food Sci Nutr*, 40(1):43-81.
- Kuberka, M., von Heimburg, D., Schoof, H., Heschel, I., and Rau, G. (2002) Magnification of pore sizes in biodegradable matrices, *Int. J. Artif. Organs.*, 25: 67-73.
- Kubo, K. and Takagi, T. (1984) The alpha 1(I) and alpha 2(I) chains of collagen separate in sodium dodecyl sulphate-polyacrylamide gel electrophoresis due to differences in sodium dodecyl sulphate binding capacities. *Coll. Relat. Res.* 4:201-208.
- Kuhn, K. (1987) Structure and Function of Collagen Types (Mayne, R., and Burgeson, R. E., eds) pp. 1-42, Academic Press, Orlando, FL
- Kuptniratsaikul, V., Tosayanonda, O., Nilganuwong, S., and Thamalikitkul, V. (2002) The Efficacy of a Muscle Exercise Program to Improve Functional Performance of the Knee in Patients with Osteoarthritis. *J Med Assoc Thailand.*, 85(1):33-39.
- Kuriyama, N., Shoji, N., Taniguchi, K., and Omote, M. (2005) Efficient peptide purification by HPLC- effect of pore size, particle size and chemistry. Kyoto, Japan: YMC Co., Ltd.
- Laemmli U. K. (1970) Cleavage of Structural Proteins during the Assembly of the Head of Bacteriophage T4. *Nature.* 227: 680-685.
- Lee, C.H., Singla, A., and Lee, Y. (2001) Biomedical applications of collagen. *Int. J. Pharm.*, 221: 1-22.

References

- Lee, S.L. (1983) Optimal conditions for long-term storage of native collagens. *Coll Relat Res* 3: 305-315.
- Levy, A.S.A., Simon, O., Shelly, J., and Gardener, M. (2006) 6-Shagaol Reduced Chronic Inflammatory Response in the Knees of Rats Treated with Complete Freund's Adjuvant, *BMC Pharmacology.*, 6(12): 1-8.
- Leyland, K.M., Hart, D.J., and Javaid, M.K. (2012) The natural history of radiographic knee osteoarthritis: A fourteen-year population-based cohort study. *Arthritis Rheum.* , 64: 2243-2251.
- Li, C.M., Zhong, Z.H., Wan, Q.H., Zhao, H., Gu, H.F., and Xiong, S.B. (2008) Preparation and thermal stability of collagen from scales of grass carp (*Ctenopharyngodon idellus*). *Euro Food Res Technol.*, 227: 1467-1473.
- Li, S.T. (1995) Tissue-derived biomaterials (collagen). In: The biomedical engineering handbook, Bronzino JD (Ed), CRC Press, Boca Raton, FL, pp 627–647.
- Lillie, L.E., Temple, N.J., Florence, L.Z. (1996) Reference values for young normal Sprague-Dawley rats: weight gain, hematology and clinical chemistry. *Hum Exp Toxicol.* 15:612–616.
- Lin, Y.K., and Liu, D.C. (2006) Comparison of physical-chemical properties of type I collagen from different species. *Food Chem.*, 99: 244-251.
- Liu, H. Y., Li, D., & Guo, S. D. (2007). Studies on collagen from the skin of channel catfish (*Ictalurus punctatus*). *Food Chem.*, 101: 621-625.
- Liu, W.T., Li, G.Y., Miao, Y.Q., and Wu, X.H. (2009) Preparation and characterization of pepsin-solubilized type I collagen from the scales of snakehead (*Ophiocephalus argus*). *J Food Biochem.*, 33: 20–37.

References

- Loeser, R.F., Goldring, S.R., and Scanzello, C.R. (2012) Osteoarthritis: A disease of the joint as an organ. *Arthritis Rheum.* , 64:1697-1707.
- Mathieu, D., Linke, J.C., and Wattel, F. (2006) Non-healing wounds. In: Handbook on hyperbaric medicine, Mathieu DE, editor. Netherlands: Springer, pp. 401-427.
- Matsui, R., Ishida, M., and Kimura, S. (1991) Characterization of an alpha 3 chain from the skin type I collagen of chum salmon (*Oncorhynchus keta*), *Comp.Biochem.Physiol.B.*, 99:171-174.
- McInnes, I.B., and O Dell J.R. (2010) State-of-the-art: rheumatoid arthritis. *Ann Rheum Dis.*, 69:1898-1906.
- McInnes, I.B., and Schett, G. (2011) The pathogenesis of rheumatoid arthritis. *N Engl J Med*, 365:2205-2219.
- Mendis, E., Rajapakse, N., and Kim, S.K. (2005) Antioxidant properties of a radical- scavenging peptide purified from enzymatically prepared fish skin gelatin hydrolysate. *J. Agric. Food Chem.*, 53(3): 581-587.
- Menke, N.B., Ward, K.R., Witten, T.M., Bonchev, D.G., and Diegelmann, R.F. (2007) Impaired wound healing. *Clin Dermatol.*, 25:19-25.
- Meszaros, A.J., Reichner, J.S., and Albina, J.E. (2000) Macrophage-induced neutrophil apoptosis. *J Immunol.*, 165:435-441.
- Miller, E.J., Piez, K.A, Reddi, A.H. (1984) Chemistry of collagens and their distribution. In: Extracellular matrix biochemistry, (Ed), Elsevier, New York, 41-82.
- Miyata, T., Taira, T., and Noishiki, Y. (1992) Collagen engineering for biomaterial use. *Clinical Materials.*, 9: 139–148.

References

- Mizuno, M. and Kuboki, Y. (2001) Osteoblast-related gene expression of bone marrow cells during the osteoblastic differentiation induced by type I collagen. *J Biochem.*, 129(1):133-8.
- Montero, P. and Borderias, J. (1989) Changes in hake muscle collagen during frozen storage due to seasonal effects. *International Journal of Refrigeration.*, 12: 220-223.
- Montero, P., Jimenez-Colmenero, F., and Borderias, J. (1991). Effect of pH and the presence of NaCl on some hydration properties of collagenous material from trout muscle and skin. *J. Sci. Food Agr.*, 54:137-146.
- Moore, R.A., Derry, S., and McQuay, H.J. (2007) Cyclo-oxygenase-2 selective inhibitors and nonsteroidal anti-inflammatory drugs: balancing gastrointestinal and cardiovascular risk. *BMC Musculoskelet Disord.*, 8:73.
- Morrissey M.T. and Park J.W. (2000) Manufacturing of surimi from light muscle fish. In: Park JW, editor. *Surimi and surimi seafood*. New York: Marcel Dekker Inc. 23- 58.
- Morrissey, M.T., Lin, J., and Ismond, A. (2005) Waste management and by-product utilization. In: *Surimi and surimi seafood*, (2nd edn), Florida, CRC Press, Taylor and Francis Group, 279-323.
- Moskowitz, R.W. (2000) Role of collagen hydrolysate in bone and joint disease. *Semin. Arthritis Rheum.*, 30: 87-99.
- Mosser, D.M., and Edwards, J.P. (2008) Exploring the full spectrum of macrophage activation. *Nat Rev Immunol.*, 8:958-969.
- Mullally, M.M., O'Callaghan, D.M., FitzGerald, R.J., Donnelly, W.J., and Dalton, J.P. (1995) Zymogen activation in pancreatic endoproteolytic preparations and influence on some whey protein characteristics. *J. Food Sci.*, 60: 227-233.

References

- Murashita, Y.Y., Nakayama., T. Hirano, S., and Ohashi. (1996) Acceleration of granulation tissue growth by hyaluronic acid in artificial skin. *Br. J. Plast. Surg.*, 49: 58- 63.
- Muyonga, J H., Colec, C. G. B. and Duodu, K. G. (2004a) Extraction and physicochemical characterisation of Nile perch (*Lates niloticus*) skin and bone gelatine. *Food Hydrocoll.*, 8: 581-592.
- Muyonga, J.H., Cole, C.G.B., and Duodu, K.G. (2004b) Characterization of acid soluble collagen from skins of young and adult Nile perch (*Lates niloticus*). *Food Chem.*, 85: 81-89.
- Mewar, D.and Wilson, A.G. (2006) Autoantibodies in rheumatoid arthritis: a review. *Biomed.Pharmacother.*, 60: 648-655.
- Myers, R.H., Montgomery D.C., and Anderson-Cook, C.M. (2008) Response Surface Methodology: Process and Product Optimization Using Designed Experiments. 3rd Edn., Wiley, New York
- Nagai N., Nakayama, Y., Zhou, Y.M., Takamizawa, K., Mori, K., and Munekata, M. (2008) Development of salmon collagen vascular graft: mechanical and biological properties and preliminary implantation study. *J. Biomed. Mater. Res. Part B.*, 87:432-439.
- Nagai, T. and Suzuki, N. (2000a) Isolation of collagen from fish waste material skin, bones and fins. *Food Chem.*, 68(3):277-81.
- Nagai, T. and Suzuki, N. (2002) preparation & partial characterization of collagen from paper nautilus (*Argonauta argo*) outer skin. *Food chem.* 76:149-153.
- Nagai, T., and Suzuki, N. (2000b) Isolation of collagen from fish waste material - skin, bone and fins. *Food Chem.*, 60: 277-281.
- Nagai, T., Araki Y., and Suzuki, N. (2002) Collagen of the skin of ocellate puffer fish (*Takifugu rubripes*). *Food Chem.*, 78: 173-177.

References

- Nagai, T., Yamashita, E., Taniguchi, K., Kanamori, N., and Suzuki N. (2001) Isolation and characterisation of collagen from the outer skin waste material of cuttlefish (*Sepia lycidas*). *Food Chem.* 72: 425-429.
- Najafian, L. and Babji, A.S. (2012) A review of fish-derived antioxidant and antimicrobial peptides: Their production, assessment, and applications. *Peptides*, 33(1): 178-185.
- Nakatani, S., Mano, H., Sampei, C., Shimizu, J., and Wada, M. (2009) Chondroprotective effect of the bioactive peptide prolyl-hydroxyproline in mouse articular cartilage *in vitro* and *in vivo*. *Osteoarthritis and Cartilage.*, 17(12):1620-1627.
- Nam, K.A., You, S.G., and Kim, S.M. (2008) Molecular and Physical Characteristics of Squid (*Todarodes pacificus*) Skin Collagens and Biological Properties of Their Enzymatic Hydrolysates. *J. Food Sci.*, 73:249-251.
- Ngo, D.H., Qian, Z.J., Ryu, B.M., Park, J.W., and Kim, S.K. (2010) *In vitro* antioxidant activity of a peptide isolated from Nile tilapia (*Oreochromis niloticus*) scale gelatin in free radical-mediated oxidative systems. *J. Funct. Foods.*, 2(2): 107-117.
- Nissler, K., Pohlers, D., and Huckel, M. (2004) Anti- CD4 monoclonal antibody treatment in acute and early chronic antigen induced arthritis: influence on macrophage activation. *Ann. Rheum.Dis.*, 63: 1470-7.
- Nomura, Y., Oohashi, K., Watanabe, M., and Kasugai, S. (2005) Increase in bone mineral density through oral administration of shark gelatine to ovariectomized rats. *Nutrition.*, 21(11-12): 1120-1126.
- Nomura, Y., Sakai, H., Ishii, Y., and Shirai, K. (1996) Preparation and some properties of type I collagen from fish scales. *Biosci Biotechnol Biochem.*, 60: 2092-2094.

References

- Noormohamed, S.E. and Ray, T. (1998) Effect of 'Compound R' on thermal burn and full-depth wound contracture in fuzzy rats. *J Burn Care Rehabil.*, 19:213-5.
- Oesser, S. and Seifert, J. (2003) Stimulation of type II collagen biosynthesis and secretion in bovine chondrocytes cultured with degraded collagen. *Cell Tissue Research.* , 311(3): 393-399.
- Oesser, S., Adam, M., Babel, W., and Seifert, J. (1999) Oral administration of ¹⁴C labelled gelatine hydrolysate leads to an accumulation of radioactivity in cartilage of mice. *J Nutr.*, 129(10):1891-1895.
- Ogawa, M., Moody, M.W., Portier, R.J., Bell, J., Schexnayder, M.A., and Losso, J.N. (2003). Biochemical properties of black drum and sheepshead seabream skin collagen. *J. Agric. Food Chem.*, 51: 8088-8092.
- Ogawa, M., Portier, R.J., Moody, M.W., Bell, J., Schexnayder, M.A., and Losso, J.N. (2004). Biochemical properties of bone and scale collagens isolated from the subtropical fish black drum (*Pogonia cromis*) and sheepshead seabream (*Archosargus probatocephalus*). *Food Chem.*, 88 (4): 495-501.
- Ohara, H., Matsumoto, H., Ito, K., Iwai, K., and Sato, K. (2007) Comparison of quantity and structures of hydroxyproline-containing peptides in human blood after oral ingestion of gelatin hydrolysates from different sources. *J Agric Food Chem.*, 55(4):1532-5.
- Paddle-Ledinek, J.E., Nasa, Z., and Cleland, H.J. (2006) Effect of different wound dressings on cell viability and proliferation. *Plast Reconstr Surg.*, 117 (Suppl):110S-8S.
- Palosuo, T, Lukka, M, and Alenius, H. (1998) Purification of filaggrin from human epidermis and measurement of antifilaggrin autoantibodies in sera from patients with

References

- rheumatoid arthritis by an enzyme-linked immunosorbent assay. *Int Arch Allergy Immunol.*, 115:294.
- Panduranga and Rao, K. (1995). Recent developments of collagen-based materials for medical applications and drug delivery. *J Biomater Sci Polym.*, 7(7):623- 645.
- Pati, F., Datta, P., Adhikari, B., Dhara, S., Ghosh, K., and Mohapatra, P.K.D. (2012) Collagen scaffolds derived from fresh water fish origin and their biocompatibility. *J Biomed Mater Res A.*, 1001068-1079.
- Pearson, C.M. and Wood, F.D. (1959) Studies of polyarthritis and other lesions induced in rats by injection of mycobacterial adjuvant. I. General clinical and pathologic characteristics and some modifying factors. *Arthritis Rheumatoid.*, 2:440-459.
- Peppas, N.A. and Khare A.R. (1993) Preparation, structure and diffusional behavior of hydrogels in controlled release. *Advanced Drug Delivery Reviews.*, 11:1–35.
- Perkins, D.N., Pappin, D.J., Creasy, D.M., and Cottrell, J.S. (1999) Probability-based protein identification by searching sequence databases using mass spectrometry data. *Electrophoresis.*, 20:3551-67.
- Persico, F.J., Pritchard, J.F., Fischer, M.C., Yorgey, K., Wong, S., and Carson, J. (1988) Effect of Tolmetin Glycine Amide, a Prodrug of Tolmetin Sodium, on Adjuvant Arthritis in the rat, *The Journal of Pharmacology and Experimental Therapeutics.*, 247(3): 889-896.
- Petibois, C., Gionnet, K., Goncalves, M., Perromat, A., Moenner, M. and Deleris, G. (2006). Analytical performances of FT-IR spectrometry and imaging for concentration measurement within biological fluids, cells, and tissues: *Analysis.* 131: 640-647.

References

- Piez, K.A. (1984) Molecular and aggregate structures of the collagens. In: Extracellular matrix biochemistry, Piez KA, Reddi AH (Ed), Elsevier, New York, pp 1–40.
- Piez, K.A. (1985) Collagen. In: Encyclopedia of polymer science and engineering, Kroschwitz JI (Ed), Wiley, New York, pp 699–727.
- Pihlanto, A. and Korhonen H. (2003) Bioactive peptides and proteins. *Adv Food Nutr Res.*, 47:175-276.
- Pihlanto-Leppala, A. (2000) Bioactive peptides derived from bovine wheyproteins: opioid and ace-inhibitory peptides. *Trends Food Sci Tech*, 11:347-356.
- Poppe, J. (1997) Gelatin. In: Thickening and gelling agents for food, Imeson A (Ed), Blackie Academic and Professional, London, pp 144-168.
- Prabjeet Singh, Benjakul S, Maqsood S, Kishimura H. (2011). Isolation and characterisation of collagen extracted from the skin of striped catfish (*Pangasianodon hypophthalmus*). *Food Chem.* 124(1):97-105.
- Prockop, D.J. and Kivirikko, K.I. (1995) Collagens: Molecular biology, diseases, and potentials for therapy. *Annu Rev Biochem.*, 64: 403-434.
- Purna, S.K. and Babu, M. (2000) Collagen based dressings—a review. *Burns.*, 26:54-62.
- Quinn, K.J., Courtney, J.M., Evans, J.H., Gaylor, J.D.S., and Reid, W.H. (1985) Principles of burn dressings. *Biomat.*, 6:369-77.
- Raabe, O., Reich, C., Wenisch, S., Hild, A., Burg-Roderfeld, M., Siebert, H.C., and Arnhold, S. (2010) Hydrolyzed fish collagen induced chondrogenic differentiation of equine adipose tissue-derived stromalcells. *Histochemistry and Cell Biology.*, 134(6): 545-554.

References

- Rajapakse, N., Mendis, E., Jung, W.K., Je J.Y. and Kim, S.K. (2005) Purification of a radical scavenging peptide from fermented mussel sauce and its antioxidant properties. *Food Res. Int.*, 38: 175-182.
- Ramshaw, J.A.M., Werkmeister, J.A., and Glattauer, V. (1995) Collagen based biomaterials. *Biotechnol Rev.*, 13: 336-382.
- Rebeca, B.D., Pena-Vera, M.T., and Diaz-Castaneda, M. (1991) Production of fish protein hydrolysates with bacterial proteases; Yield and nutritional value. *J. Food Sci.* 56: 309-314.
- Reddy, G.K., and Enwemeka, C.S. (1996) A simplified method for the analysis of hydroxyproline in biological tissues. *Clin. Biochem.* 29: 225-229.
- Regenstein, J.M. and Zhou, P. (2007) Collagen and gelatin from marine by-product. In: Maximising the value of marine by-products, Shahidi F (Ed), CRC Press, Boca Raton, Florida, pp 279-303.
- Rehn, M., Veikkola, T., Kukk-Valdre, E. Interaction of endostatin with integrins implicated in angiogenesis. (2001) *Proc. Natl. Acad. Sci.* 98:1024-1029.
- Rennekampff, H.O., Kiessig, V., Griffey, S., Greenleaf, G., and Hansbrough, J.F. (1997) Acellular human dermis promotes cultured keratinocyte engraftment. *J Burn Care Rehabil.*, 18:535-544.
- Rigby, B.J. (1968) Amino-acid composition and thermal stability of the skin collagen of the Antarctic ice-fish. *Nature.*, 219:166-167
- Roberts, P.R., Burney, J.D., Black, K.W., and Zaloga, G.P. (1999) Effect of chain length on absorption of biologically active peptides from the gastro intestinal tract. *Digestion.*, 60(4):332-337.

References

- Roreger, M. (1995) Collagen preparation for the controlled release of active substances. PCT WO 95/28964
- Roth, S.H. (2005) Nonsteroidal antiinflammatory drug gastropathy: we started it, why don't we stop it? *J Rheum.*, 32(7):1189-1191.
- Ruszczak, Z.B. (2000) Modern aspects of wound healing: an update, *Dermatol. Surg.* 26: 219-229.
- Ruszczak, Z.B. and Schwartz, R.A. (1999) Collagen uses in dermatology: an update. *Dermatology (Basel)* 199: 285- 289.
- Sachs, D.H. (1998) The immunology of xenotransplantation, in: J. Fishman, D. Sachs, R. Shaikh (Eds.), *Xenotransplantation: Scientific Frontiers and Public Policy*, *Ann. N.Y.Acad.Sci.*, 862:1- 4.
- Sadowska, M., Kolodziejska, I., and Niecikowska, C. (2003) Isolation of Collagen from the Skins of Baltic cod (*Gadus morhua*) *Food Chem.* 81: 257-262.
- Sano, A., Maeda, M., and Nagahara S. (2003) Atelocollagen for protein and gene delivery. *Advanced Drug Delivery Reviews.*, 55:1651-1677.
- Saskia Schadow, Hans-Christian, Siebert, Gunter Lochnit, Jens Kordelle, Markus Rickert, Sawitzke, A.D., Shi, H., *et.al.*, (2010) Clinical efficacy and safety of glucosamine, chondroitin sulphate, their combination, celecoxib or placebo taken to treat osteoarthritis of the knee: 2-year results from GAIT. *Ann Rheum Dis*, 69:1459-1464
- Sawitzke, A.D., She, H., Finco, M.F., Dunlop, D.D., Harris, C.L., Singer, N.G., Bradley, J.D., Silver, D., Jackson, C.G., Lane, N.E., Oddis, C.V., Wolfe, F., Lisse, J., Furst, D.E., Bingham, C.O., Reda, D.J., Moskowitz, R.W., Willians, H.J. and Clegg, D.O. (2010) Clinical efficacy and safety of glucosamine, chondroitin sulfate, their combination, celecoxib or placebo

References

- taken to treat osteoarthritis of the knee: 2-year results from GAIT. *Ann. Rheum. Dis.* 69:1459-1464.
- Schagger, H., and Von Jagow, G. (1987) Tricine-sodium dodecyl sulfate-polyacrylamidegel electrophoresis for the separation of proteins in the range from 1 to 100 kDa *Analytical Biochem.*, 166(2):368-379.
- Schellekens, G.A., de Jong, B.A., and Van den Hoogen, F.H. (1998) Citrulline is an essential constituent of antigenic determinants recognized by rheumatoid arthritis-specific autoantibodies. *J Clin Invest.*, 101:273.
- Schellekens, G.A., Visser, H., de Jong, B.A., van den Hoogen, F.H., Hazes, J.M., and Breedveld, F.C., (2000) The diagnostic properties of rheumatoid arthritis antibodies recognizing a cyclic citrullinated peptide. *Arthritis Rheum.*, 43:155-163.
- Schoof, H.J., Apel, I., Heschel, G., and Rau, (2001) Control of pore structure and sizes in freeze-dried collagen sponges, *J. Biomed. Mater. Res.* 58: 352-357.
- Scott, D.L., Shipley, M., and Dawson, A. (1998) The clinical management of rheumatoid arthritis and osteoarthritis: Strategies for improving clinical effectiveness. *Br. J. Rheumatol.*, 37:546-54.
- Senaratne, L.S., Park, P., Kim, S. (2006) Isolation and characterization of collagen from brown backed toadfish (*Lagocephalus gloveri*) skin. *Bioresour Technol.* 97:191-197.
- Shahidi, F. and Zhong, Y. (2008) Bioactive peptides. *J AOAC Int*; 9(4):914-31.
- Shanmugasundaram, N., Ravichandran, P., Reddy, P.N., Ramamurty, N., Pal, S., and Rao, K.P. (2001) Collagen-chitosan polymeric scaffolds for the in vitro culture of human epidermoid carcinoma cells. *Biomater.*, 22(14):1943-1951.

References

- Shen X.R., Kurihara H., and Takahashi K. (2007) Characterization of molecular species of collagen in scallop mantle. *Food Chem.* 102(4): 1187-1191.
- Siedler, S. and Schuller-Petrovic, S. (2000) Allogenic keratinocytes suspended in human fibrin glue used for wound healing support in chronic leg ulcers. *Arch. Dermatol.*, 136: 676-678.
- Simoës, S.I., Delgado, T.C., Lopes, R.M., Jesus, S., Ferreira, A.A., Morais, J.A., Cruz, M.E.M., Corvo, M.L., and Martins, M.B.F. (2005) Developments in the rat adjuvant arthritis model and its use in therapeutic evaluation of novel non-invasive treatment by SOD in transfersomes. *Journal of Controlled Release.*, 103: 419-434.
- Simon, G., (1965) Alterations in joint space and associated home change, in: Principles of bone X-ray diagnosis, Butlerworth and Co, Great Britain., pp. 157–163.
- Singer, A.J., Berrutti, L., Henry, C., Thode, J., and McClain, S.A., (1999) Octylcyanoacrylate for the treatment of partial-thickness burns in swine: a randomized, controlled experiment. *Acad Emer Med.*, 6:688-92.
- Singh, M.P., Stefko, J., Lumpkin, J.A., and Rosenblatt, J. (1995) The effect of electrostatic charge interactions on release rates of gentamicin from collagen matrices. *Pharm Res.*, 12: 1205-1210.
- Sionkowska, A., Skopinska-Wisniewska, J., and Wisniewski, M. (2009) Collagen synthetic polymer interactions in solution and in thin films. *J Molecular Liquids.*, 145: 135-138.
- Slade, L., and Levine, H. (1987) Polymer-chemical properties of gelatin in foods. In: Advances in meat research, collagen as a food, Pearson AM Dutson TR, Bailey AJ (Ed), 4AVI Publishers, New York, pp 251-266.

References

- Smith, W.L., De Witt, D.L. and Garavito, R.M. (2000) Cyclooxygenases: structural, cellular, and molecular biology. *Annu Rev Biochem.*, 69: 145-82.
- Song, E., Yeon Kim, S., Chun, T., Byun, H.J., and Lee, Y.M. (2006) Collagen scaffolds derived from marine source and their biocompatibility. *Biomater.*, 27:2951-2961.
- Soo C.,G. Rahbar, R.L., and Moy. (1993) The immunology of bovine collagen implants, *J. Dermatol. Surg. Oncol.* 19: 431-434.
- Spreafico, A. (2006) Osteogenic growth peptide effects on primary human osteoblast cultures: potential relevance for the treatment of glucocorticoid induced osteoporosis. *J Cell Biochem.*, 98(4): 1007-20.
- Srivastava, S., Gorham, S.D., French, D.A., Shivas, A.A., Courtney, J.M. (1990) *In vivo* evaluation and comparison of collagen, acetylated collagen and collagen/glycosaminoglycan composite films and sponges as candidate biomaterials. *Biomater.*, 11: 155-161.
- Stainsby, G. (1987) Gelatin gels. In: *Advances in meat research: Collagen on a food*. Pearson AM, Dutson TR, Bailey AJ (Ed), New York, Van Nostrand Reinhold Company Inc. 4, 209-222.
- Steffen, C., Timpl, R., and Wolff, I. (1968) Immunogenicity and specificity of collagen. V. Demonstration of three different antigenic determinants on calf collagen. *Immunology.*, 15: 135-144.
- Stemberger, A., Grimm, H., Bader, F., Rahn, H.D., and Ascherl, R. (1997) Local treatment of bone and soft tissue infections with the collagen-gentamicin sponge. *Eur J Surg.*, 163: 17-26.
- Su CH, Sun C-S, Juan S-W, Hu C-H, Ke W-T, and Sheu M.T. (1997) Fungal mycelia as the source of chitin and polysaccharides and their applications as skin substitutes. *Biomater.*,18(17):1169-74.

References

- Sudaroli, M., and Chatterjee, T.K. (2007) Evaluation of red and white seed extracts of *Abrus precatorius* Linn. Against Freund's complete adjuvant induced arthritis in rats. *J. Med. Plants. Res.*, 1: 86-94.
- Suetsuna, K., Ukeda, H., and Ochi, H. (2000) Isolation and characterization of free radical scavenging activities peptides derived from casein. *J. Nutr. Biochem.*, 11(3): 128-131.
- Sugiura H., S. Yunoki, E. Kondo, T. Ikoma, J. Tanaka, and Yasuda, K. (2009) In vivo biological responses and bioresorption of tilapia scale collagen as a potential biomaterial. *J Biomater Sci Polym.*, 20: 1353-1368.
- Suzuki, Y., Tanihara, M., Nishimura, Y., Suzuki, K., Kakimaru, Y., and Shimizu, Y. (1997) A novel wound dressing with an antibiotic delivery system stimulated by microbial infection. *ASAIO J.*, 43(5): M854.
- Swatschek D, Shatton W, Kellermann J, Muller, Keuter J. (2002) Marine sponge collagen: isolation, characterization and effects on the skin parameters surface-pH, moisture and sebum. *Eur J Pharm.*, 53(1): 107-113
- Takeshi Nagai and Nobutaka Suzuki. (2000) Isolation of collagen from fish waste material - skin, bone and fins, *Food Chem.*, 60: 277-281.
- Takeshi, K., Shoichi, N., Yasunori, K., Yasuhiko, I. (2002) Report of the committee on the classification and diagnostic criteria of diabetes mellitus (The Committee of the Japan Diabetes Society on the diagnostic criteria of diabetes mellitus) *Diabetes Res Clinical Pract.*, 55:65-85.
- Timpl, R. (1984) Immunology of the collagen, in: K.A. Piez, A.H. Reddi (Eds.), *Extracellular Matrix Biochemistry*, Elsevier, Amsterdam, 159-188.

References

- Trelstad, R.L. and Birk, D.E (1984) Collagen fibril assembly at the surface of polarized cells, in: *Extracellular matrix in Development* (R.L.Trelstad, ed) pp.513-543, Liss, New York
- Trevitt, C.R. and Singh, P.N. (2003) Variant Creutzfeldt-Jakob disease: pathology, epidemiology, and public health implications. *Am J Clin Nutr.*, 78: 651S-6S.
- Vacanti, J.P. and Langer, R.S. (1998) Preparation of three-dimensional fibrous scaffold for attaching cells to produce vascularised tissue in vivo. US Patent 5,770,193 (MIT).
- Van der Rest, M. and Garrone, R. (1991) Collagen family of proteins. *FASEB J.*, 5: 2814-2823.
- Vijayalakshmi, M.A., Lemieux, L., and Amiot, J. (1986) High performance size exclusion liquid chromatography of small molecular weight peptides from protein hydrolysates using methanol as a mobile phase additive. *J Liq Chromatogr.*, 9:3559-3576.
- Vivian Zague, Vanessa de Freitas, Marina da Costa Rosa, Geórgia, A., Ivaes de Castro, Ruy, G., Jaeger, and Gláucia, M. Machado-Santelli, (2011) Collagen Hydrolysate Intake Increases Skin Collagen Expression and Suppresses Matrix Metalloproteinase II Activity. *J Med Food.*, 14 (6): 618-624.
- Vojdani, F. (1996) Solubility: In *Methods of Testing Protein Functionality*. London: Blackie Academic and Professional, 1: 11-60.
- Wagner, W.D. (1979) A more sensitive assay discriminating galactosamine and glucosamine in mixtures. *Anal. Biochem.*, 94: 394-398.
- Wallace, D.G. and Rosenblatt, J. (2003) Collagen gel systems for sustained delivery and tissue engineering. *Advanced Drug Delivery Reviews.*, 55: 1631–1649,

References

- Walrand, S., Chiotelli, E., Noirt, F., Mwewa, S., and Lassel, T. (2008) Consumption of a functional fermented milk containing collagen hydrolysate improves the concentration of collagen-specific amino acids in plasma. *J. Agric. Food Chem.*, 56(17): 7790-7795.
- Wang, C.L., Miyata, T., Weksler, B., Rubin, A.L., and Stenzel, K.H. (1978) Collagen-induced platelet aggregation and release. I. Effects of side-chain modifications and role of arginyl residues. *Biochem Biophys Acta.*, 544: 555-567.
- Wang, L., An, X., Yang, F., Xin, Z., Zhao, L., and Hu, Q. (2008). Isolation and characterization of collagens from the skin, scale and bone of deep-sea red fish (*Sebastes mentella*). *Food Chem.*, 108(2): 616-623.
- Weadock, K.S., Miller, E.J., Keuffel, E.L., and Dunn, M.G. (1996) Effect of physical crosslinking method on collagen-fiber durability in proteolytic solutions. *J Biomed Mater Res.*, 32: 221-226.
- Whitford, D. (2005) *Proteins: Structure and Function*. John Wiley and Sons Ltd., Chichester England, pp 528.
- Williams, R.O. (1998) Rodent models of arthritis: Relevance for human disease. *Clin Exp Immunol.*, 114:330-2.
- Winter, G.D. (1962) Formation of the scab and the rate of epithelialization of superficial wounds in the skin of the young domestic pig. *Nature.*, 193: 293-294.
- Winter, G.D. (1972) In: Maibach HI, Rovee DT, editors. *Epidermal wound healing*. Chicago: Year Book Medical Publishers. p.71.
- Winter, G.D., and Scales, J.T. (1963) The effect of air drying and dressings on the surface of a wound. *Nature.*, 197: 91.
- Wong, D.W.S. (1989) *Mechanism and theory in food chemistry*. New York: Van Nostrand Reinhold Company Inc.

References

- Wood, A., Ogawa, M., Portier, R.J., Schexnayder, M., Shirley, M., and Losso, J.N. (2008) Biochemical properties of alligator (*Alligator mississippiensis*) bone collagen. *Comp. Biochem. Physiol. Biochem. Mol. Biol.*, 151:246-249.
- Wright, K.A., Nadire, K.B., Busto, P., Tubo, R., Mc Pherson, J.M., and Wentworth, B.M. (1998) Alternative delivery of keratinocytes using a polyurethane membrane and the implications for its use in the treatment of full-thickness burn injury. *Burns*; 24(1):7-17.
- Wu, J. (2004) Assessment of effectiveness of oral administration of collagen peptide on bone metabolism in growing and mature rats. *J bone min met.*, 22(6): 547-553.
- Yamada, S., Nagaoka, H., Terajima, M., Tsuda, N., Hayashi, Y., et al. (2013) Effects of fish collagen peptides on collagen post-translational modifications and mineralization in an osteoblastic cell culture system. *Dent Mater J.*, 32: 88-95
- Yannas, and Burke, J.F. (1980) Design of an artificial skin. I. Basic design principles. *J Biomed Mater Res.*, 14(1):65-8.
- Yannas, Burke, J.F., Orgill, D.P., and Skrabut, E.M. (1982) Wound tissue can utilize a polymeric template to synthesize a functional extension of skin. *Science.*, 215:174-6.
- Yong-liang Zhuang, Xue Zhao, and Ba-fang Li. (2009) Optimization of antioxidant activity by response surface methodology in hydrolysates of jellyfish (*Rhopilema esculentum*) umbrella collagen. *J. Zhejiang Univ.-SCI. B* 10(8):572-579
- Yoshihito Ishida, Hiroshi Kubota, Akisugu Yamamoto, Akira Kitamura, Hans Peter Bachinger, and Kazuhiro Nagata. (2006) Type 1 collagen in Hsp47-null cells is aggregated in endoplasmic reticulum and deficient in N-propeptide processing and fibrillogenesis. *Molecular biology of the cell.* 17: 2346-2355.

References

- Yoshikawa, T., Tanaka, H., and Kondo, M. (2004) The increase in lipid peroxidation in rat adjuvant arthritis and its inhibition by Superoxide dismutase. *Biochem. Med*, 33:320-326.
- Zague, V *et al.*, (2011) Collagen Hydrolysate Intake Increases Skin Collagen Expression and Suppresses Matrix Metalloproteinase 2 Activity. *J. Med. Food.*, 14: 618-624
- Zague, V. A (2008) New view concerning the effects of collagen hydrolysate intake on skin properties. *Arch. Derm. Res*, 300: 479-483.
- Zhang, L., and Webster, T.J. (2009) Nanotechnology and nanomaterials: Promises for improved tissue regeneration. *Nano Today.*, 4: 66-80.
- Zhang, L.N., Guo, J., Zhou, J.P., Yang, G., and Du, Y.M. (2000) Blend membranes from carboxymethylated chitosan/alginate in aqueous solution. *J. Appl. Polym. Sci.*, 77(3): 610-616.
- Zhang, W., Liu, H.T. (2002) MAPK signal pathways in the regulation of cell proliferation in mammalian cells. *Cell Res.*, 12: 9–18.
- Zhang, Y., Kouguchi, T., Shimizu, K., Sato, M., Takahata, Y., & Morimatsu, F. (2010) Chicken collagen hydrolysate reduces proinflammatory cytokine production in C57BL/6.KOR-ApoEsh1 Mice. *J Nutr Sci Vitaminol.*, 56: 208-210.
- Zhang, Y., Liu, W. T., Li, G. Y., Shi, B., Miao, Y. Q., & Wu, X. H. (2007). Isolation and partial characterization of pepsin-soluble collagen from the skin of grass carp (*Ctenopharyngodon idella*). *Food Chem.*, 103: 906-912.
- Zhang, Z.K., Li, G.Y., and Shi, B., (2006) Physicochemical Properties of Collagen, Gelatin and Collagen Hydrolysate derived from Bovine Limed Split Wastes, *J. Soc. Leather Technol. Chem.*, 90: 23-28.

References

- Zheng, Y.Q., and Wei, W. (2005) Total glucosides of paeony suppresses adjuvant arthritis in rats and intervenes cytokine-signaling between different types of synoviocytes. *International Immunopharmacology.*, 5: 1560-1573.
- Zhong, S., and Yung, L.Y. (2009) Enhanced biological stability of collagen with incorporation of PAMAM dendrimer. *J. Biomed. Mater. Res. A.*, 91: 114-122.
- Zhongkai Zhang, Guoying Li and Bi Shi. (2005) Physicochemical properties of collagen, gelatin and collagen hydrolysate derived from bovine limited split wastes. *J Soc Leath Tech Ch.*, 90:23.
- Zhu, A.P., and Fang, N. (2005) Adhesion dynamics, morphology, and organization of 3T3 fibroblast on chitosan and its derivative: The effect of O-carboxymethylation. *Biomacromolecules.*, 6(5): 2607-2614.
