

REFERENCES

- Abdul, B. and Anderson, J. D. (1973). Vigor determination in soybean by multiple criteria. *Crop Sci.* 13: 630-635.
- Adetunji, I. A. (1991). Effect of harvest date on seed quality and viability of sunflower in semi-arid tropics. *Seed Science and Technol.*, 19: 571-580.
- Aher, R. P., Surve, P. P. and Dahat, D. V. (2000). Heterosis for yield and yield components in mungbean (*Vigna radiata* (L) Wilczek). *Agri. Science Digest.* 20, 1: 5-8.
- Ajala, M. O. (1999). Influence of seed quality attributes on field emergence of pigeon peas (*Cajanus cajan*, L.) and winged beans (*Psophocarpus tetragonolobus*, L.). *Journ. of Tropical Forest Res.* 15, 1: 140-147.
- Ali, M. G., Naylor, R. E. L. and Matthews, S. (2003). The effect of Ageing (using controlled deterioration) on the germination at 21 C as an indicator of physiological quality of seed lots of fourteen Bangladeshi rice (*Oryza sativa* L.) cultivars, *Pak. J. of Biol. Sci.* 6, 10: 910-917.
- Al-Jibouri, H. A., Miller, P. A. and Robinson, H. F. (1958). Genotypic and environmental variances in upland cotton cross of interspecific origin. *Agron. Journ.* 50: 633-636.
- Allard, R. W. (1960). Genetic basis of inbreeding depression and heterosis. In *Principles of Plant Breeding*. John Wiley and Sons, Inc., New York, pp. 224-232.
- Amaritsut, W. (2004). Development of Evaluation of Seed Viability and Vigor in Soybean by Tetrazolium Test. *Ph.D. Thesis*, Suranaree University of Technology, Thailand. pp.145-152. (Available online.)
- Amin, M. U. (1999). Influence of seed size on the performance of mungbean varieties under postrice and upland cropping systems. *Asian Regional Center-AVRDC Report*. Source: [www.arc.avrdc.rg/Pdf_Files/Amin\(17-N\)](http://www.arc.avrdc.rg/Pdf_Files/Amin(17-N)).
- Anuradha, R., Balamurugan, P., Srimathi, P. and Sumathi, S. (2009). Influence of seed size on seed quality of chickpea. (*Cicer arietinum* L.). *Legume Res.*, 32, 2 : 133-135.

- Arain M.A. (1990). Grain density in relation to seedling quality and crop establishment in rice (*Oryza sativa* L.). PhD Thesis University of the Philippines. pp.155. (Available online.)
- Association of Official Seed Analysts (AOSA) (2009). Seed Vigor Testing Handbook. Contribution No. 32 to the Handbook on Seed Testing.
- Association of Official Seed Analysts (AOSA). (1983). Seed vigor testing handbook. Rules for testing seeds. *Journ. of Seed Technol.* 12-13.
- Atak, M., Kaya, M. D., Kaya, G., Kaya, M. and Khawar, K. M. (2008). Dark green colored seeds increase the seed vigor and germination ability in dry green pea (*Pisum sativum* L.) *Pak. Journ. of Bot.* 40, 6: 2345-2354.
- Austin R. B. (1972). Effect of environment before harvesting on viability. In: Roberts EH, ed. Viability of seeds. London: Chapman and Hall, pp. 114-149.
- Balbaki R. Z. and Coopeland, L.O. (1987). Vigor testing of wheat and its relationship to field emergence, storage and seed quality. *Newsletter of AOSA.* 61:15.
- Bao, S., Yuhua, H., Zong, X., Wang, L., Lichi, L., Enneking, D., Rose, I.A., Leonforte, T., Redden, R.J., Paull, J. (2008). Collection of pea (*Pisum sativum*) and faba bean (*Vicia faba*) germplasm in Yunnan. *Plant Genet. Res. Newsl.* 156: 11-22.
- Barros, A. S. do R., and Marcos, F. J. (1997). Quick tests for vigor evaluation of soybean seeds. [Portuguese]. *Revista Brasileira de Sementes.* 19, 2: 289-295.
- Barros, D. I., Dias, D. C. F. dos S., Bhering, M. C., Dias, L. A. dos S., and Araujo, E. F. (2005). Use of tetrazolium test for physiological quality evaluation of squash seeds. [Portuguese]. *Revista Brasileira de Sementes.* 27, 2: 165-171.
- Basavarajappa, B. S., Shetty, H. S. and Prakash, H. S. (1991). Membrane deterioration and other biochemical changes, associated with accelerated ageing of maize seeds. *Seed Sci. & Technol.* 19: 279-286.
- Basbag, S., Ekinici, R. and Gencer, O (2007). Combining ability and heterosis for earliness characters in line x tester population of *Gossypium hirsutum* L. *Hereditas.* 144: 185-190.

- Baskaran, K. and Muthiah, A. R. (2007). Studies on combining ability in pigeonpea {*Cajanus cajan* (L.) Millsp.}. *Legume Res.* 30, 1: 67-69.
- Bastian Elli, D., Grosjean, F., Peyronnet, C., Duparque, M. and Regnier, J. M.(1998). Feeding value of pea (*Pisum sativum*, L.). Chemical composition of different categories of pea. *Anim. Sci.*, 67, 609–619.
- Basu, R. N. (1995). Seed viability. In Seed Quality. Basic Mechanisms and Agricultural Implications, ed. A. S. Basra (Food Products Press. An Imprint of The Haworth press, Inc. New York. London. Norwood (Australia), pp. 1-44.
- Bedell, P. E. (2001). Indian Forestry Species in Seed Science and Technology. Allied Publishers Limited. Mayapuri, New Delhi, India.pp. 112-130.
- Bento, S. R. S. de O., Santos, A. E. O. dos., Melo, D. R. M. de., and Torres, S. B. (2010). Efficiency of vigor tests to evaluate physiological quality of *Erythrina velutina* Wild. seeds. [Portuguese] *Revista Brasileira de Sementes.* 32, 4: 111-117.
- Bhupendra, S. and Gangwar, C. B. S. (2005). Seed vigour evaluation in mungbean. *Farm Sci. Journ.* 14, 1: 82-84.
- Birendra, K., Verma, S. K., Govind, R., and Singh, H. P. (2012). Temperature Relations for Seed Germination Potential and Seedling Vigor in Palmarosa (*Cymbopogon martinii*). *Journal of Crop Improv.* 26: 791-801.
- Bishnoi, U. R. and Delouche, J. C. (1980). Relationship of vigour tests and seed lots to cotton seedling establishment. *Seed Sci. & Technol.* 8: 341-346.
- Borah, H. K. (2009). Studies on combining ability and heterosis in field pea (*Pisum sativum* L.). *Leg. Res.* 32, 4: 255-259.
- Bradnock, W. T. and Matthews, S. (1970). Assessing field emergence potential of wrinkled-seeded peas. *Hort. Research.*, 10: 50-58.
- Bressani, R. and Elias, L. G. (1988). Seed quality and nutritional goals in pea, lentil, faba bean and chickpea breeding. In: R.J. Summerfield (ed.), World Crops: Cool Season Food Legumes. Kluwer Academic Publishers, Dordrecht, The Netherlands. pp. 381-404

- Brocklehurst, P. A. (1985). Factors affecting seed quality of vegetable crops. *Scientific Hort.* 36: 48-57.
- Burton, G. W. and Devane, E. H. (1953). Estimating heritability in tall Festuca (*Festuca aurandianacea*) from replicated clonal material. *Agron Journ.* 45: 478-481.
- Carver, M. F. F. and Matthews, S. (1975). Respiratory measurements as indicators of field emergence ability in peas. *Seed Sci. and Technol.* 3: 871-879.
- Castillo, A. G., Hampton, J. G. and Coolbear, P. (1994). Effect of sowing date and harvest timing on seed vigor in garden pea. (*Pisum sativum* L.). *New Zealand Journal of Crop and Hort, Science.*22: 91-95.
- Ceyhan, E., Avci, M. A. and Karadas, S. (2008). Line x tester analysis in pea (*Pisum sativum* L.): identification of superior parents for seed yield and its components. *Afri. Journ. of Biotech.* 7, **16**: 2810-2817.
- Chaudhary, D. K. and Sharma, R. R. (2003). Genetic variability, correlation and path analysis for green pod yield and its components in garden pea. *Ind. Journ. of Hort.* 60, **3**: 251-256
- Chin, H. F. (1988). Storage and vigour. *Seed Sci. & Technol.* 16: 1-4.
- Chuntarachud, T., Sagwansupyakorn, C., Subhadrabandhu, S., and Sripleng, A. (1984). Seed yield and seed quality of yard long bean [*Vigna unguiculata* (L.) Walp.sub sp. sesquipedalis (L.) Verdc.] at different harvesting stages. *Journ. of Agri. Sci.* 8: 123-127.
- Copeland, L. O. (1976). Principles of Seed Science and Technology. Burgess Pub. Comp., Minnaeapolis, Minnesota. pp. 164-165.
- Copeland, L. O. and M. B. McDonald. (1995). *Principle of Seed Science andTechnology.* Chapman & Hall, New York. pp 155-162.
- Coyne, C. J., McGee, R. J., Redden, R. J., Ambrose, M. J., Furman, B. J. and Miles, C. A. (2011). Genetic Adjustment to Changing Climates: Pea. In *Crop Adaptation to Climate Change*; Yadav, S. S., Redden, R.. J., Hatfield, J. L., Lotze-Campen, H. and Hall, A. E. Eds.; Wiley Blackwell: Chichester, UK. pp. 238 250.

- Davies, D. R., Berry, G. J., Heath, M. C. and Dawkins, T. C. K. (1985). Pea (*Pisum sativum* L.). In: R. J. Summerfield and E. H. Roberts, (eds.), Williams Collins Sons and Co. Ltd, London, UK. pp. 147-198.
- Delouche, J. C. and Baskin C. C. (1973). Accelerated aging techniques for predicting the relative storability of seed lots. *Seed Sci. & Technol.* 1: 427-452.
- Delouche, J. C. and Caldwell, W. P. (1960) Seed vigor and vigor tests. *Proceedings of the Association of Official Seed Analysts.* 50: 124-129.
- Devi, P. O., Pant, S. C., Rawat, S. S., Rana, D. K., and Singh, N. I. K. (2010). Correlation coefficient and genetic divergence analysis in pea. *Ind. Journ. of Hort.* 67: (Special Issue) 160-165.
- Dewey, D. H. and Lu, K. H. (1959). A correlation and path analysis of crested wheat grass seed production. *Agronomy Journal.* 51: 515-518.
- Dhaliwal, H. S. and Singh, K. B. (1970). Combining ability and inheritance of pod and cluster number in *Phaseolous mungo* (L). *Theor. Appl. Genet.* 40: 117-120.
- Duczmal, K. W. (1981). Mechanical injury in seeds. *Acta Hort. (ISHS)* 111: 235-242. (Abstract)
- Duke, J. A. (1981). Hand book of legumes of world economic importance. Plenum Press, New York. pp. 199-265
- Dutra, A. S. and Vieira R. D. (2006). Accelerated ageing test to evaluate seed vigor in pumpkin and zucchini seeds. *Seed Sci. & Technol.* 34: 209-214.
- Ellis, R. H. and Pieta-Filho C. (1992). The development of seed quality in spring and winter cultivars of barley and wheat. *Seed Sci. Res.* 2: 9-15.
- Ellis, R. H. and Roberts E. H. (1980). Towards a rational basis for testing seed quality. *Seed production.* pp: 605-635.
- Ellis, R. H., Hong T. D., and Roberts E. H. (1987). The development of desiccation tolerance and maximum seed quality during seed maturation in six grain legumes. *Annals of Bot.* 59: 23-29.
- Ellis, T. H. N., Hofer, J. I., Timmerman-Vaughan, G. M., Coyne, C. J. and Hellens, R. P. (2011). Mendel, 150 years on. *Trends Plant Sci.* 16: 590-596.

- Espinosa, N. and Ligarreto, G. A. (2005). Evaluating combinatory ability and heterosis of seven parental *Pisum sativum* L. pea lines. (Spanish). *Agronomia Colombiana*. 23, 2: 197-206.
- Fansecò, S and Peterson F. L. (1968). Hybrid vigor in seven parent diallel crosses in common wheat (*T.aestivum* L.). *Crop Sci*. 8: 85-88.
- FAO. (2012). Food and Agriculture Organization of the United Nations, (FAOSTAT). FAO Statistical Databases. Available at <http://faostat.fao.org/default.aspx>.
- Farshadfar, M. and Farshadfar, E. (2008). Genetic variability and path analysis of chickpea (*Cicer arietinum* L.) Landraces and Lines. *Journ. of Appl. Sci*. 8, 21: 3951-3956.
- Fisher, G. and Yates, M. (1963). Statistical Tables for Biological, Agricultural and Medical Research. 6: 62-63.
- Frank, W. J. (1950). Address to the Association of Official Seed Analysts. Proceedings of the Association of Official Seed Analysts. 40: 36-39.
- Ganesh M, Singh U. P., Srivastava C. P. and Sarode, S. B. (2008). Genetic basis of heterosis for yield and related traits in pea (*Pisum sativum* L.). *Asian Journ. of Bio Sci*. 3, 1: 73-76.
- Gassim, A. A. E. (1986). Seed Quality and Vigor. Papers presented at the Seed Technology Course Cairo, Egypt. International Center for Agricultural Research in the Dry Areas ICARDA Box 5466, Aleppo, Syria pp.15-32.
- Gautam, G. V. and Vermani, A. (2005). Heterobeltiosis studies in garden pea (*Pisum sativum* L.). *Udyanika (Journ. of Hort. Sci.)*. 11, 2: 97-99.
- Gelmond, H., I. Luria, Woodstock, L. W. and Perl, M. (1978). The effect of accelerated aging of sorghum seeds on seedling vigour, *Journ. of Experimental Botany*. 29, 109: 489-495.
- Glewen K. L. and Vogel K. P. (1984). Partitioning the genetic variability for seedling growth in sand bluestem and its seed size and seedling vigor components. *Crop Sci*. 138, 24 : 200-255.

- Grando, S. and Ceccarelli, S. (1995). Seminal root morphology and coleoptiles length in wild (*Hordeum vulgare* ssp. *sontaneum*) and cultivated (*Hordeum vulgare* ssp. *vulgare*) barley. *Euphytica*, 86: 73-80.
- Gritton, E. T. (1980). Field Pea. Hybridization of Crop Plants. In: W.R. Fehr and H.H. Hadley (eds.), American Society of Agronomy, Inc., and Crop Science Society of America, Inc., Wisconsin, USA pp. 347-356.
- Gupta A. J., Singh, Y. V., Hussain, K., Bhat, S. A., Bhat, F. N. and Singh, T. K. (2007). Character association and selection parameters in garden pea (*Pisum sativum* L.). *Environ. and Ecolo.* 25 (S) 2: 331-335.
- Gupta A. J., Singh Y. V. and Verma T. S. (2006). Genetic variability and heritability in garden pea (*Pisum sativum* L.). *Ind. Journ. of Horti.* 63, 3: 332-334.
- Hadavizadeh, A. and George, R. A. T. (1989). The effect of mother plant nutrition on seed yield and seed vigour in pea (*Pisum sativum* L.) cultivar 'sprite'. *Acta Horticulturae* 253: 55-61.
- Hampton, J. G. (1994). Methods of viability and vigor testing; a critical appraisal. In Basra A.S.Ed. Basic Mechanisms and Agriculture Impelmentations. Food Proudcts Press. NewYork, USA. pp. 81-118.
- Hampton, J. G. and Coolbear, P. (1990). Potential versus actual seed performance - Can vigour testing provide an answer? *Seed Science and Technolog.* 18: 215-228.
- Hampton, J. G. and Scott, D. J. (1982). Effect of seed vigour on garden pea production. *New Zealand Journ. of Agri. Res.* 25: 289-294.
- Hampton, J. G., Brunton, B. J., Pemberton, G. M., Rowarth, J. S. and Rowarth, J. S. (2004). Temperature and time variables for accelerated ageing vigour testing of pea (*Pisum sativum* L.) seed. *Seed Sci. & Technol.* 32: 261-264.
- Hanson C. H., Robinson, S. and Comstock, R. E. (1956). Biometrical studies of yield in segregating populations of Korean Leaspedeza. *Agron. Journ.* 48: 268-272.
- Harinder S., Singh, M. and Brar P. S. (2005). Heterosis and combining ability studies for economic traits in genetically diverse lines of garden pea (*Pisum sativum* L.). *Crop Improvement.* 32, 1: 78-85.

- Harrington, J. F. (1971). The necessity of high quality vegetable seeds. *Hortscience*. 6: 550-551.
- He, C., Liu, Y., Wu, K., Yuan, M., Feng, Q., Liu, Y., Yan, Q., Guan, J., Rose, I. A. and Redden, R.J. (2008). Collecting and surveying landraces of pea (*Pisum sativum*) and faba bean (*Vicia faba*) in Qinghai province of China. *Plant Genet. Res. Newsl.* 156: 1-10.
- Heydecker, W. (1969). The vigour of seed. A review. Proceedings of the International Seed Testing Association. 34: 201-219.
- Heydecker, W. (1972). Vigor In; Viability of seeds Roberts, E.H. Chapman Hall, London, pp 209-252.
- Hiltner, L., and Ihssen, G. (1911). Uber das schlechte Auflaufen and die Auswinterung des Getreides infolge Befall isdes Saatgutes duveti *Fusarium* Landw. Jahobuch fur Bayern. 1: 20-60.
- Hopper, N. W., Overholt, J. R. and Martin, J. R. (1979). Effect of cultivar, temperature and seed size on the germination and emergence of soybeans (*Glycine max* L. Merr.). *Ann. of Bot.* 8, **44**: 301.
- Horcika, P. and Hosnedi, V. (1997). The effects of sowing and harvest terms on the seed quality of pea. *Scientia Agriculturae Bohemica*. 28, **4**: 263-270.
- Hu, W. J., Harding, S. A., Lung, J., Popko, J. L., Ralph, J., Stokke, D. D., Tsai, C. J., and Chiang, V. L. (1999). Repression of lignin biosynthesis promotes cellulose accumulation and growth in transgenic trees. *Nat. Biotechnol.* 7: 808-812.
- Hulse, J. H. (1994). Nature, composition and utilization of food legumes. In: F. J. Muehlbauer and W. J. Kaiser (eds.), Expanding the Production and Use of Cool Season Food Legumes. Kluwer Academic Publishers. Dordrecht, The Netherlands. pp. 77-97.
- Hund, A., Fracheboud, Y., Soldati, A., Frascaroli, E., Salvi, S. and Stamp, P. (2004). QTL controlling root and shoot traits of maize seedlings under cold stress. *Theor. Appl. Genet.* 109: 618-629.
- Ilbi, H., Kavak, S. and Eser, B. (2009). Cool germination test can be an alternative vigour test for maize. *Seed Sci. and technol.* 37, **2**: 516-519.

- International Seed Testing Association (ISTA) (1987). Handbook of Vigor Test Methods. 2nd ed., *International Seed Testing Association*, Zurich, Switzerland.
- International Seed Testing Association (ISTA). (2010). International rules for seed testing. Zurich, Switzerland: International Seed Testing Association.
- International Seed Testing Association. (1995). Handbook of Vigour Test Methods. 3rd ed.. Internaitonal Seed Testing Association. Zurich. Switzerland.
- Iqbal, T. M. T. and Smith, M. I. (1996). Physiological changes of pea seed quality due to aging. *Annals of Bangladesh Agriculture*. 6, 1: 17-34.
- Isely, D. (1957). Vigour Tests. Proceeding of the Association of Official Seed Analysts. 47: 176-182.
- Ishrat, A. L., Gaurav, S. S., Bahuguna, D. K. and Anwar A. (2012) {B}. Relationship between vigor traits and field mergence in rice under temperate conditions. *Progressive Research*.7, 1 : 112-116.
- Ishrat, A. L., Gaurav, S. S., Bahuguna, D. K., Ali, A. and Singh, P. B. (2009). Relationship among seed yield attributes in rice under temperate conditions. *Ind. Journ. of Plant Genet. Res.* 22, 1: 80-83.
- Ishrat, A. L., Tyagi, S. D., Bahuguna, D. K., Rajeev, K. and Nitin, K. (2012) {A}. Studies on seed vigor deterioration in pea (*Pisum sativum* L.). *HortFlora Research Spectrum.*, 1, 4: 328-333.
- Isley, D. (1950). Vigor Tests. *Proc. Assoc. Off. Seed Anal.* 47: 176-182.
- Jahagirdar, J. E. (2003). Line x tester analysis for combining ability in pigeonpea. *Ind. Journ. of Pulses Res.* 16, 1: 17-19.
- Jitendra, K., Nazima, A. and Krishan, P. (2010). Variability and character association in garden pea (*Pisum sativum* L. sub sp. *hortense* Asch. and Graebn.). *Progressive Agri.* 10, 1: 124-131.
- Johnson, H. W., Robinson, H. F. and Comstock, R. E. (1955). Estimates of genetic and environmental variability in soybean. *Agrono. Journ.* 47: 314-318.

- Johnson, T. E. and Hutchinson, E. W. (1993). Absence of strong heterosis for life span and other life history traits in *Caenorhabditis elegans*. *Genetics* 134: 465-474.
- Jones, D. B. and Peterson, M. L. (1976). Rice seedling vigor at sub optimal temperatures. *Crop Sci.* 16: 102-105.
- Joshi, A. B. and Dhawan, N. L. (1966). Genetic improvement of yield with special reference to self fertilizing crops. *Ind. Journ of Genetics.* 26 (A): 101-113.
- Jyothula, D. P. B. and Guttala, S. B. (2009). Combining ability analysis for seed yield and for some quantitative characters in field pea (*Pisum sativum* L.). *Crop Res. (Hisar).* 37, **1/3**: 174-180.
- Kaloo, G., Mathura, R., Jagdish, S., Ajay, V., Rajesh, K., and Rai, G. K. (2005). Morphological and biochemical variability in vegetable pea (*Pisum sativum* L.). *Vegetable Sci.* 32, **1**: 19-23.
- Kanade, P. B., Harer, P. N. and Kshirsagar, A. N. (2010). Correlation and path analysis studies in pigeonpea (*Cajanus cajan* (L.) Millsp.). *Journ. of Maharashtra Agri. Univs.* 35, **3**: 494-496.
- Kant, K., Sharma B., and Tyagi, M. C. (1973). Effects of maturation environment on seed size and subsequent plant growth in peas (*Pisum sativum*). *Agron. Journ.* 65: 390-394.
- Kaptsimadi, C. M., Raegen, O. and Hoften, H. (1990). Growth of four cucumber (*Cucumis sativus* L.) cultivars at suboptimal temperatures and storage behavior of their fruit at different temperatures. *Acta Horticulturae.* 287: 375-383.
- Kay, D. (1979). Food legumes. Tropical Products Institute (TPI). *TPI Crop and Product Digest* No. 3, London UK. pp. 26-47.
- Kempthorne, O. (1957). An introduction to Genetic Statistics. New York, John Wileys and Sons, Inc., London. pp. 1-545.
- Khalil, S. K., Mexal, J. G. and Murray, L. W. (2001). Soybean seed matured on different dates affect seed quality. *Pak. Journ. of Biol. Sci.* 4, **3**: 365-370.

- Khan, A. A., Lubna, S., Kawser-e-J., Mian, I. H. and Akanda, M. A. M. (2006). Effect of physical condition of pea seed on prevalence of seed-borne fungi, germination and seedling vigour. *Bangladesh Journ. of Pl. Pathol.* 22, 1/2: 85-89.
- Khan, A., Jan, A., Bashir, S. and Noor, M. (2005). Effect of nitrogen and seed size on maize crop stand and plant height. *Journ. of Agri. & Soc. Sci.* 1, 4: 380-381.
- Khattra, S., Kaur, A. J., and Singh, G. (1997). Comparison of seed quality at physiological maturity and harvest maturity of two pigeon pea (*Cajanus cajan* L. and Millsp) cultivars. *Acta Agrobotanic.*, 50 (1-2): 49-52.
- Komba, C. G., Brunton, B. J. and Hampton, J. G. (2006). Accelerated ageing vigour testing of kale (*Brassica oleracea* L. var. *acephala* DC) seed. *Seed Sci. & Techol.* 34: 205-208.
- Krishnan, P. and Rao, A. V. S. (2005). Effects of genotype and environment on seed yield and quality of rice. *The Journ. of Agri. Sci.* 143: 283-292.
- Krishnasamy, V. and Seshu, D. V. (1990). Accelerated aging in rice. *Seed Sci. & Techol.* 18: 147-156.
- Kuldeep, T., Myoung, P., Hyo, J. L., Chong, A. L., Shafiq, R., Brian, S., Kui, J. L. and Song, J. Y. (2011). Diversity for seedling vigor in wild barley (*Hordeum vulgare* L. Subsp. *Spontaneum*) germplasm. *Pak. Journ. of Bot.*, 43, 3: 2167-2173.
- Kumar, C. V. S., Sreelakshmi, C. H. and Verma, P. K. (2009). Studies on combining ability and heterosis in pigeon pea (*Cajanus cajan* L.). *Legume Res.* 32, 2: 92-97.
- Kumar, S., Singhal, N. C. and Prakash, S. (1997). Inter varietal variability for seed longevity in pea (*Pisum sativum* L.). *Ind. Journ. of Genet. and Pl. Breeding.* 57, 2: 204-209.
- Kumar, V. R. A. and Sharma, R. R. (2006). Path coefficient analysis for pod yield and its attributes in garden pea (*Pisum sativum* L. sub sp. *hortense*). *Ann. of Agri. Res.* 27, 1: 32-36.

- Lal, G. M., Meena, M. L., Kunj, C. and Singh, C. M. (2011). Assessment of genetic variability and interrelation between yield and its contributing components in field pea (*Pisum sativum* L.). *Environ. and Ecol.* 29, **3** (A): 1235-1239.
- Larson, S. U., Povlsen, F. V., Eriksen, E. N. and Pedersen, H. C. (1998). The influence of seed vigor on field performance and the evaluation of the applicability of the controlled deterioration vigor test in oil seed rape (*Brassica napus*) and pea (*Pisum sativum* L.). *Seed Sci. & Technol.* 26 **3**: 627-641.
- Lee, L. S., Lee, L. V. and Russell, T. E. (1986). Aflatoxin in Arizona cotton seed, field inoculation of bolls by *Aspergillus flavus* spores in wind driven soil. *Journ. of Amer. Oil Chem Soc.* 63: 530- 532.
- Lila, B., Vinod, K. and Maurya, S. K. (2009). Hybrid breeding for green pod quality, yield and its components in garden pea (*Pisum sativum* L.). *Annals of Hort.* 2, **2**: 161-165.
- Lohithaswa, H. C. and Dharmaraj, P. S. (2003). Implications of heterosis, combining ability and per se performance in pigeonpea. *Karnataka Journ. of Agri. Sci.* 16, **3**: 403-407.
- Lois, R., Manuel, D., Karine, G., Julie, C., Julia, B., Claudette, J., and Dominique, J. (2012). Seed germination and vigor. *Annu. Rev. of Plant Biol.* 63: 507- 533.
- Lush, J. L. (1940). Intra-sire correlations or regression of offspring on Dam as a method of estimating heritability of characteristics. *Proc. Am. Soc. Animal Prod.* 33: 293-301.
- Machado, C. G., Martins, C. C., Santana, D. G., De-Cruz, S. C. S. and Oliveira, S. S. C. de. (2011). Suitability of electrical conductivity test for seeds of *Pisum sativum* subsp. *Arvense*. *Ciencia Rural.* 41, **6**: 988-995.
- Maguire, J. D. (1962). Speed of germination-aid in selection and evaluation for seedling emergence and vigor. *Crop Sci.* 2: 176-177.
- Mahajan, J. P., Dumbre, A. D., and Bhingarde, M. T. (1998). Effect of environments, fertilizers and plant density on seed yield and quality of pigeon pea. *Journal of Maharashtra Agricultural Universities*, 22, **2**: 151-154.
- Makashera, R. (1973). The Pea. Oxonian Press Pvt. Ltd., New Delhi. pp. 50-110.

- Makasheva, R. Kh. (1983). *The Pea*. Oxonian Press Pvt. Ltd., New Delhi, India. pp. 267.
- Makkawi, M., El-Balla, M., Bishaw, Z., and Gastel, A. (1999). The relationship between seed vigour tests and field emergence in lentil (*Lens culinaris* Medikus). *Seed Sci. and Technol.* 27, 2 :657-668.
- Makkawi, M., El-Balla, M., Bishaw, Z., and Gastel, A. (2008). Correlation and path coefficient analyses of laboratory tests as predictors of field emergence in Lentil (*Lens culinaris* Medikus). *Journ. of New Seeds.* 9, 4: 284-302.
- Makoto, N., Kazunori, T., Hiroyuki, U., Erika, A., Shunsuke, I., Chiaki, M., and Hiroshi, E. (2012) Low sodium chloride priming increases seedling vigor and stress tolerance to *Ralstonia solanacearum* in tomato. *Plant Biotechnology.* 29: 9-18.
- Mark, A. Bennett (2011). Vigor and seed performance. In Consortium for inrenational seed technology training; Seed Biology Program Dept. of Horticulture and Crop Science, Ohio State University Columbus. pp 1-22.
- Matthews, S. (1980). Controlled deterioration, a new vigor test for crop seeds. In P.D. Hebblethwaite ed. *Seed production*. Butterworths London. pp. 647-660.
- Matthews, S. and Bradnock, W. T. (1968). Relationship between seed exudation and field emergence in peas and French beans. *Hort. Research.* 8: 89-93
- Matthews, S., Powell, A. A., and Rogerson, N. E. (1980). Physiological aspects of the development and storage of pea seeds and their significance to seed production. In *Seed Production*. ed. P. D. Hebblethwaite (London, UK: Butterworths), pp. 513-525.
- Maurya, C. L., Khan, A. A., and Yadav P. S., (2006). Studies on various seed vigour tests and their correlation with field emergence in oat (*Avena sativa* L.). *Farm Sci. Journ.* 15, 1: 73-74.
- Mc-Donald, Jr. M. B. (1975). A review and evaluation of seed vigor tests. *Proc. Assoc. Off. Seed Anal.* 65: 109-139.
- McDonald, M. B. J. (1977). A review and evaluation of seed vigour tests. *Proceedings of the Association of Official Seed Analysts*, 65: 109-139.

- Mckersie, B. D. and Tomer, D. T. (1982) A comparison of seed quality and seedling vigor in Birdsfoot Trefoil. *Crop Sci.* 22, 6 : 1239-1241.
- Meena, H. S., Kumar, J. and Yadav, S. S. (2004). Genetics of seed shape in chickpea (*Cicer arietinum* L.). *Annals of Agri. Res.* 25, 3: 439-441.
- Miguel, M. H. and Cicero, S. M. (1999).[A], Cold test for soybean seed vigor evaluation. (Portuguese). *Revista Brasileira de Sementes.* 21, 2: 35-42.
- Miguel, M. H. and Cicero, S. M. (1999).[B] Cold test for evaluating bean seed vigour. (Portuguese). *Scientia Agricola.* 56, 4 (S): 1233-1243.
- Miller, B. McDonald (2010). Standardization of seed vigor tests. In Seeds, trade, production and technology. Seed Biology Program Dept. of Horticulture and Crop Science, Ohio State University Columbus. pp 200-207.
- Modarresi, R., Rucker, M. and Tekrony, D. M. (2002). Accelerating ageing test for comparing wheat seed vigour. *Seed Sci. & Technol.* 30: 683-687.
- Mohan, S., Yuvinder, K., Harinder, S. and Brar, P. S. (2005). Correlation and path coefficient analysis in garden pea (*Pisum sativum* L.). *Environ. and Ecolo.* 23, 2: 315-318.
- Muasya, R. M., Lommen, W. J. M., and Struik, P. C. (2002). Differences in development of common bean (*Phaseolus vulgaris* L.) crops and pod fraction within a crop II. Seed viability and vigour. *Field Crops Research.* 75: 79-89.
- Muehlbauer, F. J. (1991). Incorporation of useful characters from germplasm resources into cultivars of food legumes. In: (A. Smith and R.S. Robertson (eds.), Legume Genetic Resources in the Semi-Arid Temperate Environments. ICARDA, Aleppo, Syria. pp. 363-393.
- Murillo-Amdor, B., Lopez-Aguilar, R., Kaya, C., Larrinaga-Mayoral, J. and Flores-Hernandez, A. (2002). Comparative effect of NaCl and polyethylene glycol in germination, emergence and seedling growth in cowpea. *Crop Sci.* 188: 235-247.
- Na Chiangmai, P., Laosuwan, P., and Waranyuwat, A. (2007). The effect of mungbean seed size on germinating ability, bean sprout ability and agronomic characters. Source: www.journal.sut.ac.th/index.php/suij/article/view/13/11.

- Naidu, N. V. and Satyanaryana, A. (1993). Heterosis and combining ability in mungbean (*Vigna radiata* (L) Wilczek). *Ind. Journ of Pulse Research*. 6, 1: 38-44.
- Nandpuri, K. A., Kumar, J. C. and Singh, H. (1973). Hertibility and interrelationship of some quantitative characters in pea (*Pisum sativum* L.). *Journ. of Research (PAU)*. 10: 309-315.
- Narwal, A. K., Pandita, M. L., Malik, Y. S. and Khurana, S. C. (2004). Correlation studies between laboratory tests and field parameters in okra. *Haryana Journ. of Hort. Sci.* 33, 3/4 : 250-253.
- Nascimento, W. M. (1994). Effect of selection on pea (*Pisum sativum* L.) seed quality. *Pesquisa Agropecu. Bras.* 29, 2: 309-313.
- Nascimento, W. M., Freitas, R. A., de Gomes, E. M. L. and Soares, A. S. (2007). Accelerated aging test on pea seeds. (Portuguese). *Horticultura Brasileira*. 25, 2: 205-209.
- Nawab, N. N., Subhani, G. M., Khalid, M., Qamar, S. and Akhtar, S. (2008). Genetic variability, correlation and path analysis studies in garden pea (*Pisum sativum* L.). *Journ. of Agri. Res.* 46, 4: 333-340.
- Nichols, M.A., Warrington, I. J. and Scott, D. J. (1978). Pre-harvest treatment effects on some quality criteria of pea seeds. *Acta Hort.* 83: 113-124.
- Nienhuis, J., Lower, R. L. and Staub, J. E. (1983). Selection for improved low temperature germination in cucumber. *J. Amer. Soc. Hort. Sci.* 108, 6: 1040-1043.
- Nonogaki, H., Bassel, G. W., Bewley, J. D. (2010). Germination still a mystery. *Plant Sci.* 179: 574-81
- Ojo, D. K. and Ariyo, O. J. (2000). Assessment of seed quality of soybean genotypes sown in three contrasting tropical field ecologies using the regression and rank-sum approaches. *Moor Journ. of Agri. Res.* 1, 1 : 16-21.
- P. G. R. O. (1981). Electrical conductivity test for vining pea seed. In Processors and Growers Research Organization information sheet number 119, Peterborough, UK.

- Pandey, N. (2004). Line x tester analysis in long-duration hybrid pigeon pea. *Legume Res.* 27, 2: 79-87.
- Pandita, V. K. and Shantha, N. (2002). Germination behaviour and field performance of garden pea (*Pisum sativum* L.) in relation to seed ageing. *Ind. Journ. of Agri. Sci.* 72, 4: 213-215.
- Pankaj B., Chaturvedi, S. K. and Awasthi, N. N. C. (2003). Character association and path coefficient analysis in chickpea. *Annals of Agri. Res.* 24, 3: 684-685.
- Panse, V. G. and Sukhatme, P.V. (1942). *Statistical Methods for Agricultural Workers*. Ind. Coun. of Agri. Res. Publication, New Delhi.
- Parera, C. A., Cantliffe, D. J., Stoffella, P. J. and Scully, B. T. (1995). Field emergence of shrunken-2 Corn Predicted by single and Multiple-vigour laboratory Tests. *Journ. of Am. Soc. Hort. Sci.*, 120, 1: 128-132.
- Parimal, B. and Chakraborty, S. K. (2009). Studies on seed parameters and seed deterioration in *Vigna radiata* L. *Advn. in Pl. Sci.* 22, 1: 89-91.
- Peacock, H. A. and Hawkins, B. S. (1970). Effect of seed source on seedling vigor, yield and lint characteristics of cotton (*Gossypium hirsutum* L.). *Crop Sci.* 10: 667-670.
- Peiffer, R. A., McKee, G. W., and Risiu, M. L. (1972). Germination and emergence of crownvetch as affected by seed maturity and depth of planting. *Agron. Journ.* 64: 172-714.
- Peksen, E. Peksen, A. Bozoglu, H. and Gulumser, A. (2004). Some seed traits and their relationships to seed germination and field emergence in pea (*Pisum sativum* L.). *Journ. of Agron.* 3, 4: 243-246.
- Perry, D. A. (1973). Interacting effects of seed vigour and the environment. In W. Heydecker (ed) *Seed ecology*. Butterworths London. pp. 311-323.
- Perry, D. A. (1980). The concept of seed vigor and its relevance to seed production techniques. In *Seed Production*, ed. P. D. Hebblethwaite (London, UK: Butterworths), pp. 585-591.
- Peterson, J. (1975). Photosynthetic productivity of tall and dwarf varieties of pea (*Pisum sativum* L.) *Rostilinna Vyroba* 21, 8: 885-895.

- Pollock, B. M. and Roos, E. E. (1972). Seed and seedling vigour. In. Seed Biology Vol.1 (Ed. Kozlowski, T.). pp: 314-318.
- Powell, A. A., Oliveira, M. de A. and Matthews, S. (1986). The role of imbibition damage in determining the vigour of white and coloured seed lots of dwarf French bean (*Phaseolus vulgaris*). *Journ. of Exp. Botany*. 37: 716-722.
- Pownall, T. L., Udenigwe, C. C., and Aluko, R. E. (2010). Amino acid composition and antioxidant properties of pea seed (*Pisum sativum* L.) enzymatic protein hydrolysate fractions. *Journ. of Agri. Food Chem.* 58, 8: 4712-4718.
- Raju, P. A. and Muthiah, A. R. (2007). Heterosis and combining ability for quantitative and qualitative traits in pigeonpea [*Cajanus cajan* (L.) Millsp.]. *Res. on Crops*. 8, 3: 625-631.
- Rech, E. G. Villela, F. A. and Tillmann, M. A. A. (1999). Quick evaluation of the physiological quality of pea seeds. *Revista Brasileira de Sementes*. 21, 2: 1-9.
- Regan, K. L., Siddique, K. H. M., Turner, N. C. and Whan, B. R. (1992). Potential for increasing early vigor and total biomass in spring wheat. Characteristics associated with early vigor. *Aust. Journ. Agric. Res.* 43: 541-553.
- Revilla, P., Butron, A., Malvar, R. A. and Ordas, A. (199). Relationships, among kernel weight, early vigor, and growth in maize. *Crop Sci.* 39: 654-658.
- Roosrokh, M., Golozani, K. G. and Javanshir, A. (2002). Relationship between seed vigour and field performance in chickpea (*Cicer arietinum* L.). *Seed and Plant*. 18, 2: 156-169.
- Sachan, C. P., Singh, H. C. and Kumar, S. (2001). Genetic analysis of various yield components in field pea. *Ind. Journ. of Pulses Res.* 14, 2: 150-151.
- Salam, J. L., Kashyap, O. P., Pandey, R. L., Nag, S. K. and Ranjan, S. K. (2007). Analysis of genetic architect of yield and its component in field pea (*Pisum sativum* L.). *Pl. Archives*. 7, 1: 229-231.
- Sardana, S., Mahajan, R. K., Gautam, N. K. and Ram, B. (2007). Genetic variability in pea (*Pisum sativum* L.) germplasm for utilization. *SABRAO Journ. of Breed. and Genet.* 39, 1: 31-41.

- Sarode, S. B., Kamal, G. and Srivastava, C. P. (2009). Line x tester analysis in pea (*Pisum sativum* L.). *International Journ. of Pl. Sci.* 4, **1**: 233-236.
- Sarwat, P., Stoddart, F. L., Marshall, D. R. and Ali, S. M. (1994). Genetic analysis of different pulse crops. *Ephytica*. 80, **1-2**: 39-48.
- Sasahara, T., Ikarashi, H. and Kambayashi, M. (1986). Genetic variations in embryo and endosperm weights, seedling growth parameters and α -amylase activity of the germinated grains in rice (*Oryza sativa* L.). *Japan. Journ. of Breed.* 36: 248-261.
- Satyawan, A., Malik, B. P. S., Ram, K. and Ram, D. (2004). Variability, correlation and path analysis in field pea (*Pisum sativum* L.). *Haryana Agri. Univ. Journ. of Res.* 34, **2**: 149-153.
- Saumya, A., Lavanya, G. R. and Babu, G. S. (2011). Character association and component analysis in garden pea (*Pisum sativum* L. hortense). *Environ. and Ecol.* 29, **1**: 133-135.
- Saumya, A., Lavanya, G. R. and Rashmi, J. (2009). Heterosis estimates of garden pea crosses (*Pisum sativum* L. hortense). *Trends in Biosci.* 2, **2**: 45-47.
- Sawicki, J. and Boros, L. (2000). Gene action in inheritance of some traits related to plant productivity in pea. *Biuletyn Instytutu Hodowli i Aklimatyzacji Roslin.* 216, **2**: 407-415.
- Schuab, S. R. P., Braccini, A. L., Scapim, C. A., Franca-Neto, J. B., Meschede, D. K., and Avila, M. R. (2007). Germination test under water stress to evaluate soybean seed vigour. *Seed Sci. and Technol.* 35, 1: 187-199.
- Scott, D. J. and Close, R. C. (1976). An assessment of seed factors affecting field emergence of garden pea seed lots. *Seed Sci. and Technol.* 4: 287-300.
- Seshu, D. V., Krishnasamy, V. and Siddique, S. B. (1987). Seed vigor in rice. In S. J. Banta, ed. *Proceeding of International Workshop on Rice Seed Health*. 16-20 March, 1987. IRRI. pp. 315- 327.
- Shephard, H. L., Naylor, R. E. L. and Stuchbury, T. (1996). The influence of seed maturity at harvest and drying method on the embryo, α -amylase activity and seed vigour in sorghum [*Sorghum bicolor* (L.) Moench]. *Seed Sci & Technol.* 24: 245-259.

- Shukla, P. S., Atul, K. and Shambhoo, P. (2009). Correlation analysis of seed yield and vigour parameters of vegetable pea varieties under moisture stress condition. *Trends in Biosciences*. 2, 1: 65-66.
- Shukla, Y. K., Kohli, U. K. and Rastogi, K. B. (1994). Seed vigor and its significant in garden pea cultivars. *South Indian Horticulture*. 42, 60: 388-389.
- Singh J. D. and Singh I. P. (2005). Studies on correlation and path coefficient analysis in field pea (*Pisum sativum* L.). *National Journ. of Pl. Improv.* 7, 1: 59-60.
- Singh, A. K. and Yadav, V. K. (2005). Character association and path coefficient analysis studies for certain metric traits in pea (*Pisum sativum* L.) under cold arid region of Ladakh. *Hort. Journ.* 18, 1: 35-38.
- Singh, B., Kumar, N. and Joshi, S. (1978). Genotypic and phenotypic variability in table pea. *Proc. Hort.* 10, 2: 55-61.
- Singh, H., Singh, M., Singh, S., and Brar, P. S. (2006). Estimation of gene effects of yield and quality characters in pea (*Pisum sativum* L.). *Environ. and Ecol.* 24, 2: 445-448.
- Singh, J. D. and Singh, I. P. (2006). Genetic variability, heritability, expected genetic advance and character association in field pea (*Pisum sativum* L.). *Legume Res.* 29, 1: 65-67.
- Singh, V. P. and Singh, I. P. (1991). Combining ability for yield and nodulation traits in mungbean (*Vigna radiata* (L) Wilczek). Golden Jubilee symposium. Genet. Res. Edu. Current trends and the next fifty years, New Delhi. *Indian Abs. II:* 485-489.
- Singh, V. P., Pathak, M. M. and Singh, R. P. (1994). Combining ability in pea. *Ind. Journ. of Pulse Research.* 7, 1 : 11-14.
- Singhabumrung, V. and Juntakool, S. (2004). Vigor test results for prediction of field emergence for sweet corn. pp.291-299. Available online on www.lib.ku.ac.th
- Sirohi, S. P. S., Ramashray, Y. and Sanjai, M. (2006). Genetic variability, correlations and path coefficient analysis for seed yield and its component characters in pea (*Pisum sativum* L.). *Plant Archives.* 6, 2: 737-740.

- Sirohi, S. P. S., Ramashray, Y., Sanjai, M., and Dubey, R. K. (2007). Association and path coefficient analysis in field pea (*Pisum sativum* var. arvence). *Ad. in Plant Sci.* 20, 2: 603-607.
- Smart, J. (1990). Grain Legumes: Evolution and genetic resources. Cambridge University Press, Cambridge, UK. pp. 200-201
- Smiciklas, K. D., Mullen, R. E., Carlson, R. E. and Knapp, A. D. (1989). Drought induced stress effects on soybean seed calcium and seed quality. *Crop Sci.* 29: 1519-1523.
- Smith, O. E. Welch, N. C. and McCoy, O. D. (1973). Studies on lettuce seed quality. II. Relationship of seed vigor to emergence, seedling weight and yield. *Journ. of Am. Soci. of Horti. Sci.* 98, 3: 552-556.
- Smykal, P., Kenicer, G., Flavell, A. J., Corander, J., Kosterin, O., Redden, R. J., Ford, R., Coyne, C. J., Maxted, N., and Ambrose, M. J., (2011). Phylogeny, phylogeography and genetic diversity of the *Pisum* genus. *Plant Genet. Res.* 9: 4-18.
- Spears, J. F., TeKrony, D. M. and Egli, D. B. (1997). Temperature during seed filling and soybean seed germination and vigour. *Seed Sci. & Technol.* 25: 233-244.
- Sprague, G. F. and Tatum L. A. (1942). General vs specific combining ability in single crosses of corn. *Journ. of Ameri. Soci. of Agrono.*, 34: 923-932.
- Sridhar, G. and Nagaraja A. (2004). Studies on relationship between vigour tests and field emergence. *Mysore Journ. of Agri. Sci.* 38, 4: 551-553.
- Summerfield, R. J. and Roberts, E. H. (1985). Recent trends in internationally oriented research on grain legumes. In: R. J. Summerfield and E. H. Roberts (eds.), Grain Legume Crops. Collins, London, UK. pp. 801-846.
- Sundstrom, F. J., Armstrong, J. E., Edwards, R. L. and Mcdowell, B. L. (1986). Relationship between laboratory indices of hot pepper seed vigour and crop greenhouse performance. *Seed Sci. & Technol.* 14: 705-714.
- Sureja A. K. and Sharma R. R. (2000). Genetic variability and heritability studies in garden pea (*Pisum sativum* L.). *Ind. Journ. of Horti.* 57, 3: 243-247.

- Tao, K. J. (1978). Factors causing variation in the conductivity test for soybean seeds. *Journ. of Seed Tech.* 3: 10-18.
- Taweekul, N., Hill G. D., McKenzie A. B. and Hill M. J. (1998). Field performance of field pea seeds with varying vigor levels. *Proceedings Annual Conference Agronomy Society of New Zealand.* 28: 99-105.
- Taylor, A. G., Lee, S. S., Beresniewicz, M. M. and Paine, D. H. (1995). Amino acid leakage from aged vegetable Seeds. *Seed Sci. & Technol.* 23: 113-122.
- Teckrony, D. M. and Egli, D. B. (1977). Relationship between laboratory indices of soybean seed vigor and field emergence. *Crop Science* 17: 573-577.
- TeKrony, D. M., Egli, D. B. and Ohillips, A. D. (1980). Effect of field weathering on the viability and vigour of soybean seed. *Journ. of Agronomy.* 72: 749-753.
- Tomer, R. P. S. and Maguire, J. D. (1990). Seed vigour studies in wheat. *Seed Sci. & Technol.* 18: 383-392.
- Toole, E. H. and Toole, V. K. (1960). Viability of stord snap bean seed as affected by threshing and processing injury. Technical Bulletin. US Deptt. of Agriculture. pp.630-639
- Torfason, W. E. and Nonnecke I. L. (1959). A study of the effects of temperature and other factors upon the germination of vegetable crops. II. Peas. *Canadian. Journ. pl. Sci.* 39: 119-124.
- Torres, R. M., Vieira R. D. and Panobianc M. (2004). Accelerated aging and seedling field emergence in soybean. *Sci. Agric. (Piracicaba, Braz.).* 61, 5: 476-480.
- Townsend, C. E. (1974). Productivity of several perennial forage legumes under irrigation and frequent cutting. Bulletin of Colorado State University Experiment Station no. 562-S. pp.220
- Trachsel, S., Messmer, R., Stamp, P., Ruta, N. and Hund, A. (2010). QTLs for early vigor of tropical maize. *Mol. Breed.* 25: 91-103.
- Tyagi, M. K. and Srivastava, C. P. (2001). Analysis of gene effects in pea. *Legume Res.* 24, 2: 71-76.
- Upadhyaya, H. D., Dwivedi, S. L., Ambrose, M., Ellis, N., Berger, J., Smýkal, P., Debouck, D., Duc, G., Dumet, D. and Flavell, A. (2011). Legume genetic

- resources: Management, diversity assessment, and utilization in crop improvement. *Euphytica*. **180**, 27-47.
- Urbano, G., Aranda, P. and Gomez-Villalva, E. (2003). Nutritional evaluation of pea (*Pisum sativum* L.) protein diets after mild hydrothermal treatment and with and without added phytase. *Journ. of Agri. Food Chem.* 51: 2415-2420.
- Usmani, M. G. and Dubey, R. K. (2007). Association and path coefficient analysis in field pea (*Pisum sativum* var. arvence). *Adv. in Pl. Sci.* 20, **2**: 603-607.
- Vanzolini, S., Araki, C. A. dos. S., Silva, A. C. T. M. da., and Nakagawa, J. (2007). Seedling length test in the evaluation of the physiological quality of soybean seeds. [Portuguese]. *Revista Brasileira de Sementes*. 29, **2**: 90-96.
- Venter van de, H. A., Barla-Szabo, G. and Ybema, S. G. (1993). A study of single and multiple stress seed vigour tests for undeteriorated seed lots of wheat. *Seed Sci. & Technol.* 21: 117-125.
- Vieira, B. G. T. L., Vieira, R. D., Krzyzanowski, F. C. and Franca Neto, J. de B. (2010). Alternative procedure for the cold test for soybean seeds. *Scientia Agricola*. 67, **5**: 540-545.
- Vieira, R. D., Neto, A. S., Mudrovitsch de Bittencourt, S. R. and Panobianco, M. (2004). Electrical conductivity of the seed soaking solution and soybean seedling emergence. *Sci Agric. (Piracicaba, Braz.)* 61, **2**: 164-168.
- Vieira, R. D., Tekorony, B. M. and Egli, D. B. (1992). Effect of drought and defoliation stress in the field on soybean seed germination and vigour. *Crop Sci.* 32: 471-475.
- Vikas and Singh S. P. (1999). Line x tester analysis in pea (*Pisum sativum* L.). *Annals of Agri. Bio. Research*. 4, **1**: 93-98.
- Vinod, S., Singh, R. J., Singh, P. K., Ratnakar, P. and Shahi, B. P. (2011). Additive, dominance and epistatic variation for seed yield and its components in pea. *Environ. and Ecol.* 29: **4** (A): 2113-2115.
- Waldron, B. L., Robins, J. G., Jenson, K. B., Palazzo, A. J., Cary, T. J. and Berdahl, J. D. (2006). Population and environmental effects on seed germination, and seedling vigor in western wheatgrass (*Pascopyrum smithii*). *Crop Sci.* 46, **2**: 503-508.

- Wang, J., Gan, Y. T., Clarke, F. and McDonald, C.L. (2006). Response of chickpea yield to high temperature stress during reproductive development. *Crop Sci.* 46, 2: 171-178.
- Wang, Y. R., Yu, L. L., You, L. and Sheng, Y. (2003). Vigour tests used to assess quality of seed lots and field emergence for several forage species. *Acta Prataculturae Sinica.* 12, 5: 62-69.
- Ware, G. W and McCallum, J. P. (1975). Producing vegetable crops. Thomson publication, Fresno California, pp 61-67.
- West, S. H. and Harris, H. C. (1963). Seed coat colors associated with physiological changes in alfalfa and crimson and white clovers. *Crop Sci.* 3: 190-193.
- Whalley, R. D. B., Mckell, C. M. and Green, L. R. (1966). Pea variability. *Crop Science* 6: 47-152.
- Whitehead, Nobel A and Russell B. (1962). In Principa mathematics. Cambridge University Press (1913) ed. Cambride University Press. pp : 100-105.
- Woltz, J., TeKrony, D. M., and Egli, D. B. (2006). Corn Seed Germination and vigor following freezing during seed development. *Crop Sci.* 46, 1: 526-535.
- Woodstock, L. W. (1969). Biochemical tests for seed vigor. *Proceedings of the International Seed Testing Association.* 34: 253-263.
- Woodstock, L. W. (1973). Physiological and biochemical tests for seed vigour. *Seed Sci. & Tech.* 1: 127-157.
- Woodstock, L. W. and Pollock, B. M. (1965). Physiological predetermination: imbibition, respiration and growth of lima bean seeds. *Science.* 150: 1031-1032. Quoted in AOSA. 1983. Seed Vigor Testing Handbook. Contribution No. 32. Association of Official Seed Analysts. Lincon, NE., USA.
- Wright T, S. (1921). Correlation and association in pea. *Journal of Agri. Research.* 20: 557-587.
- Yaklich, R.W. and Kulik, M. M. (1979) Evaluation of vigor tests in soybean seeds. Relationship of the standard germination test, seedling classification, seedling length and tetrazolium staining to field performance. *Crop Sci.* 19: 247-252

- You, X. F. (2005). Correlation between seed vigour and germination rate of melon. (Chinese). *Fujian Agri. Sci. and Technol.* 1: 20-21.
- Zanakis, G. N., Ellis, R. H. and Summerfield, R. J. (1994). A comparison of changes in vigour among three genotypes of soybean (*Glycine max L.*) during seed development and maturation in three temperature regimes. *Exp. Agr.* 30: 157-170.
- Zhoe, Z., Robards, K., Helliwell, S. and Blanchard, C. (2002). Ageing of stored rice: Change in chemical and physical attributes. *Journ. of Cereal Sci.* 35: 65-68.
- Zohary, D. and Hopf, M. (2000). *Domestication of Plants in the Old World*. Oxford University Press: Oxford, UK, 2000, pp. 57-59.