

List of Figures

Figure Title	Page no.
Fig.3.1 Location of Map of Study Area	44
Fig.3.2 Economic Status of Population around Lakwa area	47
Fig.3.3 Effluent Production from different GGSs of Lakwa Oilfield areas	49
Fig.4.1: Soil components by proportions	71
Fig.4.2: Soil Texture of the study area	77
Fig.4.3: Soil pH at different sampling sites during Monsoon and Non-Monsoon	81
Fig.4.4 : Organic Matter content in soil samples at different sites during monsoon and Non-monsoon	85
Fig.4.5: Carbonate content in soil samples at different sites during Monsoon and Nonmonsoon seasons	88
Fig.4.6(a) and (b): Variation of metal contents in soil with distance Around GGS-I Non-monsoon and Monsoon	97
Fig.4.7(a) and (b): Variation of metal contents in soil samples with distance around GGS-III non-monsoon and monsoon	98
Fig.4.8(a): Variation of metal contents in soil samples with depth around GGS-I during Non-monsoon	101
Fig.4.8(b) Variation of metal contents in soil samples with with depth around GGS-I during monsoon	102
Fig.4.9(a) Variation of metal contents in soil samples	102

with depth around GGS-III during Non-monsoon	
Fig.4.9 (b) : Variation of metal contents in soil samples	103
with depth around GGS-III during Monsoon	103
Fig.4.10(a): Total metal contents in different size fraction	108
of soil samples around GGS-I Non-monsoon	109
Fig.4.10(b): Total metal contents in different size fraction	109
of soil samples around GGS-I Monsoon	109
Fig.4.11 (a) Total metal contents in different size fraction	110
of soil samples around GGS-III, Non-monsoon	
Fig.4.11(b) Total metal contents in different size fraction of	111
soil samples GGS-III Monsoon	
Fig.5.1: Metal contents in different fractions of the tea plants	123
around GGS-I	
Fig.6.1(a)& (b):Root and Shoot length (cm) of <i>Vetiveria</i>	133
<i>zizanioides</i> of different age groups grown under	
different concentration of heavy metals	
Fig.6.2(a)& (b): Root and Shoot length (cm) of <i>Vetiveria</i>	137
<i>zizanioides</i> of different age groups grown under	
different concentration of heavy metals	
Fig.6.3: Accumulation of heavy metal on root/shoot of vetiver plant	143-144
Fig.6.4: Variation of Root essential oil from <i>Vetiveria zizanioides</i>	154
grown on soil soil treated with different	
doses of heavymetal after 120days	

Fig.6.5: Representative FTIR Spectra of Valerenol compound	156
Fig.6.5: Representative NMR Spectra of Valerenol	156
Fig.6.7 Representative GC-MS Spectra of Valerenol	157
Fig.6.8: Chemical structure of major compound Valerenol in Vetiver oil	157