

LIST OF FIGURES

- Fig. 1.1. Distribution of Lower Gondwana basins in the Peninsular India and location of Talchir basin.
- Fig. 1.2. Geological map of the Talchir basin showing the study area.
- Fig. 1.3. Geological map of the Talchir basin showing the study area and the recent classification of Damuda Group.
- Fig. 2.1. Part of survey of India Toposheet Nos. 73 H/1 and D/13 showing the physiography and drainage of the study area.
- Fig. 2.2. Geological map of the Gondwana succession north west of Angul (study area), Talchir Basin, Orissa.
- Fig. 2.3. Composite lithostratigraphic section prepared from the surface and subsurface investigation of the present area.
- Fig. 3.1. Location map of observed stratigraphic sections, observed paleocurrent data, boreholes and samples collected for petrographic analysis.
- Fig. 4.1. Diamictite (Facies A₁) of the basal part of Unit-A showing angular to subrounded pebbles and boulders.
- Fig. 4.2. Stratified diamictite (Facies A₁) of the basal part of Unit-A showing crude stratification.
- Fig. 4.3. Thinly interbedded sandstone and mudstone (Facies A₂) of Unit-A with dropstones.
- Fig. 4.4. Thick mudstone (Facies A₃) occurring in the lower part of Unit-A and C and in the middle part of Unit-B of Talchir Group.
- Fig. 4.5. Thin marlstone beds breaking into a series of concretions linearly arranged along the bedding planes within the thick mudstones (Facies A₃).
- Fig. 4.6. Interbedded sandstone and mudstone (Facies A₄) of Unit-A.
- Fig. 4.7. The conglomerate and sandstone intercalation (Facies A₅) of Unit-A.

- Fig.4.8. Low-angle cross-stratified sandstone (Facies A₆) of the upper part of Unit-A.
- Fig.4.9. Trough cross-stratified sandstone (Facies A₇) of the upper part of Unit-A. The sandstones are vertically stacked and form multistory sandstone bodies.
- Fig.4.10. Massive conglomerate (Facies A₈) occurring intimately associated with Facies A₆ and A₇ in the upper part of Unit-A.
- Fig.4.11. Interbedded sheet sandstone and mudstone (Facies A₉) of the lower part of Unit-B. The Facies is pierced by a lenticular amalgamated channel sandstone (Facies A₁₀).
- Fig.4.12. Interbedded sheet sandstone and mudstone (Facies A₉) of the upper part of Unit-B succeeded by amalgamated channel sandstone (Facies A₁₀) at the top.
- Fig.4.13. Thick amalgamated channel sandstone (Facies A₁₀) of Unit-B showing amalgamated nature, grading, concentration of pebbles and crudely developed fining upward cycles.
- Fig.4.14. Coarsening and thickening upward sandstone and mudstone sequence (Facies A₁₁) of the middle part of Unit-B near Bedasar village.
- Fig.4.15. Stromatolitic algal limestone (Facies A₁₂) of the middle part of Unit-B.
- Fig.4.16. Ripple-drift cross-laminated fine grained sandstone (Facies A₁₃) of the upper part of Unit-C of Talchir Group showing climbing ripple of the type B & C.
- Fig.4.17. Generalised lithostratigraphic section and depositional environments of Unit-A, Unit-B and Unit-C of Talchir Group of the study area.
- Fig.4.18. Interbedded carbonaceous shale - shaly coal and coal (Facies B₁) of Barakar Formation exposed along a stream section.
- Fig.4.19. Partings of sandstone with the carbonaceous shale – shaly coal and coal (Facies B₁) around which the coal splits into thin seams.

- Fig.4.20. Massive grey shale (Facies B₂) showing yellow bands and faint parallel laminations.
- Fig.4.21. Interbedded fine grained sandstone and shale (Facies B₃). The sandstone shows faint parallel laminations.
- Fig.4.22. Ripple-drift cross-laminated fine grained sandstone (Facies B₄) showing A & B type ripple-drift.
- Fig.4.23. Solitary set of large scale planar cross stratification (Facies B₅) in the medium and coarse grained sandstones of the Karharbari and Barakar Formation.
- Fig.4.24. Cosets of large scale planar cross stratification (Facies B₅) in the medium and coarse grained sandstones of the Karharbari and Barakar Formation of the present area.
- Fig.4.25. Cosets of large scale trough cross stratifications (Facies B₆) in the medium to coarse grained sandstones of the Karharbari and Barakar Formation of the present area.
- Fig.4.26. Solitary set of large scale trough cross stratification (Facies B₆) overlying a flat stratified sandstone (Facies B₇) with a scoured base.
- Fig.4.27. Flat stratified sandstone (Facies B₇) with stratification thickness ranging from 5 – 10 cm, overlain by a conglomerate bed.
- Fig.4.28. Laterally coalescing very large scale cosets of trough cross stratification (Facies B₈) in the pebbly sandstones of the Karharbari Formation of the present area.
- Fig.4.29. Lenticular conglomerate (Facies B₉) occurring at the scoured basal surface of a flat stratified sandstone.
- Fig.4.30. Boulder conglomerate (gravel) bed (Facies B₁₀) occurring at the top of the Karharbari Formation southeast of Nisha village.
- Fig.4.31. Matrix supported boulder conglomerate closely associated with clast supported boulder conglomerate or gravel (Facies B₁₀).

- Fig.4.32. Lenticular bed of trough cross-stratified pebbly sandstone (Facies B₈) occurring interbedded with clast supported conglomerate (Facies B₁₀) southeast of Nisha village.
- Fig.4.33. Generalised lithostratigraphic section and depositional environments of the Karharbari and Barakar Formation of Damuda Group of the study area.
- Fig.5.1. Sector level and regional paleocurrent pattern of the Gondwana succession of the present area.
- Fig.6.1. Photomicrograph of a medium to coarse grained sandstone of Unit-A of Talchir group showing quartz, feldspar garnet and opaque floating in an argillaceous matrix. PL, X 32.
- Fig.6.2. Same view as Fig. 6.1, Nicols crossed, X 32, showing monocrystalline nonundulatory and undulatory quartz, polycrystalline quartz, microcline and plagioclase.
- Fig.6.3. Photomicrograph of fine to medium grained sheet sandstone of Unit-B of Talchir Group showing quartz, feldspar, garnet and opaque floating in an argillaceous matrix, PL, X 32.
- Fig.6.4. Same view as Fig.6.3, Nicols crossed, X 32, showing nonundulatory and undulatory quartz, plagioclase feldspar with well developed twinning.
- Fig.6.5. Photomicrograph of the channel sandstone of Unit-B of Talchir Group showing quartz, feldspar, rock fragment, garnet and opaque floating in a calcareous cement. PL, X 32.
- Fig.6.6. Same view as Fig.6.5, Nicols crossed, X 32, showing monocrystalline nonundulatory and undulatory quartz, feldspar, polycrystalline quartz and rock fragment floating in an amorphous calcareous cement.
- Fig.6.7. Photomicrograph of the fine grained sandstone of unit-C of Talchir Group showing very fine quartz, muscovite and biotite flakes, opaque minerals set in a calcareous cement.

- Fig.6.8.** Same view as Fig.6.7, Nicols crossed, X 32, showing monocrystalline nonundulatory quartz, biotite flakes and opaque grains set in an amorphous calcareous cement.
- Fig.6.9.** Photomicrograph of the Karharbari sandstone of Damuda Group showing clear grains of quartz and feldspar cemented by ferruginous cement and argillaceous matrix. PL, X 32.
- Fig.6.10.** Same view as Fig.6.9, Nicols crossed, X 32, showing monocrystalline nonundulatory and undulatory quartz, polycrystalline quartz with numerous grains, microcline with cross-hatched twinning cemented by ferruginous cement and argillaceous matrix.
- Fig.6.11.** Photomicrograph of the medium grained sandstone of the Barakar Formation showing quartz and feldspar floating in an argillaceous matrix and ferruginous cement. PL, X 32.
- Fig.6.12.** Same view as Fig.6.11, Nicols crossed, X 32, showing monocrystalline nonundulatory and undulatory quartz, microcline with cross-hatched twinning cemented by argillaceous matrix and ferruginous cement.
- Fig.6.13.** Photomicrograph of the coarse grained sandstone of Barakar Formation showing angular to subrounded grains of quartz and feldspar floating in an argillaceous matrix.
- Fig.6.14.** Same view as Fig.6.13, Nicols crossed, X 32, showing large no. of polycrystalline quartz with numerous grains, polycrystalline quartz with 3 grains and plagioclase feldspar with well developed twinning cemented by argillaceous matrix and ferruginous cement.
- Fig.6.15.** X-ray diffraction patterns of the sandstones of the present area.
- Fig.6.16.** Classification of the sandstones of the present area.
- Fig.6.17.** Photomicrograph of the heavy minerals.
- A.** Rounded and elongated grains of ilmenite, magnetite, haematite and pyrite (opaques). PL, X 50.
 - B.** Ilmenite altering to leucoxene. PL, X 50.

- C. Garnets showing irregular outline and fractures. PL, X 50.
- D. Elongated zircon with inclusions. PL, X 50.
- E. Euhedral grain of pink tourmaline with inclusions. PL, X 100.
- F. Euhedral grain of brown tourmaline. PL, X 100.
- G. Subhedral grain of indicolite (Blue tourmaline). PL, X 100.
- H. Elongated grain of rutile. PL, X 100.
- I. Irregular grain of hypersthene with 1 set of cleavage. PL, X 100.
- J. Subhedral grain of blue-green hornblende with 1 set of cleavage. PL, X 100.
- K. Basal section of biotite flake. PL, X 100.
- L. Rounded grain of apatite with inclusions. PL, X 100.
- M. Euhedral grain of sphene. PL, X 100.
- N. Euhedral grain of anatase with dark border. PL, X 100.
- O. Elongated grain of sillimanite. CN, X 100.
- P. Rounded grain of monazite. PL, X 100.

Fig.6.18. X-ray diffraction patterns of the heavy minerals of the various formations of the Talchir Group of the present area.

Fig.6.19. X-ray diffraction patterns of the heavy minerals of the various formations of the Damuda Group of the present area.

Fig.6.20. Frequency distribution of heavy minerals in the various formations of the Talchir Group.

Fig.6.21. Frequency distribution of heavy minerals in the various formations of the Damuda Group.

Fig.7.1. Regional paleoflow pattern of the Gondwana succession of the present area.

Fig.7.2. Provenance discrimination diagram after Basu et al., 1975.

Fig.7.3. QFL and QmFLt plots of framework modes of terrigenous sandstones of the Gondwana sequence of the present area showing different provenance types after Dickinson et al., 1983.

- Fig.7.4.** Frequency distribution of heavy minerals in the various formations of the Talchir and Damuda Group of the present area.
- Fig.7.5.** Generalised lithostratigraphic section and the depositional environments of the Gondwana sequence of the present area.
- Fig.8.1.** Generalised lithostratigraphic section, its classification, depositional environments, paleoflow pattern and the provenance characteristics of the Gondwana sequence of the present area.