

LIST OF FIGURES

		Page
1.	(i) Number of tillers, dry weight per plant nitrogen per cent and nitrogen uptake in different varieties (<u>Kharif</u> , 1966)	57
	(ii) Grain yield and ear bearing tillers as influenced by deficiency (major nutrients) .	
2.	Tillers per m ² at various growth stages under dry-sown and transplanted conditions (<u>Rabi</u> and <u>Kharif</u> , 1968).	60
3.	Dry weight (g/plant) at various growth stages under dry-sown and transplanted conditions (<u>Rabi</u> and <u>Kharif</u> , 1968).	60
4.	Nitrogen per cent at various growth stages under dry-sown and transplanted conditions (<u>Rabi</u> and <u>Kharif</u> , 1968).	61
5.	Nitrogen uptake (mg/plant) at various growth stages under dry-sown and transplanted conditions (<u>Rabi</u> and <u>Kharif</u> , 1968).	61
6.	Yield, yield components, ear bearing tillers per m ² , grain-straw ratio and per cent tiller mortality under dry-sown and transplanted conditions (<u>Rabi</u> , 1968).	62
7.	Tillers per plant and nitrogen uptake (mg/plant) in different varieties at various growth stages under dry-sown conditions (<u>Kharif</u> , 1969).	63
8.	Tillers per plant and nitrogen per cent at various growth stages under 2 split and 3 split nitrogen application in different varieties (<u>Rabi</u> , 1970).	64
9.	(1) Regression lines showing correlation between grain number and spikelet number per panicle and nitrogen uptake at booting and maximum tillering (<u>Kharif</u> , 1972).	64

	Page
9. (ii) Grain number, spikelet number and nitrogen uptake at booting and maximum tillering in tillers of different ages (<u>Kharif</u> , 1972).	64
10. Per cent recovery of labelled carbon (C^{14}) in grain when fed at different stages of grain maturity (<u>Kharif</u> , 1969).	69
