

LIST OF FIGURES

<u>FIG. No.</u>	<u>TITLE</u>	<u>PAGE</u>
1.	Plot of copper(I) formed as a function of time at 25°C (0.16 mol dm ⁻³ copper sulphate + 1 g copper crystals 2-30 μm).	.. 58
2.	Plot of copper(I) formed as a function of time at 25°C (0.16 mol dm ⁻³ copper sulphate + 4 g copper crystals 2-30 μm).	.. 58
3.	Plot of copper(I) formed as a function of time at 25°C (0.16 mol dm ⁻³ copper sulphate + 8 g copper crystals 2-30 μm).	.. 58
4.	Plot of copper(I) formed as a function of time at 25°C for different copper sulphate concentrations (8 g copper crystals 2-30 μm in 20% AN).	.. 58
5.	Plot of copper(I) formed as a function of time at different temperatures (0.32 mol dm ⁻³ copper sulphate + 8 g copper crystals 2-30 μm in 20% AN).	.. 59
6.	Plot of copper(I) formed as a function of time at 25°C with different stirring speeds (0.32 mol dm ⁻³ copper sulphate + 8 g copper crystals 2-30 μm in 20% AN)	.. 59

7. Fraction of Cu^{2+} remaining (α) as a function of time parameter ($\log t$) at 25°C (0.32 mol dm^{-3} copper sulphate + 1 g copper crystals $2-30 \mu\text{m}$). 62
8. Fraction of Cu^{2+} remaining (α) as a function of time parameter ($\log t$) at 25°C (0.32 mol dm^{-3} copper sulphate + 4 g copper crystals $2-30 \mu\text{m}$). 62
9. Fraction of Cu^{2+} remaining (α) as a function of time parameter ($\log t$) at 25°C (0.32 mol dm^{-3} copper sulphate + 8 g copper crystals $2-30 \mu\text{m}$). 62
10. Fraction of Cu^{2+} remaining (α) as a function of time parameter ($\log t$) at 35°C (0.32 mol dm^{-3} copper sulphate + 4 g copper crystals $2-30 \mu\text{m}$). 69
11. Fraction of Cu^{2+} remaining (α) as a function of time parameter ($\log t$) at 55°C (0.32 mol dm^{-3} copper sulphate + 4 g copper crystals $2-30 \mu\text{m}$). 69
12. Fraction of Cu^{2+} remaining (α) as a function of time parameter ($\log t$) at 25°C for different values of surface area (0.32 mol dm^{-3} copper sulphate in 20% AN). .. 72
13. Fraction of Cu^{2+} remaining (α) as a function of time parameter ($\log t$) at 25°C with different stirring speeds (0.32 mol dm^{-3} copper sulphate + 8 g copper crystals $2-30 \mu\text{m}$ in 20% AN). .. 75

14. Plot of change of copper(I) concentration $[\text{Cu}^+]$ as a function of time from 0.40 mol dm^{-3} Cu_2SO_4 solution. .. 80
15. Plot of the rate (R) of reaction (16) as a function of initial Cu_2SO_4 concentration C_0 . .. 81
16. Fraction of Cu^+ remaining (α) as a function of time parameter ($\log t$) for different initial Cu_2SO_4 concentrations (C_0) in 10% AN. .. 83