

# CONTENTS

	PAGE No.
<b>GENERAL INTRODUCTION</b>	<b>1.1-1.23</b>
1.1 Aetiology.	1.3
1.2 Early History Of The Disease.	1.3
1.3 Prevalence.	1.4
1.4 Life Cycle of <i>Wuchereria Bancrofti</i> Parasite..	1.6
1.5 Different Forms of Filariasis.	1.8
1.5.1 Lymphatic Filariasis.	1.9
1.5.2 <i>Brugian</i> Filariasis.	1.9
1.5.3 <i>Bancroftian</i> Filariasis.	1.10
1.5.4 Occult Filariasis.	1.10
1.6 Epidemiology.	1.10
1.7 The Disease Spectrum.	1.11
1.7.1 Asymptomatic Amicrofilaraemic State (Endemic Normals).	1.12
1.7.2 Asymptomatic Microfilaremic (ASM) State.	1.12
1.7.3 The Acute Manifestations.	1.12
1.7.4 The Chronic Manifestations.	1.14
1.7.5 Tropical Pulmonary Eosinophilia (TPE).	1.14
1.8 Pathology.	1.14
1.8.1 Immunopathology.	1.17
1.9 Protective Immune Response in Lymphatic Filariasis.	1.19
<b>REVIEW OF LITERATURE</b>	<b>2.1 -2.20</b>
2.1 Immunological Profile.	2.1
2.2 Helminthic Infection & The Immune System with Special Reference to <i>W.Bancrofti</i> Infection.	2.4
2.3 Cellular and Humoral Immunity.	2.6
2.3.1 Cellular Responses.	2.7
2.3.2 Humoral Responses.	2.1
<b>BACKGROUND TO THE WORK</b>	<b>3.1-3.6</b>

<b>MATERIALS AND METHODS</b>	<b>4.1-4.28</b>
4.1 Materials Used in the Present Study and Their Source.	4.1
4.2 Study Population.	4.1
4.3 Staining of Blood Smear.	4.3
4.4 Preparation of Serum.	4.4
4.5 Detection of <i>Wuchereria bancrofti</i> parasite Antigens circulating in the blood by direct-sandwich ELISA.	4.4
4.6 Isolation of <i>W.bancrofti</i> Microfilariae from blood of ASM individuals.	4.6
4.7 Preparation of total antigens from <i>Wuchereria bancrofti</i> Microfilaria.	4.7
4.8 Preparation of <i>Wuchereria bancrofti</i> microfilaria - excretory secretory Antigens.	4.8
4.9 Preparation of <i>Setaria digitata</i> Adult soluble Antigen.	4.8
4.10 Sodium dodecyl Sulphate Polyacrylamide gel electrophoretic separation (SDS-PAGE) of Proteins:	4.9
4.11 Silver Staining of Proteins separated by SDS-PAGE.	4.11
4.12 Detection of parasite antigen specific IgG isotypes in filaria sera using Enzyme linked immunosorbent assay (ELISA)].	4.12
4.13 Immunochemical visualisation of antigens (Immunoblot) after electrophoretic transfer of proteins from SDS-Polyacrylamide gels to nitro-cellulose membranes.	4.14
4.14 Preparation of <i>Setaria digitata</i> adult soluble fractionated antigens bound to Nitro-cellulose Paper.	4.16
4.15 Isolation of Peripheral Blood Mononuclear Cells (PBMC) from human blood.	4.17
4.16 <i>In vitro</i> stimulation of PBMC for cell proliferation and cytokine secretion.	4.18
4.17 Immunoassay of Human IFN- $\gamma$	4.19
4.18 Immunoassay of Human IL-2 .	4.21
4.19 Immunoassay of Human IL-4.	4.22
4.20 Immunoassay of Human IL-5.	4.23
4.21 Immunoassay of Human IL-10.	4.25
4.22 Treatment of Microfilaria Carriers with Diethylcarbamazine Citrate (DEC) and Measurement of Parasite specific IgG4 titres in their blood:	4.27

4.23 Induction of Th1/Th2 Cytokines in the Lymphocytes of Microfilariae Carriers from different families treated with DEC and those who were not treated with DEC using 14-20 kDa Antigens of <i>S. digitata</i> adult.	4.27
4.24 Statistical Analysis.	4.28
<b>RESULTS</b>	5.1-5.55
5.1 Selection of Study Area and Study Population.	5.1
5.2 Antigens Used in the Present Study.	5.4
5.3 Parasite Antigen-Specific IgG Isotype Responses by Antigen Negative EN and ASM Individuals.	5.4
5.4 <i>Setaria digitata</i> Adult Soluble Antigens Recognised by Different IgG Isotypes Circulating in the Sera of EN and ASM Individuals as seen by Immunoblot.	5.7
5.5 Optimal Antigen Concentration Required for Stimulation of Peripheral Blood Lymphocytes <i>in-vitro</i> .	5.7
5.6 Kinetics of Induction of Maximal Cytokine Secretion in Peripheral Blood Lymphocytes by Optimal Conc. of Different Antigens.	5.22
5.7 The Effect of Freezing of Peripheral Blood Lymphocytes in Liquid Nitrogen on their Ability to Secrete Cytokine.	5.22
5.8 The Effect of Storing of Peripheral Blood Lymphocytes at Room Temperature on their Ability to Secrete Cytokine.	5.29
5.9 The Ability of <i>S. digitata</i> Adult Antigen Fractions and Other Antigens to Induce Secretion Of IL-2 And Proliferation of Peripheral Blood Lymphocytes Obtained From EN And ASM Individuals.	5.29
5.10 Differential IFN- $\gamma$ And IL-4 Secretion Induced by <i>S. digitata</i> Adult Antigen Fractions in Peripheral Blood Lymphocytes of EN And ASM Individuals.	5.33
5.11 Induction of Differential Th1/Th2 Cytokine Responses in the Lymphocytes of EN and ASM Individuals by <i>S. digitata</i> 14-20 kDa Antigens (Fr-6) and <i>W. bancrofti</i> Microfilaria-ES Antigens.	5.35
5.12 Clearance of Microfilariae from Circulation after DEC Therapy and it's Effect on Antigen Specific IgG4 titres.	5.37

5.13 Induction of Th1/Th2 Cytokines in the Lymphocytes of Individuals who cleared Microfilariae circulating in the Blood after DEC therapy using 14-20 kDa <i>Setaria digitata</i> Adult Antigens.	5.55
<b>DISCUSSION</b>	6.1-6.15
<b>CONCLUSION</b>	7.1-7.2
<b>BIBLIOGRAPHY</b>	8.1-8.31