

## LIST OF PLATES

- Plate 1 : Glandular Thelypteroid ferns of the present study.
- Plate 2 : *Candida* species tested for MIC.
- Plate 3 : Screening of antifungal activity in epidermal gland extracts of three ferns with leaf discs.
- Plate 4 : Screening of antifungal activity in epidermal gland extracts of three ferns with Potato discs.
- Plate 5 : Screening of antifungal activity in epidermal glands of three ferns with seeds.
- Plate 6 : Anti candidal activity in epidermal gland extracts of *Cyclosorus interruptus* and *Christella parasitica*.
- Plate 7 : Antibacterial screening of different parts of *Cyclosorus interruptus*.
- Plate 8 : Antibacterial activity in Crude extract (Acetone) and in Hexane extract obtained through sequential fractionation of the crude extract.
- Plate 9 : Antibacterial activity in Chloroform extract and in Ethyl acetate extract obtained through sequential fractionation of the crude extract.
- Plate 10 : Anticandidal activity in Crude extract (Acetone) and in Hexane extract obtained through sequential fractionation of the crude extract.

- Plate10A : Anticandidal activity in Chloroform extract and in Ethyl acetate extract obtained through sequential fractionation of the Crude extract.
- Plate 11 : Determination of MIC of Ethylacetate extract obtained through sequential fractionation of epidermal glands of *C. interruptus* for five different fungi by broth microdilution method.
- Plate 12 : Determination of MIC of Ethylacetate extract obtained through sequential fractionation of epidermal glands of *C. interruptus* for Three different dermatophytes by broth microdilution method.
- Plate 13 : Determination of MIC of Crude extract and different extracts (Hexane, Chloroform, Ethyl acetate) obtained through sequential fractionation of epidermal glands of *C. interruptus* for four different *Candida* species by broth microdilution method.
- Plate 14. Anticandidal (*Candida albicans*) activity of Ethyl acetate fraction of crude extract (50-300 microgram).
- Plate 15 : Anticandidal activity against clinical samples of *Candida* species in Ethyl acetate extract (obtained through sequential fractionization of Crude extract with Hexane, Chloroform, Ethyl acetate) by disc diffusion method (20-500  $\mu\text{g}$  /disc).
- Plate 16 : Anticandidal activity against samples of *Candida* species in Ethyl acetate extract (obtained through sequential fractionization of Crude extract with Haeane, Chloroform, Ethyl acetate) by disc diffusion method (200-1500  $\mu\text{g}$  /disc).

- Plate 17 : Antibacterial activity in Crude extract and sequential fractions of Hexane, Chloroform and Ethyl acetate.
- Plate 18 : Effect of Ethyl acetate fraction on biofilm of *Candida albicans* in comparison with antibiotic Fluconazole.
- Plate 19 : HPTLC – Instrumentation.
- Plate 20 : HPTLC fingerprint of crude (Acetone) extract.
- Plate 21 : HPTLC fingerprint of ethyl acetate fraction.
- Plate 22 : HPTLC fingerprints of Crude extract and ethyl acetate fraction.
- Plate 23 : HPTLC fingerprint of stearic acid, crude extract and ethyl acetate fraction.
- Plate 24 : HPLC chromatogram Crude (Acetone) extract of epidermal glands of *Cyclosorus interruptus*.
- Plate 25 : Portions of HPLC chromatogram of Crude extract and Ethyl acetate fraction to show the major difference.
- Plate 26 : FTIR Spectrum of Crude extract and Ethyl acetate fraction.
- Plate 27 : Figure to show major change in FTIR spectrum between Crude extract and Ethyl acetate extract between 1000-2000 cm<sup>-1</sup>.
- Plate 28 : Identification of active compounds by comparing TLC, Bioautography, HPLC and HPTLC.
- Plate 29 : Mechanism of action of antifungal drugs.