

CONTENTS

PAGE

Chapter I	Scope and Object of the Research	1
1.1	Choice of the main solvent used	2
1.2	Scope of the work	3
1.3	Methods of investigations	4
1.4	Summary of the work done	5
	References	8
Chapter II	General Introduction	10
2.1	The importance of solution chemistry	10
2.2	Thermodynamics of ion-pair formation	12
2.3	Viscosity	14
2.4	Conductance	19
2.5	Apparent and Partial Molal Volumes	29
2.6	Ultrasonic Velocity	32
2.7	Some recent trends in solvation models	34
2.8	A brief literature survey of the solvents used for investigation	36
	References	45
Chapter III	Experimental Section	55
3.1	Chemicals used: Their sources and purification	55
3.2	Experimental techniques	57
	References	63

Chapter IV	Thermodynamics of ion association and solvation in 2- methoxyethanol: Behaviour of tetraphenylarsonium, picrate, and tetraphenylborate ions from conductivity and ultrasonic data	64
4.1	Experimental section	65
4.2	Results	66
4.3	Discussion	69
	References	78
Chapter V	Thermodynamics of ionic association of tetraphenylphosphonium, tetraphenylarsonium, and some common cations in 2- methoxyethanol using conductivity and FT-Raman spectroscopy	87
5.1	Experimental section	88
5.2	Results	90
5.3	Discussion	92
	References	103
Chapter VI	Ultrasonic velocities and isentropic compressibilities of electrolytes in 2-methoxyethanol in the temperature range 288.15 to 308.15 K	111
6.1	Experimental section	112
6.2	Results and discussion	113
	References	119

Chapter VII	A study on the solvation phenomena of some sodium salts in 1,2-dimethoxyethane from conductance, viscosity, ultrasonic velocity, and FT-Raman spectral measurements at 298.15 K	125
7.1	Experimental section	126
7.2	Results and discussion	127
	References	137
Chapter VIII	Excess properties (excess volume, viscosity deviations and isentropic compressibility changes) of binary mixtures of N-Methylacetamide + 2-Methoxyethanol, and N-Methylacetamide + Water, at 308.15, 313.15 and 318.15 K.	143
8.1	Experimental section	146
8.2	Results	146
8.3	Discussion	148
	References	159
Chapter IX	Excess properties (excess volume, viscosity deviations and isentropic compressibility changes) of binary mixtures of N,N-Dimethylacetamide + 2-Methoxyethanol, and N,N-Dimethylacetamide + Water, at 298.15, 308.15 and 318.15 K.	168
9.1	Experimental section	169
9.2	Results	169
9.3	Discussion	171
	References	179
Chapter X	Concluding Remarks	187