

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE NO.
3.1	Outline of methodology adopted in this study	31
3.2	Map showing the regions (1 to 5) and the corresponding satellite images of sample locations	32
3.3	Pitting for sample collection	33
3.4	Picture depicting the instrument setup	38
3.5	Spectra of pure bentonite obtained from different heights of observation	40
3.6	Spectra of Kaolinite sample with different diameter of the sample spread	42
3.7	Spectra of Kaolinite sample with different sample depth	43
3.8	Line sketch of the experimental setup for this study	44
3.9	Electronic sieve shaker and the instrument setup	46
4.1	(a) Spectra of river sand of different texture	52
	(b) Curvature at 1000-1400nm region	
4.1	(c) Spectra of a river sand sample with different texture in visible region	54
4.1	(d) Result of spectral parameter of a river sand sample with different texture	59
4.2	(a) Spectra of a beach sand sample with different texture	60
4.2	Spectra of a beach sand sample with different texture b) 350-800 nm region c) 900-1400 nm region	61

FIGURE NO.	TITLE	PAGE NO.
4.2	Spectra of a beach sand sample with different texture at d) 1900-2300nm region e) 1850-2000nm region	62
4.2	(f) Result of spectral parameter of a beach sand sample with different texture	66
4.3	a) Spectra of coloured beach sand with varying grain size	67
	b) Result of spectral parameter of coloured beach sand samples with different textures	69
4.4	(a) Spectra of a red soil sample with different texture	69
	(b) Result of spectral parameter of red soil samples with different textures	72
4.5	(a) Spectra of a black soil sample with different texture	72
	(b) Correlation of spectral parameter of black soil samples with different textures	74
4.6	Spectra of black soil with different water content	78
4.7	Spectra of brown soil with different water content	90
4.8	(a) A brown soil spectra with different water content.	
	(b) Depth variation of the absorption trough in the 1400nm region	91
4.9	Spectra of a black soil with different water content	93
4.10	a) Spectra of a black soil with different water content.	94
	(b) Depth variation at 1400nm region with varying	94

FIGURE NO.	TITLE	PAGE NO.
	water content. (c) Depth variation at 1900nm region with varying water content	94
4.11	(a) Spectra of a red soil with different water content (b) Depth and width variation at 1400nm region with varying water content. (c) Depth variation at 1900nm region with varying water content	97
4.12	Spectra of a white soil sample with different water content	100
4.13	Spectra of soil samples with different organic content	105
4.14	Area under the curve (AUC) as a function of organic content	106
4.15	Peak reflectance value as a function of organic content	106
4.16	Spectra of soil samples in 650-900nm region with Different organic content	107
4.17	Slope at visible region as a function of organic content	108
4.18	Area under the curve (AUC) at visible region as a function of organic content	108
4.19	Spectra of different minerals Source (Price, J. C., 1994)	112
4.20	Spectra of standard clays (a) Bentonite (b) Kaolinite	114
4.21	Spectra of minerals from USGS spectral library (Clark et al.2007) (a) Kaolinite (b)bentonite (c) Quartz	115 116

FIGURE NO.	TITLE	PAGE NO.
4.22	Spectra of silica sand	117
4.23	(a) Images of coloured clay samples used in this study	118
	(b) Spectra of a white colour soil	118
	(c) Spectra of a red soil	119
	(d) Spectra of a black soil	120
4.24	Spectra of a beach and river sand	121
4.25	(a) Spectra of a soil (K)	121
	(b) Spectra of a soil (P3)	122
	(c) Spectra of a soil (K 33 a)	122
	(d) Spectra of a soil (Ney 2b)	123
	(e) Spectra of a soil (N6 a)	123
	(f) Spectra of a soil (K 37 a)	124
	(g) Spectra of a soil (Ney 4 a)	125
	(h) Spectra of a red soil from Guduvanchery (K2)	125
	(i) Spectra of a black soil from lake	126
4.26	(a) VNIR-SWIR spectra of sands used in this study	132
	(b) VNIR-SWIR spectra of clays used in this study	132
4.27	Spectral response of sand with different proportions of kaolin clay	135
4.28	VNIR-SWIR spectra of sand with different proportion of red clay	138
4.29	VNIR-SWIR spectra of sand with different proportion of black clay	139
4.30	VNIR-SWIR spectra of sand with different proportion of ball clay	140
4.31	Relationship between kaolin clay content and four of the spectral parameters	141

FIGURE NO.	TITLE	PAGE NO.
4.32	Relationship between red clay content and five of the spectral parameters	144
	(a) Slope at 350-550nm region as a function of clay content	
	(b) Depth of absorption at 1400nm as a function of clay content	
	(c) Depth at 1400nm at a function of clay content	
	(d) Depth at 2250nm at a function of clay content	
	(e) Area under the curve as a function of clay content	144
4.33	Relationship between black clay content and four of the spectral parameters	
	(a) Depth at 1400nm as a function of clay content	
	(b) Depth at 1900nm as a function of clay content	
	(c) Radius of curvature in 2250nm as a function of clay content	
	(d) AUC as a function of clay content	146
4.34	Relationship between ball clay content and four of the spectral parameters	
	(a) & (b) Depth at 1400 & 1900nm as a function of clay content	
	(c) Depth at 2250nm at a function of clay content	
	(d): Shift in position at a function of clay content	148
	(e) Area under the as a function of clay content	
	(f) Radius of curvature as a function of clay	149
4.35	(a) Spectra of a red soil with different density	
	(b) Spectra of a black soil with different density	154
	(c) Spectra of a kaolinite soil with different density	155

FIGURE NO.	TITLE	PAGE NO.
4.36	(a) Spectra of guduvanchery sample with the details of liquid limit and plastic limit	157
	(b) Spectra of guduvanchery sample	157
	(c)Depth at 1900nm as a function of white soil liquid limit d) Depth at 1900nm as a function of white soil plastic limit e) Depth at 1900nm as a function of black soil plastic limit	158