

LIST OF TABLES

TABLE No.	TITLE	IN BETWEEN PAGES
Table 2.1	Statewise area under production, total production and productivity of tomato (during 2007-2008).	6-7
Table 4.1	Places of collection of ToLCD infected tomato plants.	53-54
Table 4.2	Primer sequences and PCR conditions used for detection begomovirus associated genomic components.	53-54
Table 4.3	ToLCD samples from which viral genome were cloned, sequenced and characterized.	54-55
Table 4.4	Begomoviruses, their genomic components (DNA-A, DNA-B, and DNA- β) with GenBank Accession no. used for analysis.	54-55
Table 4.5	Detection of ToLCV infection by PCR from infected sample.	54-55
Table 4.6	Restriction fragment pattern of viral clones.	56-57
Table 4.7	Percentage sequence identity among genomic components of begomoviruses causing ToLCD in India.	56-57
Table 4.8	Comparision of Rep binding sequences of distinct ToLCVs occurring in India.	59-60
Table 5.1	Infectivity of infectious constructs and variation of symptoms induced on agroinoculated plants.	68-69
Table 5.2	Percentage nucleotide and amino acid sequence identities between ToLCPaV and other selected begomoviruses.	70-71
Table 5.3	Percentage nucleotide and amino acid sequence identities between ToLCPaB and other selected betasatellites.	71-72
Table 5.4	Breakpoint analysis of ToLCPaV and their putative parental sequences.	72-73
Table 6.1	Details of ToLCGV isolates and respective abbreviation used in this study.	81-82
Table 6.2	ToLCGV constructs used for agroinoculation and symptoms induced on plants.	82-83
Table 6.3	Genome organization and ORF details of three isolates	85-86

of ToLCGV.

Table 6.4	Nucleotide and amino acid percentage identity among DNA-A of ToLCGV isolates.	85-86
Table 7.1	Genome organization and ORF details of DNA-A and DNA- β of ToLCBDV-[Ran].	93-94
Table 7.2	Per cent nucleotide and amino acid sequence identities between ToLCBDV-[Ran] and other selected begomoviruses occurring in Asia.	94-95
Table 7.3	Per cent nucleotide and amino acid sequence identities between ToLCBDB and other selected betasatellites occurring in Asia.	94-95
Table 7.4	Breakpoint analysis of ToLCBDV-[Ran] and their putative parental sequences.	94-95
Table 7.5	ToLCBDV-[Ran] constructs combination(s) used for agroinoculation on plants and symptom study.	96-97
Table 8.1	Details of constructs, respective abbreviation, and probes used in betsatellite transreplication study.	102-103
Table 8.2	Constructs, their infectivity and symptoms induced on agroinoculated plants.	102-103
Table 8.3	Percentage nucleotide identities among full-length of five satellites DNA- β used in betsatellite transreplication study.	105-106
Table 8.4	Percentage nucleotide identities among SCR sequences of five satellites DNA- β used in betsatellite transreplication study.	105-106
Table 8.5	Percentage amino acid identities among β C1 ORF of five satellites DNA- β used in betsatellite transreplication study.	105-106