

REFERENCES

- Abbot, R.T. 1954. The marine molluscs of Grand Cayman Island. Monogr. Acad. Natur. Sci. Philadelphia, No. 11.
- Abidi, S.A.H., Desai, B.N. and Gauton, O.S. 1978. The length weight relationship in Sardinella sirmwal from the Andaman sea. J. Ind. Fish. Ass., 8 : 15-19.
- Abraham, K.C. 1953. Observations on the biology of Meretrix casta (Chemnitz). J. Zool. Soc. India., 5(2) : 163-190.
- Achaval, Z.C. 1984. Isoenzymes of lactate dehydrogenase in fishes of the superorder Acanthopterygii - An update. Comp. Biochem. Physiol., 79B (1) : 1-8.
- Ahmed, M. 1975. Speciation in living oysters. Adv. Mar. Biol., 13 : 357-397.
- Ahmed, M., Skibinski, D.O.F. and Beardmore, J.A. 1977. An estimate of the amount of genetic variation in the common mussel, Mytilus edulis. Biochem Genet., 15 : 833-846.
- Alston R.E. and Turner, B.L. 1963. Biochemical systematics. Prentice-Hall, New Jersey, 404 pp.
- *Annandale, N. 1916. Fauna of Chilka Lake. Mem. Ind. Mus., 5: 350-360.
- *Annandale, N. and Kemp, S. 1916. Fauna of the Chilka Lake, Mollusca, Gastropoda and Lamellibranchia. Mem. Ind. Mus., 5: 329-366.
- Ansari, Z.A., Parulekar, A.H. and Matondkar, S.G.P. 1981. Seasonal changes in meat weight and biochemical composition in the black clam, Villorita cyprinoides (Gray). Indian J. Mar. Sci., 10: 128-131.
- Ansell, A.D. 1974a. Seasonal changes in biochemical composition of the bivalve, Abra alba from the Clyde sea area. Mar. Biol., 25: 13-20.

- Ansell, A.D. 1974b.** Seasonal changes in biochemical composition of the bivalve, Chlamys septemradiata from the Clyde sea area. Mar. Biol., 25: 85-99.
- Ansell, A.D. 1974c.** Seasonal changes in biochemical composition of the bivalve, Nucula sulcata from the Clyde sea area. Mar. Biol., 25: 101-108.
- Ansell, A.D., Frenkeil, L. and Moueza, M. 1980.** Seasonal changes in tissue weight and biochemical composition for the bivalve Donax trunculus L on the Algerian coast. J. Mar. Biol. Ecol., 45 : 105-116.
- Arias, E. and Morales, E. 1977.** Comparative electropherograms of muscle proteins of Mullus surmuletus and M. barbatus. Invest. Perg., 41(2): 323-330.
- Arias, E. and Morales, E. 1980.** Comparative electropherograms of muscle proteins of Salpa salpa and Boops boops. Invest. Perg. (Barc.), 44(1): 35-41.
- Avise, J.C. 1974.** Systematic value of electrophoretic data. Syst. Zool., 23: 465-481.
- *Awati, P.R. and Rai, H.S. 1931.** Ostrea cucullata (The Bombay Oyster). Indian Zool. Memoir., III : 107 pp.
- Ayala, F.J. 1975.** Genetic differentiation during speciation process. In : Evolutionary biology, Vol. 8 Plenum Press, New York., London, pp 1-78.
- Ayala, F.J. 1976.** Molecular evolution (Sundarland : Sinauer Associates).
- Ayala, F.J., Hedgecock, D., Zumwatt, G.S. and Valetine, J.W. 1973.** Genetic variation in Tridacna maxima, an ecological analog of some unsuccessful evolutionary lineages. Evolution., 27: 177-191.
- Balasubrahmanyam, K. and Natarajan, R. 1988.** Seasonal changes in the

biochemical composition of Meretrix casta (Chemnitz) occurring in Vellar estuary. CMFRI. Bull. 42(1): 184-186.

- Balasubrahmanyan, T., Vijayaraghavan, S. and Krishnakumari. L. 1979.**
Energy content of the wedge clam, Donax spiculum. Mar. Biol., 17(4) : 318-332.
- Barnes, H. and Blackstock, J. 1973.** Estimation of lipids in marine animals and tissues : Detailed investigation of the sulpho-phosphovanillin method for total lipids. J. Exp. Mar. Biol. Ecol., 12 : 103-118.
- Barnes, H. and Healy, M.J.R. 1965.** Biometric studies on some common cirripeds. I. Balanus balanoides : measurements of the scuta and terga of animals from a wide geographical range. J. Mar. Biol. Assn. U.K., 45 : 779-789.
- Barnes, H. and Healy, M.J.R. 1969.** Biometrical studies on some common cirripeds. II. Discriminant analysis of measurements on the scuta and terga of Balanus balanus L., B. crenatus Brug., B. improviscus Darwin, B. glandula Darwin and B. amphitrite stutsburi Darwin (B. pallidus stutsburi). J. Exp. Mar. Biol. Ecol., 4 : 51-70.
- Battaglia, B. and Beardmore, J.A. 1978.** Marine organisms : Genetics, ecology and evolution. Plenum Press, New York.
- Baur, E.W. and Schorr, R.T. 1969.** Genetic polymorphism of tetrazolium oxidase in dogs. Science., 166 : 1524-1525.
- Bayne, B.L. 1976.** Marine mussels : their ecology and physiology. Bayne, B.L. (Ed.), Cambridge Univ. Press, 506 pp.
- Bayne, B.L., Buset, A., Gabbot, P.A., Livingstone, D., Lowe, D.M. and Moore, M.N. 1982.** Glycogen utilization and gametogenesis in Mytilus edulis L. Mar. Biol. Lett., 3 : 89-105.
- Beachem, T.D. 1985.** Variation and morphometric variation in pink salmon,

- Onchorynchus gorboscha in southern British columbia and Puget sound.
Can. J. Zool., 63(2) : 366-372.
- Beaumont, A.R. 1982a.** Geographic variation in allele frequencies at three loci in Chlamys opercularis from Norway to the Brittany coast.
J. Mar. Biol. Assn. U.K., 62: 243-261.
- Beaumont, A.R. 1982b.** Variations in heterozygosity at two loci between year classes of a population of Chlamys opercularis (L), from a scotish sea loch. Mar. Biol. Lett., 3: 25-33.
- Beaumont, A.R. and Beveridge, C.M. 1984.** Electrophoretic survey of genetic variation in Pecten maximus, Chlamys opercularis, C. varia and C. distorta from the Irish sea. Mar. Biol., 81 : 299-306.
- Beaumont, A.R. and Gruffydd, L.I.D. 1975.** A polymorphic system in the sarcoplasm of Chlamys opercularis. J. Cons. Nt. Explor. Mer., 36(2): 190-192.
- Beckwitt, R. 1985.** Population genetics of the sand crab Emerita anologa Simpson in S. carolina. J. Exp. Mar. Biol. Ecol., 91(1-2): 45-52.
- Bedford, J.J. and Reid, M.S. 1969.** Gel electrophoresis of proteins in the crystalline style of certain Mollusca. Comp. Biochem. Physiol., 29: 659-664.
- *Benson, 1860.** Ann. and Mag. Nat. Hist., 3rd Ser., VI, p. 260. Quoted from "Notes on Lamellibranchs in the Indian museum", (Prashad, B. 1921). Rec. Ind. Mus., XXII : 111-119, Pl. XV.
- Berger, E. 1973.** Gene-enzyme variation in three sympatric species of Littorina. Biol. Bull. (Woods Hole, Mass.), 145: 83-90.
- Bolaffi, J.L. and Booke, H.E. 1974.** Temperature effects on lactate dehydrogenase isoenzyme distribution in skeletal muscle of Fundulus heteroclitus (Pisces : Cyprinodontiformes). Comp. Biochem. Physiol., 48B:

557-564.

- Borowsky, R., Borowsky, B., Milani, H. and Greenberg, P. 1985. Amylase variation in the salt marsh amphipod, Gammarus palustris. Genetics., 111: 311-323.
- Boss, R.J. 1982. Mollusca. In: Synopsis and organisation of Living organisms, Parker, S.Q.(Ed), Mc Graw-Hill, New York, pp. 945-1166.
- *Bouch, G.R. and Ball, R.C. 1964. Influence of a diurnal oxygen pulse on fish serum proteins. Trans. Am. Fish. Soc., 94: 363-370.
- Boulva, J. 1972. Morphometrics of three sympatric cod fishes of the genera Arctogadus and Gadus. J. Fish. Res. Bd. Can., 29: 243-249.
- Brewer, G.J. 1970. An introduction to isoenzyme techniques. Academic Press, New York. 190 pp.
- Brock, V. 1978. Morphological and biochemical criteria for the separation of Cardium glaucum from Cardium edule. Ophelia., 17 : 207-214.
- Bucklin, A. and Hedegecock, D. 1982. Biochemical genetic evidence for a third species of Metridium (Coelenterata : Actiniaria). Mar. Biol., 66 : 1-7.
- Bulnheim, H.P. and Scholl, A. 1986. Genetic differentiation between populations of Talitrus saltator and Talorchestia deshayesii (Crustacea : Amphipoda) from coastal areas of the North-western European continent Mar. Biol., 92 : 525-536.
- Burch, J.Q. 1956. Taxonomic characters in Mollusca. Syst. Zool., 5 : 144.
- Buroker, N.E., Hershberger, W.K. and Chew, K.K. 1975. Genetic variation in the Pacific oyster, Crassostrea gigas. J. Fish. Res. Bd. Can., 32 : 2471-2477.
- Burton, R.S. and Feldman, M.W. 1982. Population genetics of coastal and

- estuarine invertebrates. In. Estuarine comparisons, V.S. Kennedy (Ed), Academic Press, New York. pp.537-551.
- Busselin, P. 1970.** Effects of moulting cycle and nutritional conditions on haemolymph proteins in Carcinus maenas. Comp. Biochem. Physiol., 37 : 73-83.
- Calhoun, W.B.L. and Koenig, V.L. 1970.** The distribution the soluble proteins in the eye lenses of some marine vertebrates. Comp. Biochem. Physiol., 34: 71-80.
- Carvalho, G.R. and Loney, K.H. 1989.** Biochemical genetic studies on the Pantagonian squid Loligo gahi d'Orbigny. I. Electrophoretic survey of genetic variability. J. Exp. Mar. Biol. Eco., 126 : 231-241.
- Chambers, C.C., Payne, J.F. and Kennedy, M.C. 1979.** Geographic variation in the Dwarf Crayfish, Cambarellus puer hobbs (Decapoda : Combaridae). Crustaceana., 36(1) : 39-55.
- *Chemnitz. 1785.** Neues systematisches Conchylien-Cabinet. 8 (Nurenberg).
- Choudhary, P.C. 1971.** Responses of larval Macrobrachium carcinus to variation in salinity and diet. Crustaceana., 20(2) : 113-120.
- Choudhary, P.S. and Dwivedi, N.S. 1981.** Meristic and morphometric studies of Lactarius lactarius (Schneider), off Bombay coast. J. Ind. Fish. Assoc., 10&11 : 1-4.
- Clayton, J.W., Tretiak, D.N. and Kooyman, A.H. 1971.** Genetics of multiple malate dehydrogenase isoenzymes in skeletal muscle of **Walleye** (Steizostedion istreum istreum). J. Fish. Res. Bd. Can., 28 : 1005-1008.
- *Clessin, 1879.** In Martini-Chemnitz, Conch. Cab, pp. 224-295, pl. 42, figs. 3 & 4. Quoted from "Notes on Lamellibranchs in the Indian museum" (Prashad, B. 1921). Rec. Ind. Mus. XXII : 111-119, Pl. XV.

- Cole, M.A. and Morgan, R.P. 1978a.** Genetic variation in two populations of blue crab, Callinectes sapidus. Estuaries, 3: 202-205.
- Cole, M.A. and Morgan, R.P. 1978b.** Muscle, eye and serum proteins in the blue crab, Callinectes sapidus Rathben, Comp. Biochem. Physiol., 59B : 25-26.
- Colgan, D.J. 1981.** Spatial and temporal variation in the genotypic frequencies of the mussel, Brachiodontes rostratus. Heridity., 46 : 197-208.
- Connel, J.J. 1953.** Studies on the proteins of fish skeletal muscles. I. Electrophoretic analysis of low ionic strength extracts. Biochem. J., 54 : 119-126.
- Costa, R. and Bisol, P.M. 1978.** Genetic variability in deep-sea organisms. Biol. Bull., 155 : 123-133.
- Cox, L.R. 1969.** General features of Bivalvia. In : Treatise on Invertebrate Paleontology, Part N. Vol. I : Mollusca 6, Bivalvia R.C. Moore (Ed.), Geol. Soc. Am. and Univ. of Kansas Press, Lawrence. pp. 2-129.
- Cripps, P.A. and Reish, D.J. 1973.** The effect of environmental stress on the activity of malate dehydrogenase and lactate dehydrogenase in Neanthes arenaceodentata (Annelida : Polychaeta). Comp. Biochem. Physiol., 46B : 123-133.
- Croker, R.A. and Gable, M.F. 1977.** Geographic variation in Western Atlantic populations of Gammarus oceanicus segerstrale (Amphipoda). Crustaceana., 32(1) : 55-73.
- Dando, P.R., Southward, A.J. and Crisp, D.J. 1979.** Enzyme variation in Chthamalus stellatus and C. montagui (Crustacea : Cirripedia). Evidence for the presence of the C. montagui in the Adriatic. J. Mar. Biol. Assn. U.K., 59 : 307-320.

- Dare, P.J. and Edwards, D.B. 1975.** Seasonal changes in flesh weight and biochemical composition of mussels. J. Exp. Mar. Biol. Ecol., 18: 89-97.
- Davidson, K., Roff, J.C. and Elner, L.W. 1985.** Morphological, electrophoretic and fecundity characteristics of Atlantic snow crab, Chionectes opilio, and implications for fisheries management. Can. J. Fish. Aquat. Sci., 42 : 474-482.
- *Davis, B.J. 1964.** Disc electrophoresis-II. Method and application to human serum proteins. Ann. N. Y. Acad. Sci., 121 : 404-427.
- Davis, G.M. 1967.** A systematic relationship of Pomatiopsis lapidaria and Oncomelania hupensis formosana (Prosobranchia : Hydrobiidae). Malacologia., 6(1-2) : 11-43.
- Davis, G.M. 1968a.** A systematic study of Oncomelania hupensis chini (Gastropoda : Hydrobiidae). Malacologia., 7(1) : 17-70.
- Davis, G.M. 1968b.** Biosystematic analysis of Semisulcospira trachea (Gastropoda : Pleuroceridae). Proc. Symp. Mollusca. Vol. I : 16-35.
- Davis, G.M. 1971.** Systematic studies of Bretia costula episcopalis, first intermediate host of Paraquanium westermanni in Malaysia. Proc. Acad. Natur. Sci. Philadelphia., 123(30) : 53-86.
- Davis, G.M. and Lindsay, G.K. 1967.** Disc electrophoretic analysis of molluscan individuals and populations. Malacologia., 5(2) : 311-334.
- Deshayes, 1854.** Cat. Brit. Mus. Conchifera., II. p 240. Quoted from "Notes on Lamellibranchs in the Indian museum" (Prashad, B. 1921), Rec. Ind. Mus., XXII : 111-119, pl.XV.
- Deshmukh, S. 1972.** Some aspects of the biology of Meretrix meretrix. Ph.D. Thesis, Marathwada University, Aurangabad, India.
- De Zwann, A. and Zande, D.I. 1972.** Body distribution and seasonal changes

in the glycogen content of the common sea mussel, Mytilus edulis.
Comp. Biochem. Physiol., 43A : 52-58.

- Dingle, J.R., Eagles, D.E. and Neelin, J.M. 1955.** Electrophoretic studies of proteins extractable from post-rigor cod at low ionic strength. J. Fish. Res. Bd. Can., 12 : 75-84.
- Doyle, R.W. and Hunte, W. 1981.** Genetic changes in fitness and yield of a crustacean population in a controlled environment. J. Exp. Mar. Biol., 52 : 147-156.
- Duranko, M.J., Goddard, R.H., Hoffman, W. and Lawrence, J.M. 1979.** Malate and lactate dehydrogenase activities in the pyloric caeca of Luidia clathrata (Echinodermata : Asteroidea). Comp. Biochem. Physiol., 62B : 127-128.
- Durve, V.S. and Bal, D.V. 1961.** Studies on the chemical composition of the oyster Crassostrea gryphoides (Schlotheim). J. Zool. Soc. India., 13 : 70-77.
- Durve, V.S. 1968.** On the nomenclature of two Indian backwater oysters. J. Mar. Biol. Assn. India., 9 : 173-178.
- Durve, V.S. and Dharmaraja, S.K. 1965.** A study on the dimensional relationship in the clam Meretrix casta (Chemnitz) collected from two localities. J. Mar. Biol. Assn. India., 7(1) : 69-79.
- Durve, V.S. and George, K.V. 1973.** Some observations on the index of condition of the clam, Meretrix casta (Chemnitz) in relation to mud and water qualities. Indian J. Fish., 20(2) : 326-332.
- Easterson, D.C.V. and Kandasami, D. 1988.** Biochemical changes in the oyster, Crassostrea madrasensis (Preston) with maturation. CMFRI Bull., 42 (1) : 203-205.
- *Farmur, A.S.D. 1986.** Morphometric relationship of commercially important

- species of *Penaeus* shrimp from the Arabian Gulf. Kuwait. Bull. Mar. Sci., No. 7 : 1-21.
- Ferguson, A. 1980.** Biochemical systematics and evolution : Glasgow : Blackie. 194 pp.
- Fleminger, A. and Hulsemann, K. 1974.** Systematics of distribution of the four sibling species comprising the genus *Pontinella dara* (Copepoda : Calanoidea). Fish. Bull., 72(1) : 63-120.
- Frankel, J.S. 1980.** Comparative analysis of lactate dehydrogenase isoenzymes in neural tissues of *Poecilia* (Poeciliidae). Comp. Biochem. Physiol., 69B : 881-884.
- Frankel, J.S. and Harth, N.H. 1977.** Lactate dehydrogenase ontogeny in the genus *Brachydanio* (Cyprinidae). J. Hered., 68, 81-86.
- Gabbot, P.A. 1983.** Seasonal and metabolic activities in marine mollusca In : The mollusca Vol. I, P.W. Hochachka, (Ed). Academic Press, New York. pp 165-217.
- Gabbot, P.A. and Bayne, B.L. 1973.** Biochemical effects of temperature and nutritive stress on *Mytilus edulis* L. J. Mar. Biol. Assn. U.K., 53 : 269-286.
- Galtsoff, P.S. 1931.** The weight-length relationship of the Hawaiian Pearl oyster, *Pinctada* species. Amer. Nat., 65 : 423-433.
- Galtsoff, P.S. 1964.** The American oyster, *Crassostrea virginica* (Gmelin). U.S. Fish and Wildlife Serv., Fish. Bull., 64 : 1-480.
- *Garcia, T. 1980.** Electropherogram of the muscle proteins of four species of *Sciacnidae* (Pisces) from urban waters. Rev. Rown. Biol. Sev. Biol. Anim., 25(1) : 23-25.
- Gartner-Kepkay, K.E., Dickie, L.M., Freeman, K.R. and Zouros, E. 1980.** Genetic differences and environments of mussel populations in the

- Maritime provinces. Can. J. Fish. Aquat. Sci., 37 : 775-782.
- George, E.L. and Nair, N.B. 1975.** Biochemical changes in relation to the breeding of Musculista arcuatula Yamamoto and Habe (Bivalvia : Mytilidae). Sci. Cult., 11(4) : 167-169.
- Giese, A.C. 1966.** Lipids in the economy of marine invertebrates. Physiol. Rev. 46 : 244-298.
- Giese, A.C. 1969.** A new approach to the biochemical composition of the molluscs body. Oceanogr. Mar. Biol. Ann. Rev., 7 : 175.
- Gill, P.D. 1978.** Non-genetic variation in isoenzymes of lactate dehydrogenase of Cepaea nemoralis. Comp. Biochem. Physiol., 59B : 271-276.
- Gillespie, J. and Kojima, K. 1968.** The degree of polymorphism in enzymes involved in energy production compared to that in non-specific enzymes in two D. ananassae populations. Proc. Natl. Acad. Sci. U.S.A., 61(3) : 582-585.
- Gooch, J.L. 1975.** Mechanisms of evolution and population genetics. In : Marine ecology, Vol. II, Part. I, O. Kinne. (Ed). Wiley Interscience, London. pp. 349-409.
- Gooch, J.L. and Schopf, T.J.M. 1971.** Genetic variability in the deep sea : relation to environmental variability. Evolution., 26 : 545-552.
- Gooch, J.L., Smith, B.S. and Knupp, P. 1972.** Regional survey of gene frequencies in the mud snail Nassarius obsoletus. Biol. Bull., 142 : 36-48.
- Gopalakrishnan, K. 1974.** Zoogeography of the genus Nematoscelis (Crustacea : Euphasidae). Fish. Bull., 72(4) : 1039-1076.
- Grant, W.S. and Utter, F.M. 1984.** Biochemical population genetics of Pacific herring (Clupea pallasii). Can. J. Fish. Aquat. Sci., 41 : 856-864.

- Grassle, J.F. 1972.** Species diversity, genetic variation and environmental uncertainty. In : Proc. Fifth European Marine Biology Symposium., B. Battaglia (Ed), Piccin Editore, Padua. pp. 19-26.
- Gravelly, F.H. 1941.** Shells and other animal remains found on the Madras Beach. I. Groups other than snails etc. (Mollusca : Gastropoda). Bull. Madras Govt. Mus. N.S. Hist., 5(1) : 1-112.
- Gray, 1825.** Ann. Phil., n. ser., IX, p. 136, Quoted from "Notes on Lamellibranchs in the Indian museum" (Prashad, B. 1921), Rec. Ind. Mus. XXII : 111-119, Pl. XV.
- Griffith and Pidgeon, 1833.** Animal kingdom, XII, pl. 31, fig. 5. Quoted from "Notes on Lamellibranchs in the Indian museum" (Prashad, B. 1921), Rec. Ind. Mus., XII : 111-119, Pl. XV.
- *Hamai, I. 1934a.** Relation between the weight, volume and linear dimensions in Meretrix meretrix (L). Sci. ep. Tohoku. Univ., 4(9) : 203-218.
- *Hamai, I. 1934b.** On the local variations in the shells of Meretrix meretrix (L), with special reference to growth. Sci. Rep. Tohoku. Univ., 9 : 131.
- *Hamai, I. 1935a.** A study of one case in which different environmental conditions produce different types of Meretrix meretrix. Sci. Rep. Tohoku Univ., 10 : 485.
- *Hamai, I. 1935b.** On the growth of the shell of Meretrix meretrix especially with regard to periodicity of growth relatively to the seasonal variation in the environment. Sci. Rep. Tohoku. Univ., 9 : 339.
- * Hanley, 1858.** Proc. Zool. Soc. London., XXVI, p. 543. Quoted from "Notes on Lamellibranchs in the Indian museum" (Prashad, B. 1921), Rec. Ind. Mus., XXII : 111-119, Pl. XV.
- Hanley, 1866.** Ann. Lyc. Nat. ist. Soc., New York, VII, p. 236, fig. 66.

- Quoted from "Notes on Lamellibranchs in the Indian museum" (Prashad, B. 1921), Rec. Ind. Mus., XXII : 111-119, Pl.XV.
- Heath, D.J. 1975.** Geographical variation in populations of the polymorphic isopod, Sphaeroma rugicauda. Heridity., 35(1): 99-107.
- Hedrick, P.W. 1983.** Genetics of populations. (Boston : Science Books International).
- Hedrick, P.W., Ginewan, M.E. and Ewing, E.P. 1976.** Genetic polymorphism in heterogeneous environments. Ann. Rev. Ecol. Syst., 7 : 1-32.
- Herzberg, A. and Pasteur, R. 1975.** The identification of grey mullet species by discgel electrophoresis. Aquaculture., 5 : 99-106.
- Holme, N.A. 1961.** Shell form in Venerupis rhomboides. J. Mar. Biol. Assn. U.K., 41 : 705-722.
- Hornbach, D.J., Mc Leod, M.J., Guttman, S.I. and Seilkop, S.K. 1980.** Genetic and morphological variation in the freshwater clam, Sphaerium (Bivalvia : Sphaeriidae). J. Mollus. Stud., 46: 158-170.
- Hornell, J. 1910.** Note on an attempt to ascertain the principal determining factor in oyster spawning in Madras backwaters (Madras Fish. Investigations, 1908). Madras. Fish. Bull., 4 : 25-31.
- Hornell, J. 1917.** The edible molluscs of the Madras presidency. Madras. Fish. Bull., 11 : 1-51.
- Hornell, J. 1921.** Indian molluscs, Lamellibranchia. J. Bombay. Nat. Hist. Soc., 19 : 50-52.
- Hornell, J. 1922.** The common molluscs of South India, Madras Fish. Bull., 14 : 97-215.
- Hubby, J.L. and Lewontin, R.C. 1966.** A molecular approach to the study of genic heterozygosity in natural populations. 1. The number of alleles at different loci in Drosophila pseudo-obscura. Genetics.,

54: 557-594.

- *Hubendick, B. 1951.** Recent Lymnaeidae : their variation, morphology, taxonomy, nomenclature and distribution. Kungl. Svenska Vetenskapssakademien Handlingar Fjards Serien., 3 : 1-223.
- Hulse, J.H. 1982.** Foreword. In: Bivalve culture in Asia and the Pacific. Proc Workshop held in Singapore, 16-19, Feb. 1982, International Development Research Centre, Ottawa. F.B. Davy and M. Graham (Eds). pp. 5-6.
- Ihssen, P.E., Evans, D.O., Christic, W.J., Reckahn, J.A. and Desfardine, R.L. 1981.** Life history, morphology and electrophoretic characters of five allopatric stocks of lake white fish (Coregonus clupeaformis) in the great lake region. J. Fish. Res. Bd. Can., 38: 1790-1807.
- *International Commission on Zoological Nomenclature, 1955. Opinion 338.** Acceptance of the Mesozoic fossil species Gryphaea arcuata Lamarck, 1801 as the type species of the nominal genus Gryphaea Lamarck, 1801 (Class : Pelecypoda) to the Official List of Generic Names in Zoology. Opinions and declarations rendered by the International Commission on Zoological Nomenclature, 10(5) : 125-180.
- Jayabal, R. and Kalyani, M. 1986.** Biochemical studies in the hard clam, Meretrix meretrix (L) from vellar estuary, east coast of India. Indian J. Mar. Sci., 15 : 63-64.
- Johnson, D.S. 1960.** Subspecific and intraspecific variation in some freshwater prawns of the Indo-Pacific region. In: Proc. of the Centenary and Bicentenary Congress of Biology, Singapore, Dec. 2-9, 1958. R.D. Purchon (Ed). University of Malaya Press, Singapore, pp. 259-267.
- Johnson, A.G., Utter, F.M. and Hodgins, H.O. 1972.** Electrophoretic investigation of the family Scorpeanidae. J. Fish. Res. Bd. Can., 23(10):

1599-1606.

- Jolicoeur, P. and Mosimann, J.E. 1960.** Size and shape variation in the painted turtle. A principal component analysis. Growth, 24 : 339-354.
- Jones, S. 1951.** Observations on the binomics and fishery of the brown mussel (Mytilus sp) of the Cape region of Peninsular India. J. Bombay. Nat. Hist. Soc., 49: 519-528.
- Jones, B.W. and Mackie, I.M. 1970.** An application of electrophoretic analysis of muscle myogens to taxonomic studies in the genus Merluccius. Comp. Biochem. Physiol., 32 : 267-273.
- Joseph, M.M. 1979.** Studies on the biology of the Indian backwater oyster Crassostrea madrasensis (Preston). Ph.D. thesis, Univ. Mysore, 350pp.
- Joseph, M.M. and Madhyastha, M.N. 1986.** Environmental effects on the condition and edibility of the oyster, Crassostrea madrasensis (Preston) in a tropical estuary. In : Environmental Biology, India. : R.C. Delela, M.N. Madhyastha and M.M. Joseph (Eds). pp. 183-188.
- Joshi, M.C. and Bal, D.V. 1965.** Observation on the chemical composition of the clam Katelysia marmorata Lamarck. J. Zool. Soc. India., 17(1): 108-113.
- Kannupandi, N. and Paulpandian, A.L. 1975.** Studies on the blood and muscle proteins of crabs of Porto Novo. Bull. Dept. Mar. Sci. Univ. Cochin., 7(3) : 609-622.
- Karlin, S. 1982.** Theoretical population genetics, Academic Press. N.Y.
- Karlin, A.A. and Guttman, S.I. 1979.** Niche width and heterozygosity. Isoz. Bull., 12 : 50.
- Kasinathan, R. 1974.** Bio-systematic studies on some Cyclophoridae (Mesogastropoda : Mollusca) from Peninsular India. Ph.D. thesis, Annamalai

University.

- Kasinathan, R., Desai, B.N. and Dwivedi, S.N. 1972.** Morphometry and electrophoretic studies of Cynoglossus bilineatus (Lac) from Bombay waters. J. Indian Fish. Assn., 2: 110-115.
- Katticaran, C.M. 1988.** Studies on the biology of the clam, Sunetta scripta (Linne), from the subtidal waters of Cochin. Ph.D. thesis, Cochin Univ. of Science and Technology, India. pp. 33-53.
- Kendall, M.G. 1957.** A course in multivariate analysis. Charles Griffith & Co. Ltd., London, 185 pp.
- King, M.C. and Wilson, A.C. 1975.** Evolution at two levels in humans and chimpanzees. Science., 188 : 107-116.
- Kitching, J.A., Muntz, L. and Ebling, F.J. 1968.** The ecology of Lough Ine. XV. The ecological significance of shell and body forms in Nucella. J. Anim. Ecol., 35 : 119-126.
- Koehn, R.K., Milkman, R. and Mitton, J.B. 1976.** Population genetics of marine pelecypods. IV. Selection, migration and genetic differentiation in the blue mussel Mytilus edulis. Evolution., 30 : 2-32.
- Koehn, R.K. and Mitton, J.B. 1972.** Population genetics of marine pelecypods. I. Ecological heterogeneity and evolutionary strategy at an enzyme locus. Amer. Nat., 106 : 47-56.
- Koehn, R.K., Turano, F.J. and Mitton, J.B. 1973.** Population genetics of marine pelecypods. II. Genetic differences in microhabitats of Modiolus demissus. Evolution., 27 : 100-105.
- Korringa, P. 1952.** Recent advances in oyster biology. Quart. Rev. Biol., 27 : 266-308.
- Krishnakumari, L., Rajagopal, M.D. and Vijayaraghavan, S. 1976.** Nutritive value of the rock oyster Crassostrea cucullata. Mahasagar., 8 : 203-205.

- Krishnakumar, L., Rajagopal, M.D. and Vijayaraghavan, S. 1977.** Some aspects of biology and biochemistry of the backwater clam Meretrix casta (Chemnitz). Mahasagar. Bull., 10(3): 157-163.
- Kulkarni, G.K. Nagabhushanam, R.N. and Joshi, P.K. 1980.** Electrophoretic separation of protein patterns of different tissues of four marine penaeid prawns in relation to sex. Hydrobiologia., 69(1) : 25-28.
- Kulkuhn, J.H. 1981.** Stock definition as a necessary basis for co-operative management of Great Lakes fish resources. Can J. Fish. Aquat. Sci., 1476-1478.
- Kundu, H.L. 1965.** On the marine fauna of Gulf of Kutch, Part III. Pelecypods. J. Bombay. Nat. His. Soc., 62 : 84-103.
- Kuriakose, P.S. and Nair, N.B. 1976.** The genus Perna along the coasts of India with the description of a new species Perna indica. Aquat. Biol., 1 : 25-36.
- Lakshmanan, P.T. and Nambisan, P.N.K. 1980.** Biochemical composition of the bivalve molluscs, Villorita cyprinoides. var. cochinensis (Hanley) and Meretrix casta (Chemnitz). Indian J. Mar. Sci., 9 : 65-67.
- *Lamarck, J.B. 1801.** Systeme des animaux sans vertebres. Crapelot, Paris, pp.432.
- Lear, W.H. and Misra, R.K. 1978.** Clinal variation in scale characters of Atlantic salmon (Salmo salar) based on discriminant function analysis. J. Fish. Res. Bd. Can., 35 :43-47.
- Lee, S.S. and Lim, T.K. 1973.** Electrophoretic serum protein patterns in some penaeid prawns (Crustacea : Decapoda). Malaysian Agric. J., 49 (1) : 11-15.
- Levinton, J.S. 1973.** Genetic variation in a gradient of environmental stability : Marine Bivalvia (Mollusca). Science., 180 : 75-76.

- Levinton, J.S. 1980.** Genetic divergence in estuaries. In : Estuarine Perspectives, V.S. Kennedy (Ed), Academic Press, New York). pp.509-520.
- Levinton, J.S. 1982.** Marine Ecology. Prentice-Hall, Englewood Cliffs, New Jersey.
- Levinton, J.S. and Koehn, R.K. 1976.** Population genetics of mussels. In : Marine mussels, Their Ecology and Physiology, B.L. Bayne (Ed), Cambridge University Press, Cambridge. pp.357-384.
- Lewontin, R.C. 1984.** Detecting population differences in quantitative characters as opposed to gene frequencies. Amer. Natur., 123 : 115-124.
- Lewontin, R.C. and Hubby, J.L. 1966.** A molecular approach to the study of genetic heterozygosity in natural populations. II. Amount of variation and degree of heterozygosity in natural populations of Drosophila pseudoobscura. Genetics., : 595-609.
- Lim, T.K. and Lee, S.S. 1970.** Electrophoretic studies on muscle myogen of some penaeid prawns. Mar. Biol., 5 : 83-88.
- *Linnaeus, C. 1758.** Systema Naturae (Ed), 10 : 671.
- Lindenfelser, M.E. 1980.** Subspecific variation and evolution in the fresh water prawn, Macrobrachium rosenbergii (de Man) (Decapoda : Palaemonidae) as reflected in its morphometric and allozymic variation. Ph.D. thesis, Univ. Hawaii.
- Long, G.L. 1976.** The stereospecific distribution and evolutionary significance of invertebrate lactate dehydrogenases. Comp. Biochem. Physiol., 55B : 77-83.
- Love, M.R. 1970.** The Chemical Biology of Fishes. Academic Press, London.
- Lowry, O.H., Rosenbrough, N.J., Farr, A.L. and Randall, R.J. 1951.** Protein measurement with the Folin-phenol reagent. J. Biol. Chem., 193 : 265-275.

- Lyerla, T.A. and Lyerla, J.H. 1978.** A preliminary study of electrophoretic variation of enzymes in the tunicates Clavelina picta and C. oblonga. Comp. Biochem. Physiol., 59B : 111-116.
- Mackie, I.M. and Jones, B.W. 1978.** The use of electrophoresis of the water soluble (sarcolemmic) proteins of fish muscle to differentiate the closely related species of Hake (Merluccius sp.). Comp. Biochem. Physiol., 59B : 95-98.
- Maguire, G.B. and Fielder, D.R. 1975.** Disc electrophoresis of the haemolymph proteins of some portunid crabs (Decapoda : Portunidae)-1. Physiological and Taxonomic aspects. Comp. Biochem. Physiol., 52A : 43-47.
- Mane, U.H. 1974.** The adaptations of the estuarine clam, Katelysia opima to the salinity fluctuations. Riv. deBiol., 67 : 109-139.
- Mane, U.H. and Nagabhushanam, R. 1979.** Studies on the growth and density of the clam, Paphia laterisulca at Kalabadevi estuary, Ratnagiri on the west coast of India. Malacologia., 18 (2) : 297-313.
- Manwell, C. and Ann Baker, C.M. 1963.** A sibling species of a sea cucumber discovered by starch gel electrophoresis. Comp. Biochem. Physiol., 10 : 39-53.
- Manwell, C. and Ann Baker, C.M. 1970.** Molecular biology and the Origin of Species. Univ. of Washington Press, Seattle, 394pp.
- Manwell, C., Ann Baker, C.M. and Corner, E.D.S. 1967.** Biochemical differences between Calanus finmarchicus and C. helgolandicus. J. Mar. Biol. Ass. U.K., 47 : 145-169.
- Margulis, B.A. and Pinaev, G.P. 1976.** The species specificity of the contractile protein composition of the bivalve molluscs. Comp. Biochem. Physiol., 55B 189-194.

- Markert, C.L. 1965.** Mechanism of cellular differentiation. In : Ideas in Modern Biology. Natural History Press, Garden City, New York.
- Markert, C.L. and Faulhaber, I 1963.** Lactate dehydrogenase isozyme patterns of fish. J. Exp. Zool., 159 : 319-332.
- Markert, C.L. and Moller. F. 1939.** Multiple forms of enzymes : tissue, ontogenic and species specific patterns. Proc. Natl. Acad. Sci. U.S.A., 45 : 753-763.
- Massaro, E.J. 1972.** Isozyme patterns of Coregonine fishes : Evidence for multiple cistrons for lactate and malate dehydrogenases and achromatic bands in the tissues of Prosobium cylindrecum (Pallas) and P. coulteri (Eigemann and Eigenmann). J. Expl. Zool., 179 : 247-262.
- Massaro, E.J. and Cohen, E. 1978.** An electrophoretic analysis of selected enzymes of the coconut crab, Birgus latro. Comp. Biochem. Physiol., 68(B) : 151-152.
- Mc Donald, J.H. 1985.** Size-related and geographic variation at two enzyme loci in Megalorchestia californiana (Amphipoda : Talitridae). Heredity., 54 : 359-366.
- McKenzie, J.A. 1973.** Comparative electrophoresis of tissues from blueback herring Alosa aestivalis (Mitchill and gaspareau, Alosa psedoharengus (Wilson). Comp. Biochem. Physiol., 44B : 65-68.
- Mickevich, M. and Johnson, M.S. 1976.** Congruence between morphological and allozyme data in evolutionary inference and character evolution. Syst. Zool., 25 : 260-270.
- Milkman, R. 1983.** Experimental population genetics (Stroudsburch ; Hutchinson & Ross).
- Milkman, R. and Beaty, L.D. 1970.** Large scale electrophoretic studies of allelic variation in Mytilus edulis. Biol. Bull., 139 : 430.

- Mohan, M.V. 1980.** Allometric relationship in the green mussel, Perna viridis. Indian. J. Mar. Sci., 9 : 224-226.
- Mohan, M.V. and Damodaran, R. 1981.** Allometric relationship in the clam, Sunetta scripta L. Indian J. Mar. Sci., 10(6) ; 198-200.
- Mohan, M.V., Damodaran, R and Salih, K.Y.M. 1986.** Allometric relationships in the wedge clam Donax cuneatus Gmelin. Mahasagar Bull., 19 (1) : 57-60.
- Montgomery, R. 1957.** Determination of glycogen. Arch.Biochem. Biophys., 67 : 378-386.
- Moore, H.B. 1936.** The biology of Purpura. 1. Shell variation in relation to environment. J. Mar. Biol. Assn. U.K., 21 : 61-89.
- *Morgan, G.R. 1982.** Stock and recruitment relationship in the shrimp, Penaeus semisulcatus. KSIR safat (Kuwait)., Jul : 228-241.
- Morgan, R.P., Block, S.B., Ulbanowicz, N.I. and Buys, C. 1978.** Genetic variation in the soft-shelled clam, Mya arenaria. Estuaries., 1 ; 255-258.
- Morrissey, R.E. 1981.** Alcohol dehydrogenase activity in tissues of the house cricket Acheta domesticus. Comp. Biochem. physiol., 68B : 133-135.
- Morrow, J.E. 1980.** Analysis of the dolly varden char Salvelinus malma of North Western, North American and North Eastern Siberia. In: E.K. Balan (Ed), Charrs : Salmonid fishes of the genus Salvelinus, Perspectives in Vertebrate Science. Vol. I. Dr.W.Junk, Publ., The Hague, Netherlands.
- Morton, J.E. and Yonge, C.M. 1964.** Classification and structure of the Mollusca. In : Physiology of Mollusca . K. Wilbur and C.M. Yonge (Eds), Academic Press, New York. pp.1-58.

- Moses, S.T. 1928.** A preliminary report on the anatomy and life-history of the common edible backwater oyster, Ostrea madrasensis. J. Bombay. Nat. Hist. Soc., 32 : 548-552.
- Murphy, P.G. 1978.** Collisella austrodigitalis sp. Nov : A sibling species of limpet (Acmaeidae) discovered by electrophoresis. Biol. Bull., 155 : 193-206.
- Nagabhushanam, R. and Deshmukh, R.S. 1974.** Seasonal changes in body component indices and chemical composition in the estuarine clam, Meretrix meretrix. Indian J. Fish., 21 : 531-542.
- Nagabhushanam, R. and Dhamne, K.P. 1975.** Seasonal variations in biochemical constituents of the clam Paphia laterisulca. Hydrobiologia., 54 (3) : 209-214.
- Nagabhushanam, R. and Mane, U.H. 1978.** Seasonal variation in the biochemical compositions of Mytilus viridis at Ratnagiri on the west coast of India. Hydrobiologia., 57 : 69-72.
- Nagabhushanam, R. and Thalikhedkar, P.M. 1977.** Seasonal variations in proteins, fat and glycogen of the wedge clam Donax cuneatus. Indian J. Mar. Sci., 6 : 85-87.
- Nair, S.G. and Shynamma, C.S. 1975.** Seasonal changes in the lipid and calorific values in Villorita cyprinoides var. cochinesis (Hanley). Bull. Dept. Mar. Sci. Univ. Cochin., 7 (2) : 403-407.
- Nair, U.N. and Nair, N.B. 1985.** Height-Depth (Breadth) relationship of shells in the backwater oyster Crassostrea madrasensis (Preston) of the Cochin Harbour. Fish. Technol., 22 : 87-91.
- Narang, S. 1974.** Lactate dehydrogenase of Biomphalaria glabrata (Say, 1818) (Mollusca : Pulmonata) - I. Physico-chemical characterization of lactate dehydrogenase of various tissues. Comp. Biochem. Physiol., 478 : 641-655.

- Natarajan, V., Kannupandi, T., Paulpandian, A.L. and Natarajan, R. 1975.** Electrophoretic studies on muscle proteins of some gobioids of Porto Novo. Bull. Dept. Mar. Sci. Univ. Cochin., 7 (3) : 583-589.
- Nei, M. 1975.** Molecular population genetics and evolution. Amsterdam, North Holland. 288 pp.
- Newell, N.D. 1965.** Classification of the Bivalvia. Am. Mus. Novit. No. 2206 : 1-25.
- Newell, N.D. 1969.** Classification of Bivalvia. In : Mollusca, 6. Bivalvia. R.C. Moore (Ed), pp. N 205-N 224.
- NewCombe, C.L. 1935.** Growth of Mya arenaria, L in the Bay of Fundy region. Can. J. Res., 13 (6) : 97-137.
- NewCombe, C.L. 1950.** An analysis of certain dimensional relationship of the virginia oyster, Crassostrea virginica (Gmelin). Amer. Nat., 84 : 203-204.
- NewCombe, C.L., Thompson, S.J. and Kessler, H. 1938.** Variation in growth indices of Venus mercenaria L. from widely separated environments of the Atlantic coast . Amer. Nat., 16 (1) : 1-5.
- Ornstein, L. 1964.** Disc electrophoresis. I. Background and theory. Ann. N. Y. Acad. Sci., 121 : 321-349.
- Pace, G.L. and Lindsay, G.K. 1965.** Electrophoretic studies of some diploid Bulinae. Bull. Malacol. Union., 32 : 31-32.
- Page, L.M. and Whitt, G.S. 1973.** Lactate dehydrogenase, malate dehydrogenase, and tetrazolium oxidase mobilities of darters (Etheostomatini). Comp. Biochem. physiol., 44 : 611-623.
- Paul, M.D. 1942.** Studies on the growth and breeding of certain sedentary organisms from the Madras harbour. Proc Indian Acad. Sci., 15 : 1-42.

- Pelseneer, P. 1906.** Mollusca. In : Lankester's treatise on Zoology. Adam & Charles Black London. 355pp.
- *Pfleiderer, G. and Jeckel, C.D. 1957.** Individuelle Milchsäuredehydrogenasen bei verschiedenen Säugetieren. Biochem. Z., 329 : 370-380.
- Phillips, B.F., Campbell, N.A. and Wilson, B.R. 1973.** A multivariate study of geographic variation in the whelk Dicathais. J. Exp. Mar. Biol. Ecol., 11 : 27-69.
- Pieters, H., Kluitmans, J.H., Zandee, D.I. and Cadee, G.C. 1980.** Tissue composition and reproduction of Mytilus edulis in relation to food availability. Neth. J. Sea Res., 14 : 349-361.
- Ponniah, A.G. 1985.** A study on biochemical genetics on Crassostrea madrasensis of Cochin. CMFRI Bull., 42. (1) : 189-192.
- Powell, J.R. and Taylor, C.E. 1979.** Genetic variation in ecological diverse environments. Amer. Scientist., 67 : 590-596.
- Prakash, R. and Yadav, J.P. 1989.** Electrophoretic analysis of eight gene-enzyme systems in four natural populations of Drosophila immigrans. Proc. Ind. Natl. Acad., B55 (5&6) : 387-392.
- Prashad, B. 1921.** Notes on Lamellibranchs in the Indian Museum. Rec. Ind. Mus., XXII : 111-119, pl XV.
- Prathibha, P. 1984.** Electrophoretic studies on Penaeus monodon Fabricius. M.Sc. Dissertation, Cochin. Uty. of Science and Technology, India.
- Preston, H.B. 1915.** Fauna. Brit. India, Mollusca, Gastropoda & Pelecypoda (Freshwater), pp.209.
- Preston, H.B. 1916.** Report on a collection of Mollusca from the Cochin backwaters. Rec. Ind. Mus., 12 : 27-39.
- * Prime, 1860.** Proc. Acad. Nat. Sciences, Philadelphia, p.80. Quoted from "Notes on Lamellibranchs in the Indian Museum" (Prashad. B. 1921) Réc. Ind. Mus., XXII : 111-119. pl, XV.

- * Prime, 1869. Cat. Corbiculidae. In : Amer. J. conch.,V, p.141. No.2. quoted from "Notes on Lamellibranchs in the Indian museum" (Prashad, B. 1921), Rec. Ind. Mus., XXII : 111-119, pl.XV.
- * Prime, 1870. Quoted from "Notes on Lamellibranchs in the Indian museum." (Prashad, B. 1921), Rec. Ind. Mus., XXII : 111-119, Pl.XV).
- Qasim, S.Z. and Gopinathan, C.K. 1969. Tidal cycle and the environmental features of Cochin Backwater (A tropical estuary). Proc. Ind. Acad. Sci., 69 51-94.
- Rai, H.S. 1932. The shell fisheries of the Bombay Presidency. J. Bombay. Nat. Hist. Soc., 33(4) : 893-898.
- Rainboth, M.J. Jr. and Whitt, G.S. 1974. Analysis of evolutionary relationships among shiners of the subgenus Luxilus (Teleostei: Cypriniformes, Notropis) with the lactate dehydrogenase and malate dehydrogenase isozyme systems. Comp. Biochem. Physiol., 49 : 241-252.
- Ramachandran, K. 1980. Studies on the biology of the green mussel, Perna viridis (Linnaeus). M.F.Sc thesis, Univ. Agri. Sci., Bangalore, India. 219pp.
- Ranson, G. 1948. Prodissoconques et classification desostreides vivantes. Bull. Mus. Hist. Nat. Belg., 24 : 1-12.
- *Ranson, G. 1950. La chambre promyaire et la classification zoologique des ostreides. J. Conchyliol., 90 : 195-200.
- Rao, S. 1974. Edible bivalves : Mussels and oysters. In : The Commercial Molluscs of India. R.V. Nair and K.S. Rao (Eds), ICAR. CMFRI. Bull., No. 25 : 4-49.
- Rao, K.V. 1956. Seasonal changes in the adult backwater oyster Ostrea (Crassostrea) madrasensis (Preston) from Ennur near Madras. Proc. Indian Acad. Sci., 44B : 332-356.

- Rao, K.V. 1958.** Molluscan Fisheries. In : Fisheries of West coast of India. S. Jones (Ed), pp.55-59.
- Reddy, D.K.N. 1983.** Studies on the population ecology and reproductive biology of the clam , Villorita cyprinoides (Grey). M.F. Sc. thesis, Univ. Agril. Sci. Bangalore, India. 100pp.
- *Reeve, L.A. 1857.** Conchologia jconica, 10, Monograph of the Genus Avicula. London.
- Riddell, B.E. and Leggett, W.C. 1981.** Evidence of an adaptive basis for geographic variation in body morphology and time of downstream migration of juvenile Atlantic salmon (Salmo salar). Can. J. Fish. Aquat. Sci., 38 : 308.
- *Royce, W.F. 1957.** Statistical comparison of morphological data. USWFS Spec. Sci. Report. Fisheries., No.208 : 7-28.
- Ryman, N., Allendorf, F.W. and Stahl, G. 1979.** Reproductive isolation with little genetic divergence in sympatric populations of brown trout (Salmo trutta). Genetics., 92 : 247-262.
- Ryman, N. and Stahl. G. 1981.** Genetic perspective of the identification and conservation of Scandinavian stocks of fish. Can. J. Fish. Aquat. Sci., 38 : 1562-1575.
- *Sacco, 1897.** i molluschi dei terreni terziarri del Picmonte e della, Liquria- Part 23 (Osteridae, Anomidae e Dimydia). Boll. Musei. Zool. Anat. Comp. R. Univ. Torino., 12 (298) : 99-100.
- Saila, S.B. and Flowers, J.M. 1969.** Geographic morphometric variation in the American lobster. Syst. Zool., 18 : 330-338.
- Sailas, E.G. and Pillai, P.P. 1975.** Dynamics of zooplankton in a tropical estuary (Cochin backwater), with a review on the plankton fauna of the environment. Bull. Dept. Mar. Sci. Univ. Cochin., 7(2) : 329-355.

- Salih, K.Y.M. 1979. Studies on the biochemical composition of the clam, Meretrix casta (Chemnitz) off Cochin barmouth. Bull. Dept. Mar. Sci. Univ. Cochin., 10 ; 47-73.
- Salmon. M., Ferris, S.D. Johnston, D., Hyatt, G. and Whitt, G.S. 1979. Environmental and biochemical evidence for species distinctiveness in the fiddler crabs, Uca speciosa and U. spincicarpa. Evolution., 33(1) : 182-191.
- Samuel, P.P. 1987. Biochemical genetics of selected commercially important penaeid prawns. Ph. D. thesis, CMFRI Cochin, Cochin Univ. Science and Technology, India.
- Sankaranarayanan, V.N. and Qasim, S.Z. 1969. Nutrients of the Cochin Backwater in relation to environmental characteristics. Mar. Biol., 2 : 236-247.
- Sathyamurthi, S.T. 1956. The Mollusca of Krusadi Island (in Gulf of Mannar) II. Scaphopoda, Pelecypoda and Cephalopoda. Bull. Madras Govt. Mus., (New Ser) Nat. Hist. Sect., 1(2) : 202pp, pt.7.
- Sathyamurthi, S.T. 1960. The land and fresh water Mollusca. Bull. Madras Govt. Mus. 6(4) : 143-152.
- *Schmidt, J. 1917 a. Racial Investigations. I. Zoarcis viviparus L. and the local races of the same. C. R. Trav. Lab. carlsberg, 13 (30 : 277. - 397.
- *Schmidt, J. 1917b. Racial investigations. II. Constancy investigations continued C.R. Trav. Lab. Carlsberg., 14 (1) : 1-19.
- Schmidt, J. 1918. Racial studies in fishes. I. Statistical investigations with Zoarcis viviparus L. J. Genet., 7 : 105-110.
- Schoph, T.J.M. and Gooch, J.L. 1971. Gene frequencies in a marine ectoproct : a cline in natural populations related to sea temperature. Evolution., 25 : 286-289.

- Schoph, T.J.M. and Murphy, L.S. 1973. Protein polymorphism of the hybridizing sea stars Asterius forbesi and A. vulgaris and implication for their evolution. Biol. Bull., 145 : 589-597.
- Selander, R.K., Yang, S.Y., Lewontin, R.C. and Johnson, W.E. 1970. Genetic variation in the horseshoe crab (Limulus polyphemus), a phylogenetic 'relic'. Evolution., 24 : 402-414.
- Shafee, M.S. 1976. Studies on the various allometric relationships in the intertidal green mussel, Perna viridis L. of Ennore estuary, Madras. Indian J. Fish., 23 : 1-9.
- Shafee, M.S. 1978. Variations in biochemical composition of the green mussel Perna viridis L. of Ennur estuary, Madras. Mahasagar Bull. 11(1) : 101-104.
- Sharp, J.C.K., Able, W., Leggett, W.C. and Carscadden, J.E. 1978. Utility of meristic and morphometric characters for identification of capelin (Mallotus villosus) stocks in Canadian atlantic waters. J. Fish. Res. Bd. Can., 35 : 124-130.
- Shreni, K.D. 1980. Seasonal variations in the chemical composition of the catfish, Heteropneustes fossilis (Bluch). Pro. Ind. Acad. Sci (Anim.. Sci.), 89 (2) : 191-196.
- Siciliano, M.J. and Shaw, R. 1976. Separation and visualisation of enzymes on gels. In : Chromatographic and electrophoretic techniques. Vol. II. Zone electrophoresis, I. Smith (Ed), 4th ed. Heinemann medical Books, London.pp.185-209.
- Siegismund, H.R., Simonsen, V and Kolding, S. 1985. Genetic studies of Gammarus. Hereditas., 102 : 1-13.
- Simpson, G.C. 1961. Principles of Animal Taxonomy. N.C.Dunn (Ed), Columbia Uty. Press, Columbia, 247 pp.

- Singh, S.M. and Green, R.H. 1984. Excess of allozyme homozygosity in marine molluscs and its possible biological significance. Malacologia., 25 : 569-581.
- Singh, R.S., Hickery, D.A. and David, J.R. 1982. Genetic differentiation between geographically distant populations of Drosophila melanogaster. Genetics., 101 : 235-256.
- Skaala, D., Dabic, G., Jorstad, N.L. and Naevdal, G. 1990. Interactions between natural and farmed fish populations, information from genetic markers. J. Fish. Biol., 36 : 449-460.
- *Skibinski, D.O.F., Mc Nee, A.R. and Beardmore, J.A. 1978. Protein variation in the marine bivalve Scrobicularia plana. Animal Blood Groups and Biochemical Genetics., 9 ; 223-228.
- Snedecor, G.W. and Cochran, W.G. 1967. Statistical methods. (6th edn). The Iowa State University Press, Iowa, U.S.A. 593pp.
- Sokal, R.R. and Thomas, P.A. 1968. Geographic variation of Pemphigus populi-transversus in eastern North-America : Stem mothers and new data on alates. Univ Kansas Sci. Bull., 46 : 20-251.
- Somero, G.M. and Soule, M. 1974. Genetic variation in marine fishes as a test of the niche-variation hypothesis. Nature., 249 : 670-673.
- *Soot-Ryen, T. 1955. A report on the family Mytilidae. Allen Hancock Pacific Expeditions., 20 ; 1-154, pls. 1-10. figs. 1-78.
- * Sowerby, 1878. In: Reeve, Conch. Icon., XX Velorita, pl. 1, figs. 1a-c. Quoted from "Notes on Lamellibranchs in the Indian museum" (Prashad, B. 1921), Rec. Ind.Mus., XXII : 111-119, pl.XV.
- Sreeraman, K. and Reddy, P.S. 1977. Biochemical studies in planktonic juveniles and adults of Penaeus indicus and P. monodon (Fabricius). Proc Sympo. Warmwater Zooplankton, NIO, Goa.

- Srinivasan, V.V. and Krishnaswamy, S. 1963.** Observation on the chemical composition of Martesia fragilis. Life Sci., 3 :149-151.
- Stenzel, H.B. 1971.** Oysters. In : Treatise on Invertebrate Paleontology N 953-N, Mollusca 6. K.C. Moore (Ed), Geographical Society of America Inc. and the University of Kansas, Boulder, Colorado. pp.1224.
- Suryanarayan, H. and Alexander, K.M. 1972.** Biochemical investigations on edible molluscs of Kerala. I. A study on the nutritional value of some bivalves. Fish. Technol., 9(1) : 42-47.
- Suryanarayan, H. and Nair, N.B. 1976.** Seasonal variations in the biochemical constituents of Cellana radiata. Indian. J. Mar. Sci., 5: 126-128.
- Swan, E.F. 1953.** The effect of substratum on the growth rate, shell-weight and shape of Mya arenaria. Fourth annual conference on clam research, U.S. Fish and Wildlife Serv. Boothbay Harbour. Maine: 43-46.
- Taggart, J., Ferguson, A. and Mason, F.M. 1981.** Genetic variation in Irish populations of brown trout (Salmo trutta L.) : Electrophoretic analysis of allozymes. Comp. Biochem. Physiol., 8 : 393-412.
- Taniguchi, N. 1969.** Comparative electropherograms of muscle proteins of three species of lizard fishes referable to the genus Saurida. Bull. Jap. Soc. Sci., Fish., 35 : 885-890.
- Taniguchi, N., Ochiai, A. and Miyazaki, T. 1972.** Comparative studies of the Japanese platycephalid fishes by electropherograms of muscle proteins, LDH and MDH. Jap. J. Ichthyol., 19 : 89-96.
- Taniguchi, M. and Sakata, K. 1977.** Interspecific and intraspecific variations of muscle protein in the Panamese crucian carp. 2. Starch gel electrophoresis. Jap. J. Ichthyol., 24(1): 1-11.

- ***Telebici, A. and Dimoftache, G. 1972.** Comparative aspects of the dynamics of the general biochemical composition in some bivalves of the Romanian Black Sea littoral. Cercet. Mar., 3 : 129-140.
- ***Theobald, 1876.** Cat.Shell. Brit.India. p.44. Quoted from "Notes on Lamellibranchs in the Indian museum" (Prashad, B. 1921), Rec. Ind. Mus., XXII : 111-119, Pl.XV.
- ***Thiele, 1931.** Handbuch der systematischen Weichtierkunde, Zweiter Band (Scaphopoda, Pelecypoda and Cephalopoda). I : Fischer, Stuttgart. pp.377-461.
- Thomas, M.M. 1981.** Preliminary results of electrophoretic studies on marine prawns. Indian J. Fish., 28 (1 & 2) : 292-294.
- Thomas, P.A. 1968a.** Geographic variation of the rabbit tick, Haemaphysalis leporispalustris, in North America. Univ. Kansas Sci. Bull., 47 : 787-828.
- Thomas, P.A. 1968b.** Variation and covariation in characters of the rabbit tick, Haemaphysalis leporispalustris. Univ. Kansas Sci. Bull., 47 : 829-862.
- Todd, T.N., Smith, G.R. and Cable, L.E. 1981.** Environmental and genetic contributions to morphological differentiation in Ciscoes (Coregoninae) of the Great Lakes. Can. J. Fish. Aquat. Sci., 38 : 59-67.
- Tracey, M.L., Bellet, N.F. and Gravem, C.D. 1975.** Excess allozyme homozygosity and breeding population structure in the mussel, Mytilus californianus. Mar. Biol., 32 : 303-311.
- Tsukuda, H. 1975.** Temperature dependency of the relative activities of liver lactate dehydrogenase isozymes in goldfish acclimated to different temperatures. Comp. Biochem. Physiol., 52B : 343-345.

- Tsuyuki, H. and Roberts, E. 1963.** Specific differences of some members of salmonidae based on their muscle myogen patterns. J. Fish. Res. Bd. Can., 20(1) : 101-104.
- Tsuyuki, H. and Roberts, E. 1965.** Zone electrophoretic comparison of muscle myogens and blood proteins of artificial hybrids of Salmonidae with their parental species. J. Fish. Res. Bd. Can., 22 : 763-773.
- Tsuyuki, H. and Roberts, E. 1966.** Interspecies relationships within the genus Oncorhynchus based on biochemical systematics. J. Fish. Res. Bd. Can., 23 : 101-107.
- Tsuyuki, H., Roberts, E. and Gadd, R.E.A. 1962.** Muscle proteins of Pacific salmon (Oncorhynchus) III. The separation of muscle proteins soluble in low ionic strength salt solutions by starch gel electrophoresis. Curr. J. Biochem. and Physiol., 40 ; 929-936.
- Tsuyuki, H., Roberts, E., Kerr, R.H., Uthe, J.F. and Clarke, L.W. 1967.** Comparative electropherograms of the family Catostomidae. J. Fish. Res. Bd. Can., 24 : 299-304.
- Tsuyuki, H., Roberts, E., Lowes, R.H., Hadway, W and Westerheim, J. 1968.** Contribution of protein electrophoresis to rockfish (Scorpenidae) systematics. J. Fish. Res. Bd. Can., 25(11) : 2477-2501.
- Tsuyuki, H., Roberts, E. and Vanstone, W.E. 1965a.** Comparative zone electropherograms of muscle myogens and blood haemoglobins of marine and fresh water vertebrates and their application to biochemical systematics. J. Fish. Res. Bd. Can., 22(1) : 203-213.
- Tsuyuki, H., Roberts, E., Vanstone, W.E. and Markert, J.R. 1965b.** The species specificity and constancy of muscle myogen and haemoglobin electropherograms of Onchorhynchus. J. Fish. Res. Bd. Can., 22(1) : 215-217.

- Uglow, R.F. 1969.** Haemolymph protein concentration of portunid crabs. II. The effects of imposed fasting on Carcinus maenas. Comp. Biochem. Physiol., 31 : 959-967.
- Uthe, J.F., Roberts, E., Clarke, L.W. and Tsuyuki, H. 1966.** Comparative electropherograms of representative of the families Petromyzontidae, Esocidae, Centrarchidae and Percidae. J. Fish. Res. Bd. Can., 23 : 1663-1671.
- Van Valen, L. 1965.** Morphological variation and width of ecological niche. Amer. Nat., 377 :-390.
- Venkataraman, R. and Chari, S.T. 1951.** Studies on oysters and clams : biochemical variations. Indian J. Med. Res., 39 : 533-541.
- Venkataramiah, A., Lakshmi, G.J. and Gunter, G. 1975.** A review of environmental and nutritional factors on brown shrimp, Penaeus azeticus Ives. in laboratory cultures. 10th Europ Symp. Mar. Biol., 1 : 523-547.
- Walne, P.R. 1970.** The seasonal variation of meat and glycogen content of seven populations of oysters Ostrea edulis L. and a review of the literature. Fishery Investigations. Ministry of Agriculture, Fisheries and Food, London, Ser.II, 26, 35pp.
- Whitmore, D.H. 1980.** Identification of sunfish species by muscle protein isoelectric focusing. Comp Biochem. Physio., 84B : 177-180.
- Whitt, G.S. 1970.** Genetic variation of supernatant and mitochondrial malate dehydrogenase isozymes in the teleost, Fundulus heteroclitus. Experientia., 26 : 734-736.
- *Wieland, T.A., Pfeleider, A., Haput, I. and Worner, W. 1959.** Über die verchieden beit der Milchs aller dehydrogenasen. IV. Quantitative ermittlung einiger enzyme vesteilungsmuster. Vergleichende Betrachtung Bei Verschiedencu Wirebeltien Klassen. Biochem., 332 : 1-10.

- Wilkins, N.P. 1975.** Phosphoglucose isomerase in marine molluscs. In : Isozymes. IV. Genetics and Evolution. C.L. Market (Ed), Academic Press, New York. pp.931-943.
- Wilkins, N.P. and Mathers, N.F. 1973.** Enzyme polymorphisms in the European oyster, Ostrea edulia L. Anim. Blood Groups Biochem. Genet., 4 : 41-47.
- Wilson, A.C., Maxson, L.R. and Sarich, V.M. 1974.** Two types of molecular evolution. Evidence from studies of interspecific hybridization. Proc. Natl. Acad. Sci. U.S.A., 71 : 2843-2847.
- Wilson, A.C., Sarich, V.M. and Maxon, L.R. 1973.** The importance of gene rearrangement in evolution. Evidence from studies on the rate of chromosomal protein and anatomical evolution. Pro Natl. Acad. Sci. U.S.A., 71 : 3028-3030.
- Wilson, B.R. and Summers, R. 1966.** Variation in the Zoila friendi (Gray) species complex (Gastropoda : Cypraeidae) in South-western Australia. J. Malacol. Soc. Aust., 9 : 3-24.
- Winkworth, R. 1931.** Mollusca from the Pulicat Lake. Proc. Malacol. Soc. Lond., 19 : 188-189.
- * **Wood, 1828.** Index. Test., Suppl., pl.2, fig.14. Quoted from "Notes on Lamellibranchs in the Indian museum" (Prashad, B. 1921). Rec. Ind. Mus., XXII : 111-119, pl.XV.
- * **Wright, C.A. and Ross, G.C. 1965.** Electrophoretic studies of some planorbid egg proteins. Bull. Wld. Hlth. Org., 32 : 709-712.
- Wyatt, T. and Qasim, S.Z. 1973.** Application of a model to an estuarine system. Limnol. Oceanogr., 18(2) : 301-306.
- Yonge, C.M. 1960.** Oysters. Collins, London, 209pp.
- Zouros, E. and Foltz, D.W. 1984.** Possible explanations of heterozygote deficiency in bivalve molluscs. Malacologia., 25 : 583-591.

* Not referred to the original