

---

**CONTENTS**

<b>1. Introduction</b>	
<b>2. Review of literature</b>	4
2.1 Endocytosis	5
2.2 Clathrin mediated endocytosis	5
2.3 Intracellular trafficking	6
2.4 Rab proteins: regulators of intracellular transport	6
2.5 Characteristic structural features of Rabs	7
2.6 Localization of Rab proteins	9
2.7 Regulation of intracellular transport by Rab proteins	10
2.8 Regulation of intracellular trafficking in protozoa	16
2.9 Endocytosis and intracellular trafficking in trypanosomatid parasites	17
<b>3. Chapter 1</b>	20
3.1 Introduction	21
3.2 Materials and Methods	22
3.2.1 Materials	22
3.2.2 Antibodies and recombinant proteins	22
3.2.3 Cells	22
3.2.4 Cloning and expression of Rab5 from <i>Leishmania</i>	23
3.2.5 Expression and purification of LdRab5 protein	26
3.2.6 Protein estimation	26
3.2.7 SDS-PAGE and Western blot	26
3.2.8 Preparation of antibodies against LdRab5	27
3.2.9 GTP overlay assay	27
3.2.10 GTPase assay	28
3.2.11 Generation of LdRab5 mutants	28
3.3 Results	29
3.3.1 Cloning of Rab5 homologue from <i>Leishmania donovani</i>	29

3.3.2 Expression and purification of recombinant LdRab5 in <i>E.coli</i>	32
3.3.3 Specificity of anti-LdRab5 antibody	33
3.3.4 Characterization of recombinant LdRab5	34
3.3.5 Generation of LdRab5 mutants	35
3.3.6 Confirmation of the mutations by sequencing	36
3.3.7 Characterization of LdRab5 mutants	37
3.4 Discussion	39
<b>4. Chapter 2</b>	<b>42</b>
4.1 Introduction	43
4.2 Materials and Methods	43
4.2.1 Materials	44
4.2.2 Antibodies	44
4.2.3 Biotinylation of hemoglobin	45
4.2.4 Preparation of early endosomes from <i>Leishmania</i>	45
4.2.5 Subcellular fractionation of <i>Leishmania donovani</i> endocytic compartments	46
4.2.6 <i>In vitro</i> reconstitution of endosome fusion in <i>Leishmania</i>	46
4.2.7 Treatment of endosomes with GDI	47
4.2.8 Immunodepletion of Rab5 from <i>Leishmania</i> cytosol	47
4.2.9 <i>In vitro</i> prenylation recombinant Rab proteins	47
4.2.10 Immunolocalization of Rab5 in <i>Leishmania donovani</i>	48
4.3 Results	48
4.3.1 Subcellular fractionation of <i>Leishmania donovani</i> endocytic compartments	48
4.3.2 <i>In vitro</i> reconstitution of homotypic early endosome fusion	50
4.3.3 Role of NSF-like protein in endosome fusion in <i>Leishmania</i>	52
4.3.4 Role of endocytic Rabs in intracellular trafficking of Hb	53
4.3.5 Immunolocalization of Rab5 in <i>Leishmania donovani</i>	54
4.4 Discussion	55

<b>5. Chapter 3</b>	58
5.1 Introduction	59
5.2 Materials and Methods	60
5.2.1 Materials	60
5.2.2 Preparation of late endosomes from <i>Leishmania</i>	60
5.2.3 <i>In vitro</i> reconstitution of heterotypic endosome fusion in <i>Leishmania</i>	61
5.2.4 Immunodepletion of Rab5 and Rab7 from <i>Leishmania</i> cytosol	62
5.2.5 Treatment of hemoglobin containing early endosomes with anti-hemoglobin receptor antibody	62
5.2.6 Generation of hemoglobin receptor deletion mutants	62
5.2.7 Hemoglobin binding activity of the hemoglobin receptor and its deletion mutants	62
5.3 Results	63
5.3.1 Early and late endosomes in <i>Leishmania</i> differ in their biochemical composition	63
5.3.2 Comparison of heterotypic fusion between endosomes containing receptor-mediated or fluid phase probes	63
5.3.3 Role of Rab proteins in heterotypic fusion	64
5.3.4 Generation of different deletion mutants of hemoglobin receptor and their ability to bind hemoglobin	65
5.3.5 Role of hemoglobin receptor tail in heterotypic fusion	66
5.4 Discussion	67
<b>6. Conclusions</b>	69
<b>7. Bibliography</b>	74