## **ACKNOWLEDGEMENT**

At the very onset my heartfelt thanks and fathomless gratitude goes to my research guide, Dr.R.K. Sarkar, Reader in Agronomy, University College of Agriculture, Calcutta University for his resourceful and helpful guidance and also for constant inspiration during the course of the present investigation.

I am indeed grateful to Prof. B. Bhattacharya, Dean and Dr. M.K. Majumdar, Head, University College of Agriculture, Calcutta University, for extending to me the Laboratory and Library facilities and also for resourceful suggestions during the tenure of my studies.

Since joining the Ramkrishna Ashram Krishi Vigyan Kendra, Nimpith, I have been receiving blessings from Revered Swami Buddhanandaji Maharaj, Founder of Sri Ramkrishna Ashram, Nimpith. The inspiration of this work also comes from him which compels me to express my highest regard and gratitude for him.

Swami Sadananda, Chairman, Ramkrishna Ashram Krishi Vigyan Kendra, Nimpith, has been constantly encouraging and inspiring me to submit the thesis. He has also given me necessary permission to conduct the research work in the KVK Instructional Farm. My sinceremost gratitude and regard goes to him. I am also thankful to Dr. Dipankar Saha, Training Organiser, R.K. Ashram Krishi Vigyan Kendra, for his constant support and help for preparation of this thesis.

I am extremely thankful to my teachers Dr.A.K. Sarkar and Dr. N.N. Bhowmick, both Readers in Agronomy, University College of Agriculture, Calcutta University for their help and suggestions from time to time during my studies.

Special debt is acknowledged to Dr.A.K. Basu, Former Director, Central Institute For Cotton Research ( CICR ), ICAR, Nagpur and Dr. V.T. Sundaramurthy, Former Head and Project Co-ordinator, CICR, Regional Station, Coimbatore, for supplying germplasms, allowing me to use the Institute Libraries at Nagpur and Coimbatore for collection of references and also for their constant suggestions and valuable advice during the course of the investigation. I am

also grateful to Dr.N.B. Patil, Former Director, Central Institute for Research on Cotton Technology (CIRCOT), ICAR, Bombay, for testing the fibre quality of the experimental materials at CIRCOT Laboratory, Bombay.

Dr.Ashis Chakraborty, Lecturer in Agronomy, Bidhan Chandra Krishi Viswavidyalaya and Dr.Anupam Barik, Krishi Vigyan Kendra, Nimpith, deserve special mention for rendering all possible help in preparation of this thesis.

I am proud to have such colleagues and friends who have done a commendable job for me, inspiring me and supporting me extensively in every sphere of my research work. Some of the names which need special mention are, Sri Swapan Kumar Samui, Dr. Bijaykali Mahapatra, Sri Bhaskar Mukherjee, Sri Sunirmal Das, Sri Prasanta Chattopadhyay, Sri Somnath Sengupta and Sri Niladri Dutta. I also desire to extend thanks to the Field Assistants of Agronomy Section of K.V.K., Nimpith, who have helped me a lot during the tenure of the research work.

My acknowledgement would remain incomplete unless I express my regards and thanks to my family members, specially my mother, wife, daughter, niece, brothers and sisters for extending all sorts of help and support right from the beginning in preparing the thesis.

(RATHIS CHANDRA MAJUMDAR)