

Acknowledgments

I am ever grateful and indebted to Dr. K. V. A. Ramaiah, for his excellent guidance, support and invaluable discussions at every stage of this work. Looking back over the years and reminiscing the course of this work, I found that Dr. Ramaiah always gave an overall development to the students. He is simple, generous and humble person who taught students how to develop a positive attitude while working. He always gave simple and elegant experiments. The understanding and consideration he shows towards a student during personal crisis is remarkable. I thank him for all the encouragement provided during the course of this work.

I am thankful to Prof. A. R. Reddy, Dean, School of Life Sciences, for providing me the necessary infrastructure for carrying out my research work.

I thank Prof. K. Subba Rao, FNAMs, Head, Department of Biochemistry, for providing the necessary facilities and a lively atmosphere to carryout my work.

I thank Dr. Ramanadham and Dr. Sivakumar for their keen interest in this work with whom I had, in the course of these investigations, several stimulating discussions.

I thank Prof. I. M. London and Dr. J.-J. Chen, MIT, USA for their gifts of HRI and monoclonal mammalian eIF-2 α antibodies.

I am most indebted to Ms. Janaki, with whom I was associated during every phase of this work. I have always benefited from her suggestions, criticisms and discussions. She was a motivating force and her participation and help is surely the greatest.

I like to particularly thank my laboratory colleagues for their collaboration and discussions on various aspects of this work: Janaki, Naresh, Thanuja, Sudhakar, Prabhat and Neelima.

I thank my friends Bhaskar, Karunasree, Rehman, Rajeev, Ashok, Anand, Venkateshwarulu and others for making my life happier. Special thanks to Dr. Srinivas for givingvaluable suggestions during the thesis preparation.

Above all, I thank my parents, sister and brother-in-law, whose encouragement and support was always forthcoming, but the debt, I will never be able to repay.

I finally acknowledge the University Grants Commission, New Delhi for their financial assistance in the form of Junior and Senior Research Fellowships for my Ph.D study.

Krishna...

1.0 INTRODUCTION	1
1.1 Bird's eye view of eukaryotic protein synthesis.	2
1.1.1 Initiation of protein synthesis.	2
1.1.2 Elongation and termination.	4
1.2 Regulation of initiation of protein synthesis.	5
1.3 eIF-2 structure, function and regulation.	6
1.3.1 Subunit structure and function of eIF-2.	6
1.3.2 Regulation of eIF-2 by phosphorylation.	9
1.3.2.1 Phosphorylation of eIF-2 subunits under various conditions	9
1.3.2.2 eIF-2 alpha kinases.	10
1.3.2.3 Mechanism of protein synthesis inhibition by eIF-2a phosphorylation; importance of eIF-2B activity.	14
1.4 Role of mutants of eIF-2a in understanding the translational regulation.	17
1.5 Regulation of initiation factor-2 activity in plant systems.	19
1.6 OBJECTIVES	23
2.0 MATERIALS AND METHODS	24
2.1 Preparation of heme-deficient reticulocyte lysate.	24
2.2 Measuring reticulocyte lysate protein synthesis.	24
2.3 Preparation of wheat germ lysate.	25
2.4 Wheat germ lysate protein synthesis.	26
2.5 Preparation of wheat germ eIF-2.	26
2.5.1 Assays for eukaryotic initiation factor -2.	29
2.5.1.1 Preparation of wheat germ binary complex, eIF-2.[³ H]GDP, and its dissociation <i>in vitro</i>	29
2.5.1.2 Preparation of [³⁵ S]Met.tRNA.	30
2.5.1.2.1 Formation of eIF-2.Met-tRNA complexes.	30
2.6 Estimation of reticulocyte eIF-2B activity in lysates.	30
2.7 Dissociation of wheat germ.eIF-2.[³ H]GDP, binary complex, in reticulocyte lysates.	31
2.8 Polyclonal wheat germ eIF-2 antibody.	31
2.8.1 Generation of antibodies against wheat germ eIF-2 in New Zealand	32
2.8.1.1 Isolation of control (preimmune) and anti-wheat germ eIF-2 antiserum	32
2.9 Limited proteolytic digestion of wheat germ eIF-2 subunits by Cleveland method.	33
2.9.1 Phosphorylation of purified wheat germ eIF-2 <i>in vitro</i> .	33
2.10 Phosphopeptide mapping in one dimension.	33
2.11 Separation of total ribosomes by sucrose gradient fractionation.	34
2.12 Immunoblot analysis of reticulocyte eIF-2 in the 15S complex of reticulocyte lysates.	35
2.13 Protein phosphorylation in translating reticulocyte lysates.	36
2.14 Protein phosphorylation in translating wheat germ lysates.	36
2.15 Sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE).	36
2.16 Western Blotting.	37
2.17 Protein Estimation.	38
2.18 Autoradiography.	38
2.19 Materials.	38
3.0 GENERAL OUTLINE TO RESULTS AND DISCUSSION	40
3.1 SECTION A: RESULTS AND DISCUSSION	42
3.1.1 Purification of wheat germ eIF-2.	42
3.1.2 Formation of eIF-2.[³ H]GDP and eIF-2.GTP.[³⁵ S]Met.tRNA complexes.	43
3.1.3 Phosphorylation of purified wheat germ eIF-2 by purified kinases <i>in vitro</i> .	44

3.1 4	Cross-reactivity of wheat germ and reticulocyte eIF-2 subunits by polyclonal anti-wheat germ eIF-2 antibodies	45
3.2	SECTION B: RESULTS	47
3.2.1	Wheat germ eIF-2 phosphorylation in reticulocyte lysates	47
3.2.2	Protein synthesis	48
3.2.3	Effect of wheat germ eIF-2 on [eIF-2(aP) eIF-2B) complex formation in reticulocyte lysates	49
3.2 4	eIF-2B activity in inhibited reticulocyte lysate; effect of wheat germ eIF-2	51
3.2.5	Wheat germ eIF-2 dissociation in reticulocyte lysates and in the presence of purified reticulocyte eIF-2(aP)	52
3.2.6	Distribution of reticulocyte and wheat germ eIF-2 in polysomes obtained from inhibited reticulocyte lysates treated with or without wheat germ eIF-2	54
3.2.7	Addition of poly IC (ng-µg/ml) to wheat germ lysates; effect on wheat germ translation, eIF-2 phosphorylation and the GNE activity of eIF-2	56
3.3	SECTION B: DISCUSSION	59
3.4	SECTION C: RESULTS AND DISCUSSION	65
3.4.1	Limited proteolytic digestion of phosphorylated wheat germ eIF-2 subunits	65
3.4.2	Separation of <i>S. aureus</i> protease (SV8) digestion products of phosphorylated p36 and p41-42 subunits of wheat germ eIF-2 in one dimension	66
3.4.2.1	<i>In vitro</i> phosphorylation of purified eIF-2 by CK II and HRI and separation of <i>S. aureus</i> protease (SV8) digestion products	66
3.4.2.2	<i>S. aureus</i> protease (SV8) digestion products of p36 subunit phosphorylated by CK II	66
3.4.2.3	<i>S. aureus</i> protease (SV8) digestion products of p41-42 subunit phosphorylated by CK II or HRI	67
3.4.2.4	Limited proteolytic digestion of the p41-42 subunit phosphorylated by CK II or HRI and a combination of both the enzymes (Fig. 22)	67
3.4.2.5	Comparison of the phosphopeptides of reticulocyte eIF-2α and the p41-42 subunit of wheat germ generated by higher concentrations of <i>S. aureus</i> (SV8) protease	68
4.0	SUMMARY	71
5.0	REFERENCES	76