

## LIST OF FIGURES

- Fig. 1a and 1b.** Reticulocyte lysate protein synthesis.
- Fig. 2a and 2b. Wheat germ lysate protein synthesis.
- Fig. 3.** Schema for the wheat germ **eIF-2** purification.
- Fig. 4a.** **Coomassie** stained gel of different **fractions** that were obtained **during** the purification of wheat germ eIF-2.
- Fig. 4b. Carboxy Methyl Cellulose Sephadex-C 50 (CM Sephadex) purified wheat germ eIF-2.
- Fig. 4c. Purified wheat germ and reticulocyte eIF-2 **fractions** separated on 10% **SDS-PAGE** for subunit comparison.
- Fig. 5a. Phosphorylation of reticulocyte and wheat germ eIF-2 *in vitro* by **HRI**.
- Fig. 5b. Phosphorylation of reticulocyte and wheat germ eIF-2 *in vitro* by **dsI** kinase.
- Fig. 5c.** Phosphorylation of purified wheat germ eIF-2 *in vitro* by purified CK II and partially purified HRI.
- Fig. 6a. Specificity of wheat germ eIF-2 polyclonal antibodies.
- Fig. 6b. Cross-reactivity of wheat germ eIF-2 polyclonal antibodies.
- Fig. 7. [<sup>32</sup>P]-Labeled phosphoprotein profiles of heme-deficient lysates, or **heme** (20 **μM**) and poly **IC** (75 **ng/ml**) treated lysates with or without wheat germ eIF-2 (2 **μg**).
- Fig. 8. Wheat germ eIF-2 protects protein synthesis in heme-deficient (Fig. 8a), heme and poly **IC-treated** (Fig. 8b) reticulocyte lysate.
- Fig. 9. Wheat germ eIF-2 does not mitigate the protein synthesis inhibition in reticulocyte lysates caused by **cycloheximide**.
- Fig. 10.** Immunoblot analysis of free eIF-2 and eIF-2 in the **15S** complex in 10-30% sucrose gradient fractions of normal and inhibited reticulocyte lysates.
- Fig. 11. Western immunoblot analysis of 15S complex fraction obtained from reticulocyte lysates supplemented with or without wheat germ eIF-2 to determine the interaction between phosphorylated wheat germ eIF-2 and reticulocyte eIF-2B.
- Fig. 12. Kinetics of eIF-2B activity in **hemin** and poly **IC-treated** reticulocyte lysates in the presence and absence of wheat germ eIF-2.
- Fig. 13. Guanine nucleotide exchange on wheat germ eIF-2 in reticulocyte lysates.
- Fig. **14.** Effect of phosphorylation of wheat germ eIF-2 by HRI *in vitro* on the dissociation of wheat germ **eIF-2**·[<sup>3</sup>H]GDP binary complex.

- Fig. 15. Polysome profiles of reticulocyte lysates treated with heme, heme and poly IC and heme, poly IC and wheat germ eIF-2**
- Fig. 16.** Western immunoblot analysis of the 10-50% sucrose gradient fractions of reticulocyte lysates supplemented with or without wheat germ eIF-2; detection of the presence of wheat germ and reticulocyte eIF-2 in the polysome fractions
- Fig. 17.** Wheat germ lysate protein synthesis in the presence of varying concentrations of poly IC.
- Fig. 18.** [<sup>32</sup>P] phosphoprotein profiles of wheat germ lysates in the presence of different concentrations of poly IC and wheat germ eIF-2
- Fig. 19.** Effect of poly IC-treated wheat germ lysate on wheat germ eIF-2.[<sup>3</sup>H]GDP complex dissociation
- Fig. 20.** Phosphorylation of wheat germ eIF-2 subunits by HRI and CK II *in vitro* for phosphopeptide analysis.
- Fig. 21.** Cleland partial peptide digestion of phosphorylated wheat germ eIF-2 subunits
- Fig. 22.** SV8 protease digestion products of p41-42 subunit of wheat germ eIF-2 phosphorylated by HRI, CK II and HRI + CK II added together or at different time points.
- Fig. 23.** Cleland mapping of phosphorylated p41-42 subunit of wheat germ eIF-2 and alpha-subunit of reticulocyte eIF-2 in the presence of SV8 protease enzyme; effect of two concentrations of the protease