

REFERENCES

- Anonymous 1973. The New Encyclopaedia Britannica. Vol. VII, Encyclopaedia Britannica Inc. 15th Edition, Helen Hemingway Benton Publisher: 1973-74: 133.
- Anonymous 1975. Statistical Abstract. Bureau of Applied Economics and Statistics, Government of West Bengal (India).
- Al-Jibouri, H.A. Miller P.A. and Robinson, H.F. 1958. Genotypic and environmental variance and covariances in an upland cotton cross of interspecific origin. Agron. J., 50: 633-637.
- Arunachalam, V. and Ram Jawhar 1967. Geographic Diversity in relation to genetic divergence. Ind. J. Genet., 27 : 369-80.
- * Babdzhanov, F.A. and A. Mauturdyev, B. 1971. (Correlated relationships between staple length and other economic characters of hybrid cotton) Nauch tr Tashkent-S-kh in-t., 22: 94-100 (Rus) from Referativnyi zhurnal. 1972. 2155* 295.
- Balakataiah, K. 1973. Path coefficient analysis in upland cotton (G. hirsutum L.). Ind. J. Agric. Sci. 43(7): 681-683.
- Balakataiah, K. and Butany, W.T. 1971. Correlated response to selection for boll number in upland cotton (G. hirsutum L.). Ind. J. Agric. Sci. 41(11): 920-924.

- Baluch, M.A.A. 1963. A note on selection index technique for yield improvement in cotton. *Pakist. Cotton.* 7: 133-136.
- Bandyopadhyaya, S.B. 1964. Assessment of Jute fibre- Bundle strength-I. *Jute Bulletin.* 27(8): 1-10.
- Bandyopadhyaya, S.B. 1965. Assessment of Jute fibre- Bundle Strength-II. *Jute Bulletin.* 27(11): 1-8.
- Bandyopadhyaya, S.B. and Chatterjee, U. 1960. Assessment of jute fibre. *Textile Trends.* 5(6): 260-264.
- Barkat, M.A., Fachry, S.I. and Kahil, M.A. 1971. Relative salt tolerance in 5 varieties of Egyptian cotton. *Agril. Res. Review. Arab Republic of Egypt.* 49(2): 191-200.
- Bhaumick, P.K. 1971. D.Phil. Thesis. Calcutta University.
- Burton, S. and Devane, E.H. 1953. Estimating heritability in tall Fescue (*Festuca arundinacea*) from replicated clonal material. *Agron. J.* 45 : 478-481.
- Chakravarty, A.C. 1973. Fibre characters involved in raw jute grading and their estimation. *Textile Trends.* 59: 1-9.
- Chandranath, P.S. 1973. Correlations and Path coefficient of ginning percentage in cotton, Madras. *Agric. Jour.* 60: 1526-1528.

Comstock, R.E. and Robinson, H.F. 1952.

**Estimation of average dominance of genes
in Heterosis . Iowa State College Press.
Ames. Iowa. , 494 - 516.**

Dewey, D.R. and Lu, K.H. 1959. A correlation and path coefficient analysis of crested wheat grass seed production. Agron. J. 51 : 515-518.

*El Latif, H.M.A. and EL Mazar, M.F. 1970. Variability and correlation studies in some families of Menoufi variety of Egyptian cotton. Agril. Res. Review, Arab Republic Egypt. 48(6): 63-70.

Federer, W.T. 1955. Experimental Design, Theory and Application. The Macmillan Company, London. 283-287.

Fisher, R.A. 1918. The correlations among relatives on the supposition of Mendelian inheritance. Trans. Royal Soc. of Edinburgh L.II : 399-433.

Fisher, R.A. 1954. The discrimination of groups by means of multiple measurements. Statistical Methods, Oliver & Boyd. 285-292.

Gupta, S.P., Singh, T.H. and Phul, P.S. 1972. Analysis of quantitative variability in upland cotton. Ind. J. Agric. Sci. 42(3): 215-218.

Hutchinson, J. 1937. The Wealth of India, Raw Material (1956) Vol. IV : F-G. Council of Scientific and Industrial Research, New Delhi.

*Johannsen, W. 1909. Elemente der exakten Erblchkeitslehre. Jena.
Gustav. Fischer.

Johnson, H.W., Robinson, H.F. and Comstock, R.E. 1955. Genotypic
and phenotypic correlations in soybeans and
their implications in selection. Agron. J.
47(10) : 477-483.

Kamalanathan, S. and Ponnaiya, B.W.X. 1964. Construction of selection
index for lint yield in cotton (G. arboreum L.)
Madras Agric. J. 51: 358.

Khan, Siraj-Ud-Din, M. 1963. Stigma length an index of fibre
length in Pak cottons. W. Pak. J. Agric. Res.
1(2): 110-115.

Kirby, R.H. 1952. Vegetable fibres. Leonard Hill (Books) Limited,
London.

Kundu, B.C. 1942. The anatomy of two Indian fibre plants Cannabis
and Corchorus with special reference to fibre
distribution and development. J. Indian Bot.
21 : 93-128.

Mehridirtta, P.D., Phul, P.S. and Arora, D.N. 1971. Genetic diversity
in relation to fodder yield and its component in
sorghum. Ind. J. Genet. 31(2): 300.

Murty, B.R. and Arunachalam, V. 1966. The nature of divergence in
relation to Breeding system in some crop plants.
Ind. J. Genet. 26A : 188-198.

- *Nagubin, Ya. D. and Suvlatov, S. 1972. (The inheritance of weight of crude cotton per boll in hybrids of species G. hirsutum) . Dozl Acad. Fanhoi RSS Tojikistan 15(1): 46-49 (from Referativnyi zhurnal (1972): 7 : 55 : 139).
- Patil, M.S. and Mathapati, S.N. 1976. Studies on genetic variability for yield and its components in cotton (G. hirsutum). Mysore Journal of Agric.Sci. 10(1): 17-23.
- Patil, M.S. and Mensinkai, S.W. 1972. Selection index for yield in diploid cotton (G. arboreum) . Madras Agri. Jour. 59(11): 660-662.
- *Pudenko, L.S. 1968. (Correlations between some economically valuable characters of cotton). In materialy pyatoi ob e dinennoi konfesentsii Molody kh Uchenykh posel skma khoyaistva Uzbekistan (Selektsii genetika Semenovodstvo Kholopchatnikai Lyutserroy Tashkent, Uzbek SSR, Fan: 162-166.
- Rao, C.R. 1952. Advanced statistical method for Biometrical Research. John Willey & Sons, Inc., New York.
- Ramanujam, S. and Rai, B. 1963. Analysis of yield components in Brassica campestris var yellow sarson. Ind. J. Genet. 23 : 312-19.
- *Savov, P. 1962. Results of the crossing of geographically distant cotton varieties. Izv Centr Nauczisled Inst hidroth. Abst. Bulg.Sci. lit. 7 : Abst 80 : 20.

- Santhanam, V. 1967. Breeding Procedures for cotton. I.C.A.R. Tech. Bull. (Agri.) Series No.10.I.C.A.R. 6.
- Singh, B.B. 1973. Nature of variation and association between characters related to yield and quality in G. hirsutum L. Madras Agric. J. 60(9): 1882-1884.
- Singh, R.B. and Bains, S.S. 1968. Genetic divergence of upland cotton. Ind. J. Genet. 28 : 151-157.
- Singh, H.G., Govil, S.K. and Chaturvedi, M.C. 1971. Correlations and heritability of some characters in cotton (G. arboreum L.). Allahabad Fmr. 45(5): 515-517.
- Singh, R.B. and Gupta, M.P. 1968. Multivariate analysis of divergence of upland cotton. Ind. J. Genet. 28: 151-157.
- Sivasubramanian, V. 1962. Selection of suitable indices for determining of earliness of cotton varieties under rice fallows condition. Madras Agric. J. 4: 117-120.
- Ter-Avanesyan, D.V. 1967. Variability of the hybrid generation of cotton plants obtained through crossing geographically removed specimens by the use of the limited pollination method. Ind. J. Agric. Sci. 37: 120-127.
- Tukey, J.W. 1954. Causation, regression and path analysis. Chapter 3 of Statistics and Mathematics in biology edited by O. Kempthorne et al. Iowa State College Press, Ames.

- *Vadogreeva, L.G. 1975. (Signed of earliness of fibred cotton)
Khbopkovodstvo No.1-32(Ru)(Taken from Plant
Breeding Abst. Commonwealth Bureaux of Agri.
45(7): 5610).
- Venkata Raman, N. 1963. Formulation of selection index for yield
in combodia cotton (G. hirsutum L.) correlation
study of yield components - a review. Madras
Agric. J. 50: 179-182.
- Venkata Raman, N. and Santhanam, V. 1962. Correlation studies on
yield components in combodia cotton (G.hirsutum L.)
Ind. Cott. Gr. Rev. 16: 131-137.
- Wright, S. 1921. Correlation and Causation. J. Agric. Res.
20: 557-585.
- Wright, S. 1934. The method of path coefficient. Am. Math. Statist.
5: 161-215.

* Original not seen.