

CONTENTS

	Page No.	
CHAPTER 1 : <u>GENERAL INTRODUCTION</u>		
1.0	Introduction	1
1.1	Theory of free radical vinyl polymerization	1
1.2	Kinetic chain length and degree of polymerization	5
1.3	k_p^2/k_t ratio	6
1.4	Chain transfer	7
1.5	Inhibition and retardation	8
1.6	Photopolymerization	8
1.7	Energetic characteristics (activation energy)	11
1.8	Non-ideality in the kinetics of radical vinyl polymerization	12
CHAPTER 2 : <u>LITERATURE REVIEW</u>		
2.0	Introduction	14
2.1	Dye-sensitized polymerization	14
2.1.1	Polymerization in the presence of oxygen	15
2.1.2	Polymerization in the absence of oxygen	17
2.2	Peroxide-amine combinations as initiators for vinyl polymerization	21
2.3	Vinyl polymerization induced by Michler's ketone under photo-condition	25
2.4	Main theme of the present thesis	28

CHAPTER 3 : EXPERIMENTAL (GENERAL)

3.0	Purification of monomer - methyl methacrylate (MMA)	30
3.1	Purification of solvents	30
3.2	Purification of initiator components	30
3.2.1	Benzoyl peroxide (Bz_2O_2)	30
3.2.2	Dyes or dye intermediates	30
3.3	Purification of nitrogen gas	32
3.4	Dilatometric technique of polymerization	32
3.5	Equipments and their descriptions for dilatometric measurement of polymerization	32
3.5.1	Dilatometer	32
3.5.2	Thermostatic bath	33
3.5.3	Light source	33
3.6	Experimental procedure to follow the course of polymerization of MMA taken in the dilatometer	34
3.6.1	Procedure	34
3.6.2	Calculation	34
3.6.3	Rate of polymerization of MMA in the diluted system	36
3.7	Isolation and purification of polymer	36
3.8	Determination of polymer molecular weight from intrinsic viscosity	36
3.9	Spectrophotometric studies of initiator systems	38

3.10	Spectrophotometric analysis of polymers obtained	39
------	--	----

CHAPTER 4 : RESULTS AND DISCUSSIONS

4.1	Photopolymerization of MMA at 40°C using Rhodamine 6G anhydrobase (Rh) and benzoyl peroxide combination as photoinitiator	40
4.1.0	Introduction	40
4.1.1	Spectrophotometric studies of Rhodamine 6G (anhydrobase) - Bz ₂ O ₂ system	41
4.1.2	Initiator exponent	41
4.1.3	k_p^2/k_t value	42
4.1.4	Initiator transfer	44
4.1.5	Activation Energy	46
4.1.6	Photopolymerization of MMA in diluted systems	47
4.1.7	End-Group analysis	47
4.1.8	Mechanism	49
4.1.9	Termination	49
4.2	Photopolymerization of MMA at 40°C using Auramine O anhydrobase (Au) and benzoyl peroxide combination as photoinitiator	57
4.2.0	Introduction	57
4.2.1	Spectrophotometric studies of Auramine-O anhydrobase (Au)-Bz ₂ O ₂ system	58
4.2.2	Initiator Exponent	58
4.2.3	k_p^2/k_t value	59
4.2.4	Initiator transfer	59

4.2.5	Activation energy	61
4.2.6	Photopolymerization of MMA in diluted systems	63
4.2.7	End-group analysis	65
4.2.8	Mechanism	65
4.2.9	Termination	66
4.3	Photopolymerization of MMA at 40°C using Michler's ketone (MK) and benzoyl peroxide combination as photoinitiator	72
4.3.0	Introduction	72
4.3.1	Spectrophotometric analysis of Michler's ketone (MK)-Bz ₂ O ₂ system	73
4.3.2	Initiator exponent	73
4.3.3	k_p^2/k_t value	74
4.3.4	Activation energy	74
4.3.5	Photopolymerization of MMA in diluted systems	76
4.3.6	Examination of solvent dependency of the radical generation/initiator process	78
4.3.7	End group analysis	79
4.3.8	Mechanism	79
4.3.9	Termination	82
4.4	Thermal polymerization of MMA at 40°C using crystal violet lactone (CVL) and benzoyl peroxide combination as photoinitiator	87
4.4.0	Introduction	87
4.4.1	Spectrophotometric analysis of crystal violet lactone (CVL)-Bz ₂ O ₂ initiator system	88

	Page No.
4.4.2 Initiator exponent	88
4.4.3 k_p^2/k_t value	90
4.4.4 Activation energy	90
4.4.5 Initiator transfer	91
4.4.6 Thermal polymerization of MMA in diluted systems	92
4.4.7 Examination of solvent depend- ing of the radical generation /initiation process	94
4.4.8 End-group analysis	96
4.4.9 Mechanism	96
4.4.10 Termination	97
 SUMMARY	 102
 REFERENCE	 114