

## LIST OF TABLES

1). Table 3.3-T1 : Binding energies (BE) of nuclei in sd shell with universal sd interaction by spectral distribution methods along with experimental values and those by shell model. Column  $\bar{A}$  represents BE obtained using the simple Ratcliff procedure, column  $\bar{B}$  contains the BE corrected by integration up to an excited state, and column  $\bar{C}$ , in addition, incorporates the  $(\gamma_1, \gamma_2)$  correction as given by equation (B-8) of appendix B.

page 35

2). Table 3.3-T2 : Ground state occupancies of  $1d_{5/2}$ ,  $2s_{1/2}$ , and  $1d_{3/2}$  orbits as predicted by shell model and by spectral distribution methods using the universal sd interaction.

page 38

3). Tables 4.1-T1(a,b,c,d) : One -body matrix elements  $\epsilon^{11}(rs)$  for the Gamow-Teller and isovector M1 operators for sd and fp shell.

page 45

4). Table 4.2-T1 : Strength sums for  $\beta^-$ -decays of some even-even fp shell nuclei calculated using approximate sum rule expression of eq.(4.2-8a) with occupancies obtained by three different methods.

page 50

5). Table 4.3-T1 : Centroids and widths of H and K (the sum rule operator) along with their correlation coefficient  $\zeta_{HK}$  and the expansion of K in terms of polynomials in energy upto the second term in the scalar-T space (see eq.(4.3-2)) for even-even self-conjugate fp shell nuclei. The ground state binding energy used here is taken from Haq et al.

page 53

6). Table 4.3-T2 : FP shell Gamow-Teller  $\beta^-$  strength sums from the ground state with the final isospin ( $T'$ ) decomposition for a number of even and odd A neutron-rich Fe and Ni isotopes, evaluated in scalar-T space using both R- and S- strength methods. For R-strength sums we present the CLT result (i.e.  $K_0 + K_1(E) = K_{CLT}(E)$ ). Here  $K_0$  and  $K_1$  are successive terms in the R-strength expansion of the sum rule strength. The three values of final isospin are  $T-1 = T_<$ ,  $T$ , and  $T+1 = T_>$ . The  $\beta^+$  strength sum can be obtained from the  $\beta^-$  strength sum using the Ikeda constraints  $S_{\beta^-} - S_{\beta^+} = 3(N-Z)$ .

page 57

6). Table 4.3-T3 : The expression for the density operators  $\rho_{rs}^\nu(x, \tau)$  for  $x \leq 2$ .

page 58

7). Table 4.3-T4 : Configuration input coefficients  $C_{rs}^{\omega_T}(x, \tau)$  for GT operator in fp shell.

page 59

8). Table 4.3-5a : Configuration averaged  $\beta^-$  strength sum for some Fe and Ni isotopes with decomposition according to the final isospin along with the available shell model values.

page 60

9). Table 4.3-T5b : Scalar and configuration averaged  $\beta^+$  strength sum for Fe and Ni isotopes along with the shell model results.

page 60

10). Table 4.3-T6 : Correlation coefficient  $\zeta_{HK}$  of the Hamiltonian with  $K = O^+O$  where  $O$  is the excitation operator for both GT and  $M1^{\Delta T=1}$  excitations for five self-conjugate nuclei in the sd shell.

page 66

11). Table 4.3-T7a :  $\beta^-$  -strength sums from the ground state to all final states with isospin  $T' = T-1 = T_<$ ,  $T' = T$ , and  $T' = T+1 = T_>$  for some nuclei in sd shell . The strength sum is evaluated by using both R- and S-strength methods. For the R-strength the table gives the strength sums calculated including the first term beyond the CLT term (i.e.  $K_2(E)$ ) for which correction due to non-zero skewness and excess ( $\gamma_1, \gamma_2$ ) is also made. The numbers within the parenthesis are the shell model sum rule strengths obtained by Hino et al.

page 70-71

12). Table 4.3-T7b : Isovector M1 sum rule strengths for some sd shell nuclei with final isospin decomposition (in this  $T' = T, T+1$  only). We display  $M_0(E)$  calculated by both R- and S strength methods. For the R-strength method the term  $K_2(E)$  is calculated for both zero and non-zero values of  $\gamma_1$  and  $\gamma_2$ . The numbers within the parenthesis are the values obtained by the shell model calculation of Hino et al.

page 72-73

13). Table 5.2-T1 : The marginal strength centroids and strength widths along with the strength correlation coefficient (defined by equation (2.4-5a) and (2.4-5b) respectively) for five self-conjugate sd shell nuclei, with GT and  $M1^{\Delta T=1}$  excitation operators.

page 80

14). Table 6.3-T1 : Comparisons of calculated and experimental halflives.

page 97

15). Table 6.3-T2 :  $\beta^-$  decay rates for the nuclei  $^{69}\text{Cu}$ ,  $^{67}\text{Ni}$ ,  $^{63}\text{Co}$  and  $^{62}\text{Co}$ . The number of excited states known experimentally for  $^{69}\text{Cu}$ ,  $^{63}\text{Co}$  and  $^{62}\text{Co}$  are 7, 12, and 25 respectively, whereas for  $^{67}\text{Ni}$ , no excited state is available. The numbers within the parenthesis are the rates from the ground state only.  $Y_e = 0.50$ .

page 98-100