

REFERENCES

-
- Abdallah KS, Cao Y, Wei DJ. 2011. Epidemiologic investigation of extraintestinal pathogenic *E. coli* (ExPEC) based on PCR phylogenetic group and fimH single nucleotide polymorphisms (SNPs) in China. *Int J Mol Epidemiol Genet* 2(4):339-53.
 - Al-Hasani K, Navarro-Garcia F, Huerta J, Sakellaris H, Adler B. 2009. The immunogenic SigA enterotoxin of *Shigella flexneri* 2a binds to HEp-2 cells and induces fodrin redistribution in intoxicated epithelial cells. *PLoS One* 4(12):e8223.
 - AL-Hasani K, Rajakumar KK, Bulach D, Robins-Browne R, Adler B, Sakellaris H. 2001. Genetic organization of the *she* pathogenicity island in *Shigella flexneri* 2a. *Microb Pathog* 30(1):1-8.
 - Al-Hasani K, Henderson IR, Sakellaris H, Rajkumar K, Grant T, Nataro JP et al., 2000. The *sigA* gene which is borne on the *she* pathogenicity island of *Shigella flexneri* 2a encodes an exported cytopathic protease involved in intestinal fluid accumulation. *Infect Immun* 68(5):2457-63.
 - Allen PM, Roberts I, Boulnois GJ, Saunders JR, Hart CA. 1987. Contribution of capsular polysaccharide and surface properties to virulence of *Escherichia coli* K1. *Infect Immun* 55(1):2662–8.
 - Alos JI, Lambert T, Courvalin P. 1993. Comparison of two molecular methods for tracing nosocomial transmission of *Escherichia coli* K1 in a neonatal unit. *J Clin Microbiol* 31(7):1704–09.
 - Baltimore RS, Huie SM, Meek JI, Schuchat A, O’Brein KL. 2001. Early-onset neonatal sepsis in the era of group B Streptococcal prevention. *Pediatrics* 108(5):1094-8.
 - Bassani DG, Kumar R, Awasthi S, Morris SK, Paul VK, Shet A, et al., 2010. Causes of neonatal and child mortality in India: a nationally representative mortality survey. *Lancet* 376(9755):1853-60.
 - Basu S. 2015. Neonatal sepsis: the gut connection. *Eur J Clin Microbiol Infect Dis* 34(2):215-22.
 - Belperio JA, Keane MP, Arenberg DA, Addison CL, Ehlert JE, Burdick MD et al. 2000. CXC chemokines in angiogenesis. *J Leukoc Biol* 68(1):1-8.

-
- Benitz WE. 2010. Adjunct laboratory tests in the diagnosis of early-onset neonatal sepsis. *Clin Perinatal* 37(2):421-38.
 - Benjelloun-Touimi Z, Sansonetti PJ, Parsot C. 1995. SepA, the major extracellular protein of *Shigella flexneri*: autonomous secretion and involvement in tissue invasion. *Mol Microbiol* 17(1):123-35.
 - Benjelloun-Touimi Z, Si Tahar M, Montecucco C, Sansonetti PJ, Parsot C. 1998. SepA, the 110 kDa protein secreted by *Shigella flexneri*: two-domain structure and proteolytic activity. *Microbiology* 144(pt 7):1815–22.
 - Bernstein HD. 2007. Are bacterial “autotransporters” really transporters? *Trends Microbiol* 15(10):441–7.
 - Bhakdi S, Mackman N, Nicaud JM, Holland IB. 1986. *Escherichia coli* hemolysin may damage target cell membranes by generating transmembrane pores. *Infect Immun* 52(1):63–9.
 - Bhandari V, Wang C, Rinder C, Rinder H. 2008. Hematologic profile of sepsis in neonates: neutrophil CD64 as a diagnostic marker. *Pediatrics* 121(1):129-34.
 - Bhat YR, Lewis LE, Vandana KE. 2011. Bacterial isolates of early-onset neonatal sepsis and their antibiotic susceptibility pattern between 1998 and 2004: an audit from a center in India. *Ital J Pediatr* 37:32.
 - Bianconi S, Gudavalli M, Sutija VG, Lopez AL, Barillas-Arias L, Ron N. 2007. Ranitidine and late-onset sepsis in the neonatal intensive care unit. *J Perinat Med* 35(2):147–50.
 - Bien J, Sokolova O, Bozko P. 2012. Role of Uropathogenic *Escherichia coli* Virulence Factors in Development of Urinary Tract Infection and Kidney Damage. *Int J Nephrol* Article ID 681473, 15 pages doi:10.1155/2012/681473.
 - Bingen E, Picard B, Brahim N, Mathy S, Desjardins P, Elion J, et al. 1998. Phylogenetic analysis of *Escherichia coli* strains causing neonatal meningitis suggests horizontal gene transfer from a predominant pool of highly virulent B2 group strains. *J Infect Dis* 177(3):642-50.

-
- Bizzarro MJ, Raskind C, Baltimore RS, Gallagher PG. 2005. Seventy-five years of neonatal sepsis at Yale:1928-2003. *Pediatrics* 116(3):595-602.
 - Boisen N, Perez FR, Scheutz F, Krogfelt KA, Nataro JP. 2009. Short Report: High prevalence of serine protease autotransporter cytotoxins among strains of enteroaggregative *Escherichia coli*. *Am J Trop Med Hyg* 80(2):294-301.
 - Bonadio WA, Hegenbarth M, Zachariason M. 1990. Correlating reported fever in young infants with subsequent temperature patterns and rate of serious bacterial infections. *Pediatr Infect Dis J* 9(3):158.
 - Borden EC and Chin P. 1994. Interleukin-6: a cytokine with potential diagnostic and therapeutic roles. *J Lab Clin Med* 123(6):824-9.
 - Bradford MM. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72:248-54.
 - Bruce MC, Poncz L, Klinger JD, Stern RC, Tomashefski JF, Dearborn DG. 1985. Biochemical and pathologic evidence for proteolytic destruction of lung connective tissue in cystic fibrosis. *Am Rev Respir Dis* 132(3):529-35.
 - Brunder W, Schmidt H, Karch H. 1997. EspP, a novel extracellular serine protease of enterohaemorrhagic *Escherichia coli* O157:H7 cleaves human coagulation factor V. *Mol Microbiol* 24(4):767-78.
 - Buttery JP. 2002. Blood cultures in newborns and children: optimising an everyday test. *Archiv Dis Childhood*. 87(1):F25-8.
 - Camacho-Gonzalez A, Spearman PW, Stoll BJ. 2013. Neonatal infectious diseases evaluation of neonatal sepsis. *Pediatr Clin N Am* 60(2):367-89.
 - Cappello RE, Estrada-Gutierrez G, Irls C, Giono-Cerezo S, Bloch RJ, Nataro JP. 2011. Effects of the plasmid-encoded toxin of enteroaggregative *Escherichia coli* on focal adhesion complexes. *FEMS Immunol Med Microbiol* 61(3):301-14.
 - Cavalieri SJ, Bohach GA, Snyder IS. 1984. *Escherichia coli* α -hemolysin: characteristics and probable role in pathogenicity. *Microbiol Rev* 48(4):326-43.

-
- Cetinkaya M, Ozkan H, Köksal N, Celebi S, Hacimustafaoğlu M. 2009. Comparison of serum amyloid A concentrations with those of C-reactive protein and procalcitonin in diagnosis and follow-up of neonatal sepsis in premature infants. *J Perinatol* 29(3):225-31.
 - Chacko B, Sohi I. 2005. Early onset neonatal sepsis. *Indian J Pediatr* 72(1):23-6.
 - Chapman TA, Wu XY, Barchia I, Bettelheim K, Driesen S, Trott DJ et al. 2006. Comparison of virulence gene profiles of *Escherichia coli* strains isolated from healthy and diarrheic swine. *Appl Environ Microbiol* 72(7):4782-95.
 - Chen M, Tofighi R, Bao W, Aspevall O, Jahnulainen T, Gustafsson LE, et al. 2006. Carbon monoxide prevents apoptosis induced by uropathogenic *Escherichia coli* toxins. *Pediatr Nephrol* 21(3):382–9.
 - Chiesa C, Panero A, Osborn JF, Simonetti AF, Pacifico L. 2004. Diagnosis of neonatal sepsis: a clinical and laboratory challenge. *Clin Chem* 50(2):279-87.
 - Clermont O, Banacorsi S, Bingen E. 2000. Rapid and simple determination of the *Escherichia coli* phylogenetic group. *Appl Environ Microbiol* 66(10):4555-8.
 - Cohen J. 2002. The immunopathogenesis of sepsis. *Nature* 420(6917):885–91.
 - Cooperstock MS. 1974. Inactivation of endotoxin by polymyxin B. *Antimicrob Agents Chemother* 6(4):422-5.
 - Corrigan JJ Jr, Bell BM. 1971. Endotoxin-induced intravascular coagulation: prevention with polymyxin B sulfate. *J Lab Clin Med* 77(5):802-10.
 - Couto RC, Carvalho EA, Pedrosa TM, Pedroso ER, Neto MC, Biscione FM. 2007. A 10-year prospective surveillance of nosocomial infections in neonatal intensive care units. *Am J Infect Control* 35(3):183-9.
 - Darmstadt GL, Bhutta ZA, Cousens S, Adam T, Walker N, de Bernis L, et al. 2005. Evidence based, cost-effective interventions: how many newborn babies can we save? *Lancet* 365(9463):977–88.

-
- Das P, Singh AK, Mukherjee S, Rajendran K, Saha DR, Koley H, et al. 2013. Composition of *Escherichia coli* population in the neonatal gut: phylogroups and virulence determinants. *J Med Microbiol* 62(Pt 11):1680-7.
 - Dautin N, Barnard TJ, Anderson DE, and Bernstein HD. 2007a. Cleavage of a bacterial autotransporter by an evolutionarily convergent autocatalytic mechanism. *EMBO J* 26(7):1942-52.
 - Dautin N, Bernstein HD. 2007b. Protein secretion in Gram-negative bacteria via the autotransporter pathway. *Annu Rev Microbiol* 61:89-112.
 - Dautin N. 2010. Serine Protease Autotransporters of *Enterobacteriaceae* (SPATEs): Biogenesis and function. *Toxins* 2(6):1179-1206.
 - De Felice C, Flori ML, Pellegrino M, Toti P, Stanghellini E, Molinu A, et al. 2002. Predictive value of skin color for illness severity in the high risk newborn. *Pediatr Res* 51(1):100-5.
 - De Lorenzo V, Martinez JL. 1988. Aerobactin production as a virulence factor: a reevaluation. *Eur J Clin Microbiol Infect Dis* 7(5):621-9.
 - De Lorenzo V, Neilands JB. 1986. Characterization of *iucA* and *iucC* genes of the aerobactin system of plasmid ColV-K30 in *Escherichia coli*. *J Bacteriol* 167(1):350-5.
 - De Souza HL, De Carvalho VR, Romeiro FG, Sasaki LY, Keller R, Rodrigues J. 2012. Mucosa-associated but not luminal *Escherichia coli* is augmented in Crohn's disease and ulcerative colitis. *Gut Pathog* 4(1):21.
 - DeForge LE, Remick DG. 1991. Kinetics of TNF, IL-6, and IL-8 gene expression in LPS-stimulated human whole blood. *Biochem Biophys Res Commun* 174(1):18-24.
 - Diard M, Garry L, Selva M, Mosser T, Denamur E, Matic I. 2010. Pathogenicity associated islands in extraintestinal pathogenic *Escherichia coli* are fitness elements involved in intestinal colonization. *J Bacteriol* 192(19):4885-93.
 - Dinarello CA. 1997. Proinflammatory and anti-inflammatory cytokines as mediators in the pathogenesis of septic shock. *Chest* 112(6):321S-9S.

-
- Djafari S, Ebel F, Deibel C, Kramer S, Hudel M, Chakraborty T. 1997. Characterization of an exported protease from Shiga toxin-producing *Escherichia coli*. *Mol Microbiol* 25(4):771-84.
 - Dobrindt U, Chowdary MG, Krumbholz G, Hacker J. 2010. Genome dynamics and its impact on evolution of *Escherichia coli*. *Med Microbiol Immunol* 199(3):145–54.
 - Dolly JO, Aoki KR. 2006. The structure and mode of action of different botulinum toxins. *Eur J Neurol* 13:1-9.
 - Dombrovskiy VY, Martin AA, Sunderram J, Paz HL. 2007. Rapid increase in hospitalization and mortality rates for severe sepsis in the United States: a trend analysis from 1993 to 2003. *Crit Care Med* 35(5):1244–50.
 - Dulay AT, Buhimschi IA, Zhao G, Luo G, Abdel-Razeq S, Cackovic M, et al. 2008. Nucleated red blood cells are a direct response to mediators of inflammation in newborns with early-onset neonatal sepsis. *Am J Obstet Gynecol* 198(4):426 e1- e9.
 - Duriez P, Clermont O, Bonacorsi S, Bingen E, Chaventre A, Elion J, et al. 2001. Commensal *Escherichia coli* isolates are phylogenetically distributed among geographically distinct human populations. *Microbiology* 147(Pt 6):1671-6.
 - Dutta PR, Cappello R, Garcia FN, Nataro JP. 2002. Functional comparison of serine protease autotransporters of *Enterobacteriaceae*. *Infect Immun* 70(12):7105-13.
 - Eberspacher B, Hugo F, Bhakdi S. 1989. Quantitative study of the binding and hemolytic efficiency of *Escherichia coli* hemolysin. *Infect Immun* 57(3):983–8.
 - Edmond K, Zaidi A. 2010. New approaches to preventing, diagnosing and treating neonatal sepsis. *PLoS Med* 7(3):e1000213.
 - Eslava C, Navarro-Garcia F, Czeczulin JR, Henderson IR, Cravioto A, Nataro JP. 1998. Pet, an autotransporter enterotoxin from enteroaggregative *Escherichia coli*. *Infect Immun* 66(7):3155-63.

-
- Etienne-Manneville S, Hall A. 2002. Rho GTPases in cell biology. *Nature* 420(6916):629–35.
 - Fahlman C, Jacobsen FW, Veiby OP, McNiece IK, Blomhoff HK, and Jacobsen SEW. 1994. Tumor necrosis factor- α (TNF- α) potently enhances *in vitro* macrophage production from primitive murine hematopoietic progenitor cells in combination with stem cell factor and interleukin-7: novel stimulatory role of p55 TNF receptors. *Blood* 84(5):1528–33.
 - Fattori E, Cappelletti M, Costa P, Sellitto C, Cantoni L, Carelli M, et al. 1994. Defective inflammatory response in interleukin 6-deficient mice. *J Exp Med* 180(4):1243-50.
 - Ferrari D, Pizzirani C, Adinolfi E, Forchap S, Sitta B, Turchet L, et al. 2004. The antibiotic polymyxin B modulates P2X7 receptor function. *J Immunol* 173(7):4652-60.
 - Fink DL, Cope LD, Hansen EJ, Geme JW, 3rd. 2001. The *Hemophilus influenzae* Hap autotransporter is a chymotrypsin clan serine protease and undergoes autoproteolysis via an intermolecular mechanism. *J Biol Chem* 276(42):39492-500.
 - Fiorentini C, Fabbri A, Matarrese P, Falzano L, Boquet P, Malorni W. 1997. Hinderance of apoptosis and phagocytic behaviour induced by *Escherichia coli* cytotoxic necrotizing factor 1. Two related activities in epithelial cells. *Biochem Biophys Res Commun* 241(2):341–6.
 - Fischer H, Ellstrom P, Ekstrom K, Gustafsson L, Gustafsson M, Syanborg C. 2007. Ceramide as a TLR4 agonist; a putative signalling intermediate between sphingolipid receptors for microbial ligands and TLR4. *Cellular Microbiology* 9(5):1239–51.
 - Franz AR, Bauer K, Schalk A. 2004. Measurement of interleukin 8 in combination with C-reactive protein reduced unnecessary antibiotic therapy in newborn infants: a multicenter, randomized, controlled trial. *Pediatrics* 114(6):1–8.
 - Gabay C, Kushner I. 1999. Acute-phase proteins and other systemic responses to inflammation. *N Engl J Med* 340(6):448–54.

-
- Galanos C, Luderitz O, Rietschel ET, Westphal O, Brade H, Brade L, et al. 1985. Synthetic and natural *Escherichia coli* free lipid A express identical endotoxic activities. *Eur J Biochem* 148(1):1–5.
 - Ganatra HA, Stoll BJ, Zaidi AK. 2010. International perspective on early-onset neonatal sepsis. *Clin Perinatol* 37(2):501-23.
 - Garcia FG, Nager AL. 2002. Jaundice as an early diagnostic sign of urinary tract infection in infancy. *Pediatrics* 109(5):846-51.
 - Genevrois S, Steeghs L, Roholl P, Letesson JJ, van der Ley P. 2003. The Omp85 protein of *Neisseria meningitidis* is required for lipid export to the outer membrane. *EMBO J* 22(8):1780-9.
 - Gerard C, Bruyns C, Marchant A, Abramowicz D, Vandenabeele P, Delvaux A. 1993. Interleukin 10 reduces the release of tumor necrosis factor and prevents lethality in experimental endotoxemia. *J Exp Med* 177(2):547-50.
 - Gerdes JS. 2004. Diagnosis and management of bacterial infections in the neonates. *Pediatr Clin North Am.* 51(4):939-59.
 - Gheibi SH, Hagh S, Soleimani SH. 2005. Mortality and septicemia in neonates admitted into the NICU of Imam Khomeini Hospital of Urmia (Persian). *Med J Tabriz Uni Med Sci* 27(3):69-73.
 - Goetz DH, Holmes MA, Borregaard N, Bluhm ME, Raymond KN, Strong RK. 2002. The neutrophil lipocalin NGAL is a bacteriostatic agent that interferes with siderophore-mediated iron acquisition. *Mol Cell* 10(5):1033–43.
 - Gogia S, Sachdev HS. 2009. Neonatal vitamin A supplementation for prevention of mortality and morbidity in infancy: systematic review of randomized controlled trials. *BMJ* 338: b919.
 - Gogos CA, Drosou E, Bassaris HP, Skoutelis A. 2000. Pro- versus anti-inflammatory cytokine profile in patients with severe sepsis: a marker for prognosis and future therapeutic options. *J Infect Dis* 181(1):176–80.
 - Gophna U, Barlev M, Hacker J, Ron EZ. 2001b. Internalization of *Escherichia coli* serotype O78 by eukaryotic cells. *Infect Dis Rev Suppl* 3:37–42.

-
- Gophna U, Barlev M, Seiffers R, Oelschläger TA, Hacker J, Ron EZ. 2001c. Curli fibers mediate internalization of *Escherichia coli* by eukaryotic cells. *Infect Immun* 69(4):2659–65.
 - Gophna U, Ideses D, Rosen R, Grundland A, Ron EZ. 2004. OmpA of a septicemic *Escherichia coli* 078 secretion and convergent evolution. *Int J Med Microbiol* 294(6):373-81.
 - Gophna U, Oelschläger TA, Hacker J, Ron EZ. 2001a. *Yersinia* HPI in septicemic *Escherichia coli* strains isolated from diverse hosts. *FEMS Microbiol Lett* 196(1):57–60.
 - Gophna U, Oelschläger TA, Hacker J, Ron EZ. 2002. Role of fibronectin in curli-mediated internalization. *FEMS Microbiol Lett* 212(1):55–8.
 - Gophna U, Parket A, Hacker J, Ron EZ. 2003. A novel ColV plasmid encoding type IV pili. *Microbiology* 149(pt 1):177–84.
 - Griffin MP, Lake DE, O'Shea TM, Moorman JR. 2007. Heart rate characteristics and clinical signs in neonatal sepsis. *Pediatr Res* 61(2):222-7.
 - Grozdanov L, Raasch C, Schulze J, Sonnenborn U, Gottschalk G, Hacker J, et al. 2004. Analysis of the genome structure of the nonpathogenic probiotic *Escherichia coli* strain Nissle 1917. *J Bacteriol* 186(16):5432-41.
 - Guignot J, Chaplais C, Coconnier-Polter MH, and Servin AL. 2007. The secreted autotransporter toxin, Sat, functions as a virulence factor in Afa/Dr diffusely adhering *Escherichia coli* by promoting lesions in tight junction of polarized epithelial cells. *Cell Microbiol* 9(1):204-21.
 - Guyer DM, Henderson IR, Nataro JP, Mobley HL. 2000. Identification of Sat, an autotransporter toxin produced by uropathogenic *Escherichia coli*. *Mol Microbiol* 38(1): 53-66.
 - Guyer DM, Radulovic S, Jones FE, Mobley HL. 2002. Sat, the secreted autotransporter toxin of uropathogenic *Escherichia coli*, is a vacuolating cytotoxin for bladder and kidney epithelial cells. *Infect Immun* 70(8):4539–46.

-
-
- Guyer DM, Henderson IR, Nataro JP, Mobley HLT. 2000. Identification of Sat, an autotransporter toxin produced by uropathogenic *Escherichia coli*. *Mol Microbiol* 38(1):53-66.
 - Hack CE, De Groot ER, Felt-Bersma RJ, Nuijens JH, Strack Van Schijndel RJ, Eerenberg-Belmer AJ, et al. 1989. Increased plasma levels of interleukin-6 in sepsis. *Blood* 74(5):1704–10.
 - Harbarth S, Sudre P, Dharan S, Cadenas M, Pittet D. 1999. Outbreak of *Enterobacter cloacae* related to understaffing overcrowding and poor hygiene practices. *Infect Cont Hosp Epid* 20(9):598-603.
 - Hartleib S, Prager R, Hedenström I, Löfdahl S, Tschäpe H. 2003. Prevalence of the new, SPII-like, pathogenicity island ETT2 among *Escherichia coli*. *Int J Med Microbiol* 292(7-8):487–93.
 - Hase CC, Finkelstein RA. 1993. Bacterial extracellular zinc-containing metalloproteases. *Microbiol Rev* 57:823-37.
 - Healy CM, Baker CJ. 2007. Maternal immunization. *Pediatr Infect Dis J* 26(10):945–8.
 - Heck LW, Alarcon PG, Kulhavy RM, Morihara K, Russell MW, Mestecky JF. 1990. Degradation of IgA proteins by *Pseudomonas aeruginosa* elastase. *J Immunol* 144(6):2253-7.
 - Heck LW, Morihara K, McRae WB, Miller EJ. 1986. Specific cleavage of human type III and IV collagens by *Pseudomonas aeruginosa* elastase. *Infect Immun* 51(1):115-8.
 - Heimer SR, Rasko DA, Lockett CV, Johnson DE, Mobley HL. 2004. Autotransporter genes *pic* and *tsh* are associated with *Escherichia coli* strains that cause acute pyelonephritis and are expressed during urinary tract infection. *Infect Immun* 72(1):593-7.
 - Heimer SR, Rasko DA, Lockett CV, Johnson DE, Mobley HL. 2004. Autotransporter genes *pic* and *tsh* are associated with *Escherichia coli* strains that cause acute pyelonephritis and are expressed during urinary tract infection. *Infect Immun* 72(1):593–7.

-
- Heinzmann A, Ahlert I, Kurz T, Berner R, Deichmann KA. 2004. Association study suggests opposite effects of polymorphisms within IL-8 on bronchial asthma and respiratory syncytical virus bronchiolitis. *J Allergy Clin Immunol* 114(3):671-6.
 - Henderson IR, Czeczulin J, Eslava C, Noriega F, Nataro JP. 1999. Characterization of *pic*, a secreted protease of *Shigella flexneri* and enteroaggregative *Escherichia coli*. *Infect Immun* 67(11):5587-96.
 - Henderson IR, Hicks S, Navarro-Garcia F, Elias WP, Philips AD, Nataro JP. 1999. Involvement of the enteroaggregative *Escherichia coli* plasmid-encoded toxin in causing human intestinal damage. *Infect Immun* 67(10):5338-44.
 - Henderson IR, Nataro JP. 2001. Virulence functions of autotransporter proteins. *Infect Immun* 69(3):1231-43.
 - Henderson IR, Navarro-Garcia F, Nataro JP. 1998. The great escape: structure and function of the autotransporter proteins. *Trends Microbiol* 6(9):370-8.
 - Herzer PJ, Inouye S, Inouye M, Whittam TS. 1990. Phylogenetic distribution of branched RNA-linked multicopy single-stranded DNA among natural isolates of *Escherichia coli*. *J Bacteriol* 172(11):6175-81.
 - Hofer N, Zacharias E, Müller W, Resch B. 2012. An update on the use of C-reactive protein in early onset neonatal sepsis: current insights and new tasks. *Neonatology* 102(1):25-36.
 - Hotchkiss RS, Karl IE. 2003. The Pathophysiology and Treatment of Sepsis. *N Engl J Med* 348(2):138-50.
 - Hull RA, Gill RE, Hsu P. 1981. Construction and expression of recombinant plasmids encoding type 1 or D-mannose-resistant pili from a urinary tract infection *Escherichia coli* isolate. *Infect Immun* 33(3):933-8.
 - Ideses D, Gophna U, Paitan Y, Chaudhuri RR, Pallen MJ, Ron EZ. 2005. A degenerate type III secretion system from septicemic *Escherichia coli* contributes to pathogenesis. *J Bacteriol* 187(23):8164-71.
 - Janben T, Schwarz C, Preikschat P, Voss M, Philipp HC, Wieler LH. 2001. Virulence-associated genes in avian pathogenic *Escherichia coli* (APEC)

- isolated from internal organs of poultry having died from coli bacillosis. *Int J Med Microbiol* 291(5):371–8.
- Johanson IM, Plos K, Marklund BI, Svanborg C. 1993. *Pap*, *papG* and *prsG* DNA sequences in *Escherichia coli* from the fecal flora and the urinary tract. *Microb Pathog* 15(2):121-9.
 - Johnson JR, Clabots C, Kuskowski MA. 2008. Multiple-host sharing, long-term persistence, and virulence of *Escherichia coli* clones from human and animal household members. *J Clin Microbiol* 46(12):4078–82.
 - Johnson JR, Delavari P, Kuskowski M, Stell AL. 2001. Phylogenetic distribution of extraintestinal virulence-associated traits in *Escherichia coli*. *J Infect Dis* 183(1):78-88.
 - Johnson JR, Moseley S, Roberts P, Stamm WE. 1988. Aerobactin and other virulence factor genes among strains of *Escherichia coli* causing urosepsis: association with patient characteristics. *Infect Immun* 56(2):405-12.
 - Johnson JR, Russo TA. 2001. Extraintestinal pathogenic *Escherichia coli*: “The other bad *E. coli*”. *J Lab Clin Med* 139(3):155-62.
 - Johnson JR, Stapleton AE, Russo TA, Scheutz F, Brown JJ, Maslow JN. 1997. Characteristics and prevalence within serogroup O4 of a J96-like clonal group of uropathogenic *Escherichia coli* O4:H5 containing the class I and class III alleles of *papG*. *Infect Immun* 65(6):2153-9.
 - Johnson JR, Stell AL. 2000. Extended virulence genotypes of *Escherichia coli* strains from patients with urosepsis in relation to phylogeny and host compromise. *J Infect Dis* 181(1):261-72.
 - Johnson JR. 1991. Virulence factors in *Escherichia coli* urinary tract infection. *Clin Microbiol Rev* 4(1):80-128.
 - Joshi SG, Ghole VS, Niphadkar KB. 2000. Neonatal gram-negative bacteremia. *Indian J Pediatr* 67(1):27-32.
 - Jung HC, Eckmann L, Yang SK, Panja A, Fierer J, Morzycka-Wroblewska E, et al. 1995. A distinct array of proinflammatory cytokines is expressed in

- human colon epithelial cells in response to bacterial invasion. *J Clin Investig* 95(1):55-65.
- Kaftan H, Kinney JS. 1998. Early onset neonatal bacterial infections. *Semin Perinatol* 22(1):15-24.
 - Kaistha N, Mehta M, Singla N, Garg R, Chander J. 2010. Neonatal septicemia isolates and resistance patterns in a tertiary care hospital of North India. *J Infect Dev Ctries* 4(1):55-7.
 - Kaper JB, Nataro JP, Mobley HLT. 2004. Pathogenic *Escherichia coli*. *Nat Rev Microbiol* 2(2):123-40.
 - Karch H, Schubert S, Zhang D, Zhang W, Schmidt H, Oelschläger TA, et al. 1999. A genomic island, termed high-pathogenicity island, is present in certain non-O157 Shiga toxin-producing *Escherichia coli* clonal lineages. *Infect Immun* 67(11):5994-6001.
 - Karlowicz MG, Buescher ES, Surka AE. 2000. Fulminant late-onset sepsis in a neonatal intensive care unit, 1988-1997, and the impact of avoiding empiric vancomycin therapy. *Pediatrics* 106(6):1387-90.
 - Karthikeyan G, Premkumar K. 2001. Neonatal sepsis: *Staphylococcus aureus* as the predominant pathogen. *Indian J Pediatr* 68(8):715-7.
 - Keane WF, Welch R, Gekker G, and Peterson PK. 1987. Mechanism of *Escherichia coli* α -hemolysin-induced injury to isolated renal tubular cells. *Am J Pathol* 126(2):350-7.
 - Kellogg JA, Ferrentino FL, Goodstein MH, Liss J, Shapiro SL, Bankert DA. 1997. Frequency of low level bacteremia in infants from birth to two months of age. *Pediatr Infect Dis J* 16(4):381-5.
 - Kenny B. 2002. Mechanism of action of EPEC type III effector molecules. *Int J Med Microbiol* 291(6-7):469-77.
 - Kim KJ, Elliott SJ, Di Cello F, Stins MF, Kim KS. 2003. The K1 capsule modulates trafficking of *E. coli*-containing vacuoles and enhances intracellular bacterial survival in human brain microvascular endothelial cells. *Cell Microbiol* 5(4):245-52.

-
- Kohler CD, Dobrindt U. 2011. What defines extraintestinal pathogenic *Escherichia coli*? *Int J Med Microbiol* 301(8): 642-7.
 - Kostakioti M, Stathopoulos C. 2004. Functional analysis of the Tsh autotransporter from an avian pathogenic *Escherichia coli* strain. *Infect Immun* 72(10):5548-54.
 - Kotlowski R, Bernstein CN, Sepehri S, Krause DO. 2007. High prevalence of *Escherichia coli* belonging to the B2+D phylogenetic group in inflammatory bowel disease. *Gut* 56(5):669-75.
 - Koushik Roy, Scott Bartels, Firdausi Qadri, James M. Fleckenstein. 2010. Enterotoxigenic *Escherichia coli* elicits immune responses to multiple surface proteins. *Infect Immun* 78(7):3027-35.
 - Kristóf K, Kocsis E, Nagy K. 2009. Clinical microbiology of early onset and late-onset neonatal sepsis, particularly among preterm babies. *Acta Microbiol Immunol Hung* 56(1):21-51.
 - Kuhnert P, Hacker J, Mühldorfer I, Burnens AP, Nicolet J, Frey J. 1997. Detection system for *Escherichia coli* specific virulence genes: absence of virulence determinants in B and C strains. *Appl Environ Microbiol* 63(2):703-09.
 - Labigne-Roussel A, Falkow S. 1988. Distribution and degree of heterogeneity of the afimbrial-adhesin-encoding operon (*afa*) among uropathogenic *Escherichia coli* isolates. *Infect Immun* 56(3):640-8.
 - Labigne-Roussel AF, Lark D, Schoolnik G, Falkow S. 1984. Cloning and expression of an afimbrial adhesin (AFA-I) responsible for P blood group-independent, mannose-resistant hemagglutination from a pyelonephritic *Escherichia coli* strain. *Infect Immun* 46(1):251-9.
 - Laestadius A, Richter-Dahlfors A, Aperia A. 2002. Dual effects of *Escherichia coli* α -hemolysin on rat renal proximal tubule cells. *Kidney Int* 62(6):2035-42.
 - Lahariya C, Paul VK. 2010. Burden, Differentials, and Causes of Child Deaths in India. *Indian J Pediatr* 77(11):1312-21.

-
- Lam HS, Ng PC. 2008. Biochemical markers of neonatal sepsis. *Pathology* 40(2):141-8.
 - Lawn JE, Cousens S, Zupan J. 2005. 4 million neonatal deaths: when? Where? Why? *Lancet* 365(9462):891-900
 - Le Gall T, Clermont O, Gouriou S, Picard B. 2007. Extraintestinal virulence is a coincidental by-product of commensalism in B2 phylogenetic group *Escherichia coli* strains. *Mol Biol Evol* 24(11):2373–84.
 - Lemonnier M, Landraud L, Lemichez E. 2007. Rho GTPase-activating bacterial toxins: from bacterial virulence regulation to eukaryotic cell biology. *FEMS Microbiol Rev* 31(5):515–34.
 - Levy O. 2007. Innate immunity of the newborn: basic mechanisms and clinical correlates. *Nat Rev Immunol* 7(5):379–90.
 - Lin CY, Hsu CH, Huang FY, Chang JH, Hung HY, Kao HA, et al. 2011. The changing face of early-onset neonatal sepsis after the implementation of amaternal group B *Streptococcus* screening and intrapartum prophylaxis policy-a study in one medical center. *Pediatr Neonatol* 52(2):78–84.
 - Liu L, Oza S, Hogan D, Perin J, Rudan I, Lawn JE, et al. 2015. Global, regional, and national causes of child mortality in 2000–13, with projections to inform post-2015 priorities: an updated systematic analysis. *Lancet* 385(9966):430-40.
 - Loisa P, Rinne T, Laine S, Hurme M, Kaukinen S. 2003. Anti-inflammatory cytokine response and the development of multiple organ failure in severe sepsis. *Acta Anaesthesiol Scand* 47(3):319–25.
 - Louvois J. Acute bacterial meningitis in the newborn. 1994. *J Antimicrob Chemother* 34(suppl A):61-73.
 - Lund B, Marklund BI, Stromberg N, Lindberg F, Karlsson KA, Normark S. 1988. Uropathogenic *Escherichia coli* can express serologically identical pili of different receptor binding specificities. *Mol Microbiol* 2(2):255-63.
 - Luo Q, Kumar P, Vickers TJ, Sheikh A, Lewis WG, Rasko DA, et al. 2014. Enterotoxigenic *Escherichia coli* Secretes a Highly Conserved Mucin-

Degrading Metalloprotease To Effectively Engage Intestinal Epithelial Cells. *Infect Immun* 82(2):509-21.

- Machado JR, Soave DF, Vinícius da Silva M, Borges de Menezes L, Etchebehere RM, Goncalves dos Reis Monteiro ML, et al. 2014. Neonatal Sepsis and Inflammatory Mediators. *Mediators of Inflammation*. Volume 2014, Article ID 269681, 10 pages.
- Macharashvili N, Kourbatova E, Butsashvili M, Tsertsvadze T, McNutt LA, Leonard MK. 2009. Etiology of neonatal bloodstream infections in Tbilisi, Republic of Georgia. *Int J Infect Dis* 13(4):499-505.
- Malaguarnera M, Di Fazio I, Romeo MA, Restuccia S, Laurino A, Trovato BA. 1997. Elevation of interleukin 6 levels in patients with chronic hepatitis due to hepatitis C virus. *J Gastroenterol* 32(2):211–5.
- Marre R, Kreft B, Hacker J. 1990. Genetically engineered S and F1C fimbriae differ in their contribution to adherence of *Escherichia coli* to cultured renal tubular cells. *Infect Immun* 58(10):3434–7.
- Mayor-Lynn K, Gonz´alez-Quintero VH, O’Sullivan MJ, Hartstein AI, Roger S, Tamayo M. 2005. Comparison of early-onset neonatal sepsis caused by *Escherichia coli* and group B *Streptococcus*. *Am J Obstetric Gynaecol* 192(5):1437–9.
- Medzhitov R, Janeway CA Jr. 1999. Innate immune induction of the adaptive immune response. *Cold Spring Harb Symp Quant Biol* 64:429-35.
- Medzhitov R, Preston-Hurlburt P, Janeway CA Jr. 1997. A human homologue of the *Drosophila* Toll protein signals activation of adaptive immunity. *Nature* 388:394–7.
- Mehr S, Doyle LW. 2000. Cytokines as markers of bacterial sepsis in newborn infants: a review. *Pediatr Infect Dis J* 19(9):879-87.
- Mellies JL, Navarro-Garcia F, Okeke I, Frederickson J, Nataro JP, Kaper JB. 2001. *espC* pathogenicity island of enteropathogenic *Escherichia coli* encodes an enterotoxin. *Infect Immun* 69(1):315-24.

-
- Messai FM, Bidet P, Caro V, Diancourt L, Brian V, Aujard Y, et al. 2011. *Escherichia coli* isolates causing bacteremia via gut translocation and urinary tract infection in young infants exhibit different virulence genotypes. *J Infect Dis* 203(12):1844-9.
 - Mills M, Meysick KC, O'Brien AD. 2000. Cytotoxic necrotizing factor type 1 of uropathogenic *Escherichia coli* kills cultured human uroepithelial 5637 cells by an apoptotic Mechanism. *Infect Immun* 68(10):5869–80.
 - Moch T, Hoschutzky H, Hacker J, Kroncke KD, Jann K. 1987. Isolation and characterization of the α -sialyl- β -2,3-galactosyl-specific adhesin from fimbriated *Escherichia coli*. *Proc Natl Acad Sci USA* 84(10):3462-6.
 - Mokady D, Gophna U, Ron EZ. 2005. Virulence factors of septicemic *Escherichia coli* strains. *Int J Med Microbiol* 295(6-7):455-62.
 - Moore D. 2004. Nosocomial infections in newborn nurseries and neonatal intensive care units: In: Mayhall C.G., *Hospital epidemiology and infection control*. 3rd ed. Pa: Lippincott Williams & Wilkins, USA, p852-83.
 - Moore RA, Bates NC, Hancock RE. 1986. Interaction of polycationic antibiotics with *Pseudomonas aeruginosa* lipopolysaccharide and lipid A studied by using dansyl-polymyxin. *Antimicrob Agents Chemother* 29(3):496–500.
 - Moriela DG, Bertoldia I, Spagnuoloa A, Marchia S, Rosini R, Nesta B, et al. 2010. Identification of protective and broadly conserved vaccine antigens from the genome of extraintestinal pathogenic *Escherichia coli*. *Proc Natl Acad Sci* 107(20):9072-7.
 - Mullany LC, Darmstadt GL, Tielsch JM. 2006. Safety and impact of chlorhexidine antiseptics interventions for improving neonatal health in developing countries. *Pediatr Infect Dis J* 25(8):665–75.
 - Multu M, Aslan Y, Saygin B, Yilmaz G, Bayramoglu G, Koksai I. 2011. Neonatal Sepsis Caused by Gram-negative Bacteria in a Neonatal Intensive Care Unit: A Six Years Analysis. *HK J Paediatr* 16:253-7.
 - Mulvey MA. 2002. Adhesion and entry of uropathogenic *Escherichia coli*. *Cellular Microbiology* 4(5):257–71.

-
- Nakjang S, Ndeh DA, Wipat A, Bolam DN, Hirt RP. 2012. A novel extracellular metallopeptidase domain shared by animal host-associated mutualistic and pathogenic microbes. *PLoS One* 7(1):e30287.
 - Nast-Kolb D, Wayd Has C, Gippner-Steppert C, Schneider I, Trupka A, Ruchholtz S, et al. 1997. Indicators of the posttraumatic inflammatory response correlate with organ failure in patients with multiple injuries. *J Trauma* 42(3):446–54; discussion 454-55.
 - Navarro-Garcia F, Canizalez-Roman A, Bao Quan S, Nataro JP, Azamar Y. 2004. The serine protease motif of EspC from enteropathogenic *Escherichia coli* produces epithelial damage by a mechanism different from that of Pet toxin from enteroaggregative *E. coli*. *Infect. Immun* 72(6): 3609–21.
 - Navarro-Garcia F, Gutierrez-Jimenez J, Garcia-Tovar C, Castro LA, Salazar-Gonzalez H, Cordova V. 2010. Pic, an autotransporter protein secreted by different pathogens in the *Enterobacteriaceae* family, is a potent mucus secretagogue. *Infect Immun* 78(10): 4101-9.
 - Navarro-Garcia F, Eslava C, Villaseca JM, Lopez-Revilla R, Czeczulin JR, Srinivas S, et al. 1998. *In vitro* effects of a high-molecular-weight heat-labile enterotoxin from enteroaggregative *Escherichia coli*. *Infect Immun* 66(7):3149-54.
 - Neilands JB, Bindereif A, Montgomerie JZ. 1985. Genetic basis of iron assimilation in pathogenic *Escherichia coli*. *Curr Top Microbiol Immunol* 118:179-95.
 - Netea MG, van der Meer JW, van Deuren M, Kullberg BJ. 2003. Proinflammatory cytokines and sepsis syndrome: not enough, or too much of a good thing? *Trends in Immun* 24(5):254-8.
 - Ng PC, Lam HS. 2010. Biomarkers for late-onset neonatal sepsis: cytokines and beyond. *Clin Perinatol* 37(3):599-610.
 - Nizet V, Klein JO. Bacterial sepsis and meningitis. Chapter 6. 225-278. DOI: 10.1016/B978-1-4160-6400-8.00006-7.
 - Nougayrede JP, Fernandes PJ, Donnenberg MS. 2003. Adhesion of enteropathogenic *Escherichia coli* to host cells. *Cell Microbiol* 5(6):359–72.

-
-
- Nowicki B, Rhen M, Vaisanen-Rhen V, Pere A, Korhonen TK. 1985. Fractionation of a bacterial cell population by adsorption to erythrocytes and yeast cells. *FEMS Microbiol Lett* 26(1):35-40.
 - O'Neill LAJ. 2008. The interleukin-1 receptor/Toll-like receptor superfamily: 10 years of progress. *Immunol Rev* 226(1):10–18.
 - Ohlsson A, Shah VS. 2009. Intrapartum antibiotics for known maternal Group B Streptococcal colonization. *Cochrane Database Syst Rev*. pp CD007467.
 - Okusawa S, Gelfand JA, Ikejima T, Connolly RJ, Dinarello CA. 1988. Interleukin 1 induces a shock-like state in rabbits. Synergism with tumor necrosis factor and the effect of cyclooxygenase inhibition. *J Clin Invest* 81(4):1162–72.
 - Ostolaza H, Goni FM. 1995. Interaction of the bacterial protein toxin alpha-haemolysin with model membranes: protein binding does not always lead to lytic activity. *FEBS Lett* 371(1):303–06.
 - Otto BR, Van Dooren SJ, Dozois CM, Luirink J, Oudega B. 2002. *Escherichia coli* hemoglobin protease autotransporter contributes to synergistic abscess formation and heme-dependent growth of *Bacteroides fragilis*. *Infect Immun* 70(1):5-10.
 - Otto BR, van Dooren SJ, Nuijens JH, Luirink J, Oudega B. 1998. Characterization of a hemoglobin protease secreted by the pathogenic *Escherichia coli* strain EB1. *J Exp Med* 188(6): 1091-1103.
 - Otto G, Sandberg T, Marklund BI, Ulleryd P, Svanborg C. 1993. Virulence factors and *pap* genotype in *Escherichia coli* isolates from women with acute pyelonephritis, with or without bacteremia. *Clin Infect Dis* 17(3):448-56.
 - Paolucci M, Landini MP, Sambri V. 2012. How can the microbiologist help in diagnosing neonatal sepsis? *Int J Pediatr* Volume 2012, Article ID 120139, 14 pages doi:10.1155/2012/120139
 - Parameswaran N, Patial S. 2010. Tumor necrosis factor- α signalling in macrophages. *Crit Rev Eukaryot Gene Expr* 20(2):87–103.

-
- Parham NJ, Pollard SJ, Desvaux M, Scott-Tucker A, Liu C, Fivian A, et al. 2005. Distribution of the Serine Protease Autotransporters of the *Enterobacteriaceae* among extraintestinal clinical isolates of *Escherichia coli*. *J Clin Microbiol* 43(8):4076-82.
 - Parham NJ, Srinivasan U, Desvaux M, Foxman B, Marrs CF, Henderson IR. 2004. PicU, a second serine protease autotransporter of uropathogenic *Escherichia coli*. *FEMS Microbiol Lett* 230(1):73–83.
 - Parkkinen J, Finne J, Achtman M, Vaisanen V, Korhonen TK. 1983. *Escherichia coli* strains binding neuraminyl α 2-3 galactosides. *Biochem Biophys Res Commun* 111(2):456-61.
 - Parreira VR, Gyles CL. 2003. A Novel Pathogenicity Island Integrated Adjacent to the *thrW* tRNA Gene of Avian Pathogenic *Escherichia coli* Encodes a Vacuolating Autotransporter Toxin. *Infect Immun* 71(9):5087-96.
 - Patel SK, Dotson J, Allen KP, Fleckenstein JM. 2004. Identification and molecular characterization of EatA, an autotransporter protein of enterotoxigenic *Escherichia coli*. *Infect Immun* 72(3):1786-94.
 - Penders J, Thijs C, Vink C, Stelma FF, Snijders B, Kummeling I, et al. 2006. Factors influencing the composition of the intestinal microbiota in early infancy. *Pediatrics* 118(2):511-21.
 - Picard B, Garcia JS, Gouriou S, Duriez P, Brahimi N, Bingen E, et al. 1999. The link between phylogeny and virulence in *Escherichia coli* extraintestinal infection. *Infect Immun* 67(2):546-53.
 - Pickler R, Brown L, McGrath J, Lyon D, Rattican D, Cheng CY, et al., 2010. Integrated review of cytokines in maternal, cord, and newborn blood: part II—associations with early infection and increased risk of neurologic damage in preterm infants. *Biol Res Nurs* 11(4):377–86.
 - Pierrakos C, Vincent JL. 2010. Sepsis biomarkers: a review. *Crit Care* 14(1):R15. doi:10.1186/cc8872.
 - Pittet D, Boyce JM. 2001. Hand hygiene and patient care: pursuing the Semmelweis legacy. *Lancet Infect Dis* 1(1):9-20.

-
- Poltorak A, He X, Smirnova I, Liu MY, Van Huffel C, Du X, et al. 1998. Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: mutations in Tlr4 gene. *Science* 282(5396):2085–8.
 - Preiser JC, Schmartz D, Van der Linden P, Content J, Vanden Bussche P, Buurman W, et al. 1991. Interleukin-6 administration has no acute hemodynamic or hematologic effect in the dog. *Cytokine* 3(1):1–4.
 - Provence DL, Curtiss R, 3rd. 1992. Role of *crl* in avian pathogenic *Escherichia coli*: a knockout mutation of *crl* does not affect hemagglutination activity, fibronectin binding, or Curli production. *Infect Immun* 60(11):4460-7.
 - Provence DL, Curtiss R, 3rd. 1994. Isolation and characterization of a gene involved in hemagglutination by an avian pathogenic *Escherichia coli* strain. *Infect Immun* 62(4):1369-80.
 - Raetz CR, Whitfield C. 2002. Lipopolysaccharide endotoxins. *Annu Rev Biochem* 71: 635–700.
 - Rajakumar K, Sasakawa C, Adler B. 1997. Use of a novel approach, termed island probing, identifies the *Shigella flexneri she* pathogenicity island which encodes a homolog of the immunoglobulin A protease-like family of proteins. *Infect Immun* 65(11):4606-14.
 - Rasko DA, Webster DR, Sahl JW, Bashir A. 2011. Origins of the *E. coli* strain causing an outbreak of hemolytic-uremic syndrome in Germany. *N Engl J Med* 365(8):709-17.
 - Reinhart K, Bauer M, Riedemann NC, Hartog CS. 2012. New Approaches to Sepsis: Molecular Diagnostics and Biomarkers. *Clin Microbiol Rev* 25(4):609–34.
 - Ren CP, Chaudhuri RR, Fivian A, Bailey CM, Antonio M, Barnes WM, et al. 2004. The *ETT2* gene cluster, encoding a second type III secretion system from *Escherichia coli*, is present in the majority of strains but has undergone widespread mutational attrition. *J Bacteriol* 186(11):3547–60.
 - Restieri C, Garriss G, Locas MC, Dozois CM. 2007. Autotransporter-Encoding Sequences are phylogenetically distributed among *Escherichia coli* clinical isolates and reference strains. *Appl Environ Microbiol* 73(5):1553-62.

-
- Rhee V, Mullany LC, Khattry SK, Katz J, LeClerq SC, Darmstadt GL, et al. 2008. Maternal and birth attendant hand washing and neonatal mortality in southern Nepal. *Arch Pediatr Adolesc Med* 162(7):603–08.
 - Robak T, Gladalska A, Stepień H, Robak E. 1998. Serum levels of interleukin-6 type cytokines and soluble interleukin-6 receptor in patients with rheumatoid arthritis. *Mediators Inflamm* 7(5):347–53.
 - Ron EZ. 2010. Distribution and evolution of virulence factors in septicemic *Escherichia coli*. *Int J Med Microbiol* 300(6):367-70.
 - Roy I, Jain A, Kumar M, Agarwal SK. 2002. Bacteriology of neonatal septicemia in a tertiary care Hospital of Northern India. *Indian J Med Microbiol* 20(3):156-9.
 - Roy S, Singh AK, Viswanathan R, Nandy RK, Basu S. 2011. Transmission of imipenem resistance determinants during the course of an outbreak of NDM-1 *Escherichia coli* in a sick newborn care unit. *J Antimicrob Chemother* 66(12):2773-80.
 - Ruiz-Perez F, Nataro JP. 2014. Bacterial serine proteases secreted by the autotransporter pathway: classification, specificity, and role in virulence. *Cell Mol Life Sci* 71(5):745-70.
 - Ruiz-Perez F, Wahid R, Faherty CS, Kolappaswamy K. 2011. Serine protease autotransporters from *Shigella flexneri* and pathogenic *Escherichia coli* target a broad range of leukocyte glycoproteins. *Proc Natl Acad Sci U S A* 108(31):12881-6.
 - Ruiz-Perez, F, Henderson IR, Leyton DL, Rossiter AE, Zhang Y, Nataro JP. 2009. Roles of periplasmic chaperone proteins in the biogenesis of serine protease autotransporters of *Enterobacteriaceae*. *J Bacteriol* 191(21):6571-83.
 - Russo TA, Johnson JR. 2000. Proposal for a new inclusive designation for extraintestinal pathogenic isolates of *Escherichia coli*: ExPEC. *J Infect Dis* 181(5):1753-4.
 - Russo TA, Carlino UB, Mong A, Jodush ST. 1999. Identification of genes in an extraintestinal isolate of *Escherichia coli* with increased expression after exposure to human urine. *Infect Immun* 67(10):5306–14.

-
- Russo TA, Davidson BA, Genagon SA, Warholc NM, Macdonald U, Pawlicki PD, et al. 2005. *E. coli* virulence factor hemolysin induces neutrophil apoptosis and necrosis/lysis in vitro and necrosis/lysis and lung injury in a rat pneumonia mode. *Am J Physiol* 289(2):L207–16.
 - Russo TA, Johnson JR. 2003. Medical and economic impact of extraintestinal infections due to *Escherichia coli*: focus on an increasingly important endemic problem. *Microb Infect* 5(5):449-56.
 - Russo TA, McFadden CD, Carlino-MacDonald UB, Beanan JM, Barnard TJ, Johnson JR. 2002. IroN functions as a siderophore receptor and is a urovirulence factor in an extraintestinal pathogenic isolate of *Escherichia coli*. *Infect Immun* 70(12):7156–60.
 - Salek-Ardakani S, Stuart AD, Arrand JE, Lyons S, Arrand JR, Mackett M. 2002. High level expression and purification of the Epstein-Barr virus encoded cytokine viral interleukin 10: efficient removal of endotoxin. *Cytokine* 17(1):1-13.
 - Salender RK, Korhonen TK, Vaisanen-Rhen V, Williams PH, Pattison PE, Caugant DA. 1986. Genetic relationships and clonal structure of strains of *Escherichia coli* causing neonatal septicemia and meningitis. *Infect Immun* 52(1):213-22.
 - Sandt CH, Hill CH. 2000. Four different genes responsible for nonimmune immunoglobulin-binding activities within a single strain of *Escherichia coli*. *Infect Immun* 68(4):2205–14.
 - Sanghvi KP, Tudehope DI. 1996. Neonatal bacterial sepsis in a neonatal intensive care unit: a 5 year analysis. *J Paediatr Child Health* 32(4):333–8.
 - Sankar MJ, Agarwal R, Deorari AK, Paul VK. 2008. Sepsis in the newborn. *Indian J Pediatr* 75(3):261-6.
 - Sarkar M, Chaudhuri K. 2004. Association of adherence and motility in interleukin 8 induction in human intestinal epithelial cells by *Vibrio cholerae*. *Microb Infect* 6(7):676-85.
 - Scheller J, Rose-John S. 2006. Interleukin-6 and its receptor: from bench to bedside. *Med Microbiol Immun* 195(4):173–83.

-
- Schelonka RL, Chai MK, Yoder BA, Hensley D, Brockett RM, Ascher DP. 1996. Volume of blood required to detect common neonatal pathogens. *J Pediatr* 129(2):275-8.
 - Schindler R, Mancilla J, Endres S, Ghorbani R, Clark SC, Dinarello CA. 1990. Correlations and interactions in the production of interleukin-6 (IL-6), IL-1, and tumor necrosis factor (TNF) in human blood mononuclear cells: IL-6 suppresses IL-1 and TNF. *Blood* 75(1):40-7.
 - Scholl D, Adhya S, Merrill C. 2005. *Escherichia coli* K1's capsule is a barrier to bacteriophage T7. *Appl Environ Microbiol* 71(8):4872-4.
 - Schouten M, Wiersinga WJ, Levi M, Van Der Poll T. 2008. Inflammation, endothelium, and coagulation in sepsis. *J Leukoc Biol* 83(3):536-45.
 - Schubert S, Rakin A, Karch H, Carniel E, Heesemann, J. 1998. Prevalence of the “high-pathogenicity island” of *Yersinia* species among *Escherichia coli* strains that are pathogenic to humans. *Infect Immun* 66(2):480-5.
 - Schulte W, Bernhagen J, Bucala R. 2013. Cytokines in Sepsis: Potent Immunoregulators and Potential Therapeutic Targets—An Updated View. *Mediators of Inflammation*. Volume 2013, Article ID 165974, 16 pages.
 - Schultz C, Strunk T, Temming P, Matzke N, Hartel C. 2007. Reduced IL-10 production and -receptor expression in neonatal T lymphocytes. *Acta Paediatrica* 96(8):1122-5.
 - Sexton PM, Christopoulos G, Christopoulos A, Nylén ES, Snider RH Jr, Becker KL. 2008. Procalcitonin has bioactivity at calcitonin receptor family complexes: potential mediator implications in sepsis. *Crit Care Med* 36(5):1637-40.
 - Shah BA, Padbury JF. 2014. Neonatal sepsis. *Virulence* 5(1):170-8.
 - Shah SS, Ehrenkranz RA, Gallagher PG. 1999. Increasing incidence of Gram-negative rod bacteremia in a newborn intensive care unit. *Pediatr Infect Dis J* 18(7):591-5.

-
-
- Shannon E, Noveck R, Sandoval F, Kamath B, Kearney M. 2007. Thalidomide suppressed interleukin-6 but not tumor necrosis factor-alpha in volunteers with experimental endotoxemia. *Transl Res* 150(5):275–80.
 - Sikora JP, Chlebna Sokol D, Krzyzanska-Oberbek A. 2001. Proinflammatory cytokines (IL-6, IL-8), cytokine inhibitors (IL-6sR, sTNFRII) and anti-inflammatory cytokines (IL-10, IL-13) in the pathogenesis of sepsis in newborns and infants. *Arch Immunol Ther Expe* 49(5):399-404.
 - Silveria RC, Procianoy RS. 1999. Evaluation of interleukin-6, tumor necrosis factor-alpha and interleukin-1beta for early diagnosis of neonatal sepsis. *Acta Paediatr* 88(6):647-50.
 - Smith YC, Grande KK, Rasmussen SB, O'Brien AD. 2006. Novel three-dimensional organoid model for evaluation of the interaction of uropathogenic *Escherichia coli* with terminally differentiated human urothelial cells. *Infect and Immun* 74(1):750–7.
 - Stanilova SA, Karakolev ZT, Dimov GS, Dobрева JG, Miteva LD, Slavov ES, et al. 2005. High interleukin 12 and low interleukin 10 production after *in vitro* stimulation detected in sepsis survivors. *Intensive Care Med* 31(3):401–07.
 - Stoll BJ, Hansen N, Fanaroff AA, Wright LL, Carlo WA, Ehrenkranz RA, et al. 2002. Changes in pathogens causing early-onset sepsis in very-low-birth-weight infants. *N Engl J Med* 347(4):240–7.
 - Stoll BJ, Hansen NI, Higgins RD, Fanaroff AA, Duara S, Goldberg R, et al. 2005. Very low birth weight preterm infants with early onset neonatal sepsis: the predominance of gram-negative infections continues in the National Institute of Child Health and Human Development Neonatal Research Network, 2002-2003. *Pediatr Infect Dis J* 24(7):635-9.
 - Stoll BJ, Hansen NI, Sánchez PJ, Faix RG, Poindexter BB, Van Meurs KP, et al. 2011. Early onset neonatal sepsis: the burden of group B Streptococcal and *E. coli* disease continues. *Pediatrics* 127(5):817-26.

-
- Stoll BJ, Holman RC, Schuchat A. 1998. Decline in sepsis-associated neonatal and infant deaths in the United States, 1979 through 1994. *Pediatrics* 102(2):1-7.
 - Stoll BJ, Zaidi AK. 2010. International perspective on early-onset neonatal sepsis. *Clin Perinatol* 37(2):501-23.
 - Spurbeck RR, Dinh PC Jr, Walk ST, Stapleton AE, Hooton TM, Nolan LK, et al. 2012. *Escherichia coli* isolates that carry *vat*, *fyuA*, *chuA*, and *yfcV* efficiently colonize the urinary tract. *Infect Immun* 80(12):4115-22.
 - Syngkon A, Elluri S, Koley H, Rompikuntal PK, Saha DR, Chakrabarti MK, et al. 2010. Studies on a novel serine protease of a $\Delta hapA\Delta prtV$ *Vibrio cholerae* O1 strain and its role in hemorrhagic response in the rabbit ileal loop model. *PloS One* 5(9):1-11.
 - Taddei CR, Moreno ACR, Filho AF, Montemor LPG, Martinez MB. 2003. Prevalence of secreted autotransporter toxin gene among diffusely adhering *Escherichia coli* isolated from stools of children. *FEMS Microbiol Lett* 227(2):249-53.
 - Tang BM, Huang SJ, McLean AS. 2010. Genome-wide transcription profiling of human sepsis: a systematic review. *Crit Care* 14(6):R237. doi: 10.1186/cc9392.
 - Tenaillon O, Skurnik D, Picard B, Denamur E. 2010. The population genetics of commensal *Escherichia coli*. *Nat Rev Microbiol* 8:207–17.
 - Tobias PS, Mathison J, Mintz D, Lee JD, Kravchenko V, Kato K, et al. 1992. Participation of lipopolysaccharide-binding protein in lipopolysaccharide-dependent macrophage activation. *Am J Respir Cell Mol Biol* 7(3):239–45.
 - Tripathi S, Malik GK. 2010. Neonatal Sepsis: past, present and future; a review article. *Internet Journal of Medical Update* 5(2):45-54.
 - Tsai CH, Chen YY, Wang KG, Chen CY, Chen CP. 2012. Characteristics of early-onset neonatal sepsis caused by *Escherichia coli*. *Taiw J Obs Gynec* 51(1):26-30.

-
- Tullus K, Brauner A, Frykland B, Munkhammar T, Rabsch W, Reissbrodt R, et al., 1992. Host factors versus virulence-associated bacterial characteristics in neonatal and infantile bacteraemia and meningitis caused by *Escherichia coli*. *J Med Microbiol* 36(3):203-08.
 - Turner D, Hammerman C, Rudensky B, Schlesinger Y, Schimmel MS. 2006. The role of procalcitonin as a predictor of nosocomial sepsis in preterm infants. *Acta Paediatr* 95(12):1571-6.
 - Turunen R, Nupponen I, Siitonen S, Repo H, Anderson S. 2006. Onset of mechanical ventilation is associated with rapid activation of circulating phagocytes in preterm infants. *Pediatrics* 117(2):448-54.
 - Uhlen P, Laestadius A, Jahnukainen T, Soderblom T, Backhed F, Celsi G. 2000. α -Haemolysin of uropathogenic *E. coli* induces Ca^{2+} oscillations in renal epithelial cells. *Nature* 405(6787):694–7.
 - Ulevitch RJ, Tobias PS. 1995. Receptor-dependent mechanisms of cell stimulation by bacterial endotoxin. *Annu Rev Immunol* 13:437–57.
 - Usein CR, Grigore LA, Georgescu RM, Baltoiu MC, Condei M, Telean MD. 2011. Phylogenetic background and extraintestinal virulence genotypes of *Escherichia coli* vaginal strains isolated from adult women. *Revista Română de Medicină de Laborator* 19:37-44.
 - Valvano MA, Crosa JH. 1984. Aerobactin iron transport genes commonly encoded by certain ColV plasmids occur in the chromosome of a human invasive strain of *Escherichia coli* K1. *Infect Immun* 46(1):159–67.
 - Van der Kleij D, Van den Biggelaar AH, Kruize YC, Retra K, Fillie Y, Schmitz M, et al. 2004. Responses to Toll-like receptor ligands in children living in areas where schistosome infections are endemic. *J Infect Dis* 189(6):1044-51.
 - Veiga E, Sugawara E, Nikaido H, de Lorenzo V, Fernandez LA. 2002. Export of autotransported proteins proceeds through an oligomeric ring shaped by C-terminal domains. *EMBO J* 21(9):2122–31.

-
- Vidal JE, Navarro-Garcia F. 2006. Efficient translocation of EspC into epithelial cells depends on enteropathogenic *Escherichia coli* and host cell contact. *Infect Immun* 74(4):2293-303.
 - Vidal JE, Navarro-Garcia F. 2008. EspC translocation into epithelial cells by enteropathogenic *Escherichia coli* requires a concerted participation of type V and III secretion systems. *Cell Microbiol* 10(10):1975-86.
 - Viswanathan R, Singh AK, Basu S, Chatterjee S, Sardar S, Isaacs D. 2012. Multi-drug resistant gram negative bacilli causing early neonatal sepsis in India. *Arch Dis Child Fetal Neonatal Ed* 97(3):F182-7.
 - Waters D, Jawad I, Ahmad A. 2011. Aetiology of community-acquired neonatal sepsis in low- and middle-income countries. *Journal of global health* 1(2):154-70.
 - Watt S, Lanotte P, Mereguetti L, Moulin-Schouleur M, Picard B, Quentin R. 2003. *Escherichia coli* strains from pregnant women and neonates: intraspecies genetic distribution and prevalence of virulence factors. *J Clin Microbiol* 41(5):1929–35.
 - Weighardt H, Heidecke C, Westerholt A, Emmanuilidis K, Maier S, Veit M, et al. 2002. Impaired monocyte IL-12 production before surgery as a predictive factor for the lethal outcome of postoperative sepsis. *Ann Surg* 235(4):560–7.
 - Weston EJ, Pondo T, Lewis MM, Martell-Cleary P, Morin C, Jewell B, et al. 2011. The burden of invasive early-onset neonatal sepsis in the United States, 2005-2008. *Pediatr Infect Dis J* 30(11):937-41.
 - Wiles TJ, Kulesus RR, Mulvey MA. 2008. Origins and virulence mechanisms of uropathogenic *Escherichia coli*. *Exp Mol Pathol* 85(1):11–9.
 - Witsell AL, Schook LB. 1992. Tumor necrosis factor alpha is an autocrine growth regulator during macrophage differentiation. *Proceedings of the National Academy of Sciences of the United States of America* 89(10) 4754–8.
 - World Health Organization. 1999. Serious infections in young infants in developing countries: rationale for a multicentre study. *Paediatr Infect Dis J* 18(10):54-7.

-
- Wullt B, Bergsten G, Connell H, Rollano P, Gebretsadik N, Hull R, et al. 2000. P fimbriae enhance the early establishment of *Escherichia coli* in the human urinary tract. *Mol Microbiol* 38(3):456–64.
 - Wurfel MM, Kunitake ST, Lichenstein H, Kane JP, Wright SD. 1994. Lipopolysaccharide (LPS)-binding protein is carried on lipoproteins and acts as a cofactor in the neutralization of LPS. *J Exp Med* 180(3):1025–35.
 - Xicohtencatl-Cortes J, Saldana Z, Deng W, Castaneda E, Freer E, Tarr PI et al. 2010. Bacterial macroscopic rope-like fibers with cytopathic and adhesive properties. *J Biol Chem* 285(42):32336-42.
 - Yalaz M, Cetin H, Akisu M. 2006. Neonatal nosocomial sepsis in a level-111 NICU evaluation of the causative agents and antimicrobial susceptibilities. *Turkish J Paediatr* 48(1):13-18.
 - Yamamoto S, Terai A, Yuri K, Kurazono H, Takeda Y, Yoshida O. 1995. Detection of urovirulence factors in *Escherichia coli* by multiplex polymerase chain reaction. *FEMS Immunol Med Microbiol* 12(2):85-90.
 - Yen YT, Kostakioti M, Henderson IR, Stathopoulos C. 2008. Common themes and variations in serine protease autotransporters. *Trends Microbiol* 16(8):370-9.
 - Young JAT, Collier RJ. 2007. Anthrax Toxin: Receptor Binding, Internalization, Pore Formation, and Translocation. *Ann Rev Biochem* 76:243-65.
 - Zaidi AKM, Huskins WC, Thaver D, Bhutta ZA, Abbas Z, Goldmann DA. 2005. Hospital-acquired neonatal infections in developing countries. *Lancet* 365(9465):1175-88.
 - Zaidi AKM, Thaver D, Ali SA, Khan TA. 2009. Pathogens associated with sepsis in newborns and young infants in developing countries. *Pediatr Infect Dis J* 28(1):S10-S18.
 - Zgur-Bertok D, Modric E, Grabnar M. 1990. Aerobactin uptake system, ColV production, and drug resistance encoded by a plasmid from an urinary tract infection *Escherichia coli* strain of human origin. *Can J Microbiol* 36(4): 297–9.