

ACKNOWLEDGEMENTS

I owe my deep sense of gratitude and indebtedness to Dr. Sukh Dev, D.Sc., FNA, presently Director, Malti-Chem. Research Centre, Baroda, who in addition to his conspicuous and meticulous guidance, gave me an opportunity to use modern armoury of organic chemist in its entirety of aspects. I deem myself most fortunate to be in his research group and am very grateful for all his kindness.

I am grateful to Dr. A.S. Gupta for his prompt help and encouragements. I pay my warmest thanks to Drs. S.C. Sethi, H.R. Sonawane and S. Krishnappa, for their whole-hearted co-operation. My sincere thanks are also due to Dr. H.P.S. Chawla for his helpful discussions.

I owe a lot to my elder brother, Dr. A.S. Narula, for his constant advice, inspiration and valuable suggestions.

I wish to thank Glass Blowing, Micro Analysis, Spectroscopic (UV, IR, NMR, GLC and Mass spectra) Sections for their prompt assistance. I am also thankful to all my friends and colleagues for their help and co-operation.

I thank the Council of Scientific and Industrial Research, New Delhi, for the award of Junior Research Fellowship during the tenure of this work. I also thank the Director, National Chemical Laboratory, Poona 8, for his kind permission to submit this work in the form of a thesis.

Finally, I am thankful to Mr. S. Venkataraman for the excellent typing of this thesis and Dr. Ashok K. Kulkarni for his assistance in compiling the same.

APS Narula

NCL, Poona 8

(A. P. S. NARULA)

February 17, 1975