

List of Figures

Figure	Title of the figure	Page no.
3.1	Map showing Bio-geographic zones of the State West Bengal	28
3.2	Map showing survey sites in respect of altitudinal gradients of West Bengal	31
3.3	Map showing survey sites in respect of vegetation patterns of West Bengal	32
4.1	Family-wise distribution of butterflies in different bio-geographic provinces of West Bengal	36
5.1	Photographs of Papilionid butterflies found in West Bengal. a. <i>Graphium cloanthus</i> (82-85 mm), b. <i>Graphium sarpedon</i> (80-90 mm), c. <i>Graphium doson</i> (70-80 mm), d. <i>Graphium chironides</i> (75-100 mm), e. <i>Graphium agamemnon</i> (85-100 mm), f. <i>Graphium antiphates</i> (80-85 mm), g. <i>Graphium eurous</i> (60-70 mm), h. <i>Graphium nomius</i> (75-90 mm), i. <i>Graphium aristeus</i> (70-80 mm), j. <i>Graphium macareus</i> (80-100 mm)	69
5.2	Photographs of Papilionid butterflies found in West Bengal. a. <i>Graphium xenocles</i> (85-120 mm), b. <i>Papilio agestor</i> (83-120 mm), c. <i>Papilio epycides</i> (70-90 mm), d. <i>Papilio clytia</i> f. <i>clytia</i> (90-100 mm), e. <i>Papilio clytia</i> f. <i>dissimilis</i> (90-100 mm), f. <i>Papilio polytes</i> female (90-100 mm), g. <i>Papilio polytes</i> male (90-100 mm), h. <i>Papilio castor</i> female (100-130 mm), i. <i>Papilio castor</i> male (100-130 mm), j. <i>Papilio helenus</i> (110-130 mm)	70
5.3	Photographs of Papilionid butterflies found in West Bengal. a. <i>Papilio nephelus</i> (115-130 mm), b. <i>Papilio memnon</i> female f. <i>butlerianus</i> (120-150 mm), c. <i>Papilio memnon</i> male f. <i>agenor</i> (120-150 mm), d. <i>Papilio memnon</i> male f. <i>agenor</i> (120-150 mm), e. <i>Papilio polymnestor</i> (120-150 mm), f. <i>Papilio protenor</i> (100-130 mm), g. <i>Papilio protenor</i> (100-130 mm), h. <i>Papilio alcmenor</i> (110-130 mm), i. <i>Papilio bootes</i> (110-120 mm), j. <i>Papilio demoleus</i> (80-100 mm)	71
5.4	Photographs of Papilionid butterflies found in West Bengal. a. <i>Papilio paris</i> (90-140 mm), b. <i>Papilio paris</i> (90-140 mm), c. <i>Papilio polyctor</i> (90-130 mm), d. <i>Papilio crino</i> (80-100 mm), e. <i>Atrophaneura polyeuctes</i> (100-140 mm), f. <i>Atrophaneura varuna</i> (88-136 mm), g. <i>Atrophaneura aristolochiae</i> (80-110 mm), h. <i>Atrophaneura hector</i> (90-110 mm), i. <i>Troides helena</i> (140-170 mm), j. <i>Troides helena</i> (140-170 mm)	72
5.5	Photographs of Pierid butterflies found in West Bengal. a. <i>Eurema andersoni</i> (38-45 mm), b. <i>Eurema hecabe</i> (40-50 mm), c. <i>Eurema blanda</i> (40-45 mm), d. <i>Eurema brigitta</i> (30-40 mm), e. <i>Eurema laeta</i> (30-45 mm), f. <i>Gandaca harina</i> (35-45 mm), g. <i>Dercas verhuelli</i> (60-70 mm), h. <i>Catopsilia Pomona</i> (55-80 mm), i. <i>Catopsilia pyranthe</i> (50-70 mm), j. <i>Colias fieldii</i> (40-65 mm)	87
5.6	Photographs of Pierid butterflies found in West Bengal. a. <i>Colotis amata</i> (35-50 mm), b. <i>Ixias marianne</i> female (50-55 mm), c. <i>Ixias marianne</i> male (50-55 mm), d. <i>Ixias pyrene</i> (50-70 mm), e. <i>Hebomoia glaucippe</i> (80-100 mm), f. <i>Pareronia valeria</i> female f. <i>philomela</i> (65-80 mm), g. <i>Pareronia valeria</i> female (65-80 mm), h. <i>Pareronia valeria</i> male (65-80 mm), i. <i>Appias libythea</i> female (50-60 mm), j. <i>Appias libythea</i> male (50-60 mm)	88

Figure	Title of the figure	Page no.
5.7	Photographs of Pierid butterflies found in West Bengal. a. <i>Appias lycida</i> (55-70 mm), b. <i>Appias albina</i> (60-75 mm), c. <i>Appias indra</i> (60-70 mm), d. <i>Appias lalage</i> (55-80 mm), e. <i>Pieris canidia</i> (45-60 mm), f. <i>Pontia daplidice</i> (45-50 mm), g. <i>Cepora nerissa</i> (40-65 mm), h. <i>Cepora nerissa</i> (40-65mm), i. <i>Cepora nadina</i> (55-65 mm), j. <i>Prioneris thestylis</i> (70-90 mm)	89
5.8	Photographs of Pierid butterflies found in West Bengal. a. <i>Prioneris clemathe</i> (80-90 mm), b. <i>Delias belladonna</i> (79-100 mm), c. <i>Delias pasithoe</i> (70-90 mm), d. <i>Delias acalis</i> (80-100 mm), e. <i>Delias descombesi</i> (65-90 mm), f. <i>Delias hyparete</i> (72-88 mm), g. <i>Delias eucharis</i> (66-83 mm), h. <i>Leptosia nina</i> (35-50 mm), i. <i>Belenois aurota</i> (40-55 mm), j. <i>Belenois aurota</i> (40-55 mm)	90
5.9	Photographs of Lycaenid butterflies found in West Bengal. a. <i>Poritia hewitsoni</i> (31-38 mm), b. <i>Spalgis epius</i> (20-30 mm), c. <i>Curetis thetis</i> (40-48 mm), d. <i>Curetis bulis</i> (35-45 mm), e. <i>Curetis acuta</i> (35-42 mm), f. <i>Arhopala pseudocentaurus</i> (53-62 mm), g. <i>Arhopala amantes</i> (45-57 mm), h. <i>Arhopala perimuta</i> (30-37 mm), i. <i>Arhopala rama</i> (34-40 mm), j. <i>Arhopala eumolphus</i> (44-50 mm)	130
5.10	Photographs of Lycaenid butterflies found in West Bengal. a. <i>Arhopala eumolphus</i> (44-50 mm), b. <i>Arhopala atrax</i> (34-40 mm), c. <i>Arhopala fulla</i> (36-38 mm), d. <i>Arhopala paramuta</i> (38-44 mm), e. <i>Mahathala ameria</i> (38-42 mm), f. <i>Flos adriana</i> (38-44 mm), g. <i>Flos asoka</i> (40-44 mm), h. <i>Flos chinensis</i> (43-46 mm), i. <i>Surendra quercetorum</i> (30-40 mm), j. <i>Amblypodia anita</i> (45-52 mm)	131
5.11	Photographs of Lycaenid butterflies found in West Bengal. a. <i>Iraota timoleon</i> (40-48 mm), b. <i>Loxura atymnus</i> (36-40 mm), c. <i>Horaga onyx</i> (27-33 mm), d. <i>Rathinda amor</i> (26-28 mm), e. <i>Ticherra acte</i> (34-38 mm), f. <i>Cheritra freja</i> (38-42 mm), g. <i>Dacalana cotys</i> (33-37 mm), h. <i>Creon cleobis</i> (27-38 mm), i. <i>Tajuria jehana</i> (30-37 mm), j. <i>Remelana jangala</i> (32-42 mm)	132
5.12	Photographs of Lycaenid butterflies found in West Bengal. a. <i>Hypolycaena erylus</i> (32-36 mm), b. <i>Zeltus amasa</i> (28-32 mm), c. <i>Chliaria kina</i> (26-29 mm), d. <i>Chliaria othona</i> (24-27 mm), e. <i>Deudorix isocrates</i> (34-50 mm), f. <i>Sinthusa nasaka</i> (28-32 mm), g. <i>Sinthusa chandrana</i> (28-32 mm), h. <i>Rapala pheretima</i> (38-42 mm), i. <i>Rapala iarbus</i> (33-41 mm), j. <i>Rapala manea</i> (30-33 mm)	133
5.13	Photographs of Lycaenid butterflies found in West Bengal. a. <i>Rapala varuna</i> (28-29 mm), b. <i>Rapala buxaria</i> (38-42 mm), c. <i>Rapala tara</i> (33-41 mm), d. <i>Rapala nissa</i> (34-38 mm), e. <i>Catapaecilma elegans</i> (28-32 mm), f. <i>Acupicta lohita</i> (30-42 mm), g. <i>Acupicta syama</i> (26-40 mm), h. <i>Cigaritis vulcanus</i> (26-34 mm), i. <i>Cigaritis ictis</i> (27-35 mm), j. <i>Spindasis nipalicus</i> (36-42 mm)	134
5.14	Photographs of Lycaenid butterflies found in West Bengal. a. <i>Heliophorus epicles</i> (30-36 mm), b. <i>Anthene emolus</i> (28-35 mm), c. <i>Anthene lycaenina</i> (24-29 mm), d. <i>Caleta elna</i> (30-34 mm), e. <i>Castalius rosimon</i> (24-34 mm), f. <i>Tarucus nara</i> (23-28 mm), g. <i>Tarucus ananda</i> (22-28 mm), h. <i>Talicauda nyseus</i> (30-36 mm), i. <i>Leptotes plinius</i> (22-30 mm), j. <i>Petrelaea dana</i> (24-28 mm)	135

Figure	Title of the figure	Page no.
5.15	Photographs of Lycaenid butterflies found in West Bengal. a. <i>Nacaduba kurava</i> (30-38 mm), b. <i>Nacaduba beroe</i> (28-34 mm), c. <i>Nacaduba berenice plumbeomicans</i> (22-30 mm), d. <i>Prosotas aluta coelestis</i> (23-25 mm), e. <i>Prosotas nora</i> (18-25 mm), f. <i>Prosotas dubiosa indica</i> (22-26 mm), g. <i>Prosotas bhutea</i> (25-28 mm), h. <i>Jamides bochus</i> (25-34 mm), i. <i>Jamides celeno</i> (27-40 mm), j. <i>Jamides alecto</i> (30-44 mm)	136
5.16	Photographs of Lycaenid butterflies found in West Bengal. a. <i>Catochrysops panormus</i> (25-35 mm), b. <i>Catochrysops strabo</i> (25-35 mm), c. <i>Lampides boeticus</i> (24-36 mm), d. <i>Zizzeria karsandra</i> (18-24 mm), e. <i>Pseudozizeeria maha</i> (26-30 mm), f. <i>Zizina otis</i> (19-26 mm), g. <i>Zizula hylax</i> (16-24 mm), h. <i>Chilades putli</i> (15-22 mm), i. <i>Azanus ubaldus</i> (20-25 mm), j. <i>Megisba malaya</i> (19-30 mm)	137
5.17	Photographs of Lycaenid butterflies found in West Bengal. a. <i>Neopithecops zalmora</i> (16-30 mm), b. <i>Euchrysops cnejus</i> (25-33 mm), c. <i>Chilades lajus</i> dry-season-form (26-30 mm), d. <i>Chilades lajus</i> (26-30 mm), e. <i>Everes lacturnus</i> (22-28 mm), f. <i>Chilades pandava</i> (25-35 mm), g. <i>Chilades parrhasius</i> (22-26 mm), h. <i>Acytolepis puspa</i> (28-35 mm), i. <i>Celastrina lavendularis</i> (28-34 mm), j. <i>Udara albocaerulea</i> (26-34 mm)	138
5.18	Photographs of Riodinid butterflies found in West Bengal. a. <i>Abisara fylla</i> female (50-60 mm), b. <i>Abisara fylla</i> male (50-60 mm), c. <i>Abisara echerius</i> (40-50 mm), d. <i>Abisara echerius</i> (40-50 mm), e. <i>Zemeros flegyas</i> (35-40 mm), f. <i>Zemeros flegyas</i> (35-40 mm)	141
5.19	Photographs of Nymphalid butterflies found in West Bengal. a. <i>Libythea lepita</i> (45-50 mm), b. <i>Libythea myrrha</i> (45-55 mm), c. <i>Tirumala limniace</i> (90-100 mm), d. <i>Tirumala septentrionis</i> (75-95 mm), e. <i>Danaus genutia</i> (72-100 mm), f. <i>Danaus chrysippus</i> (70-80 mm), g. <i>Parantica aglea</i> (70-85 mm), h. <i>Parantica sita</i> (85-110 mm), i. <i>Parantica melaneus</i> (85-95 mm), j. <i>Euploea sylvester</i> (95-105 mm)	189
5.20	Photographs of Nymphalid butterflies found in West Bengal. a. <i>Euploea mulciber</i> female (90-100 mm), b. <i>Euploea mulciber</i> male (90-100 mm), c. <i>Euploea algea</i> (95-100 mm), d. <i>Euploea radamanthus</i> (80-90 mm), e. <i>Euploea klugii kollari</i> (85-100 mm), f. <i>Euploea core</i> (85-95 mm), g. <i>Polyura athamas</i> (60-75 mm), h. <i>Polyura dolon</i> (80-105 mm), i. <i>Polyura eudamippus</i> (100-120 mm), j. <i>Charaxes bernardus</i> male f. <i>corax</i> (85-110 mm)	190
5.21	Photographs of Nymphalid butterflies found in West Bengal. a. <i>Charaxes bernardus</i> (85-110 mm), b. <i>Charaxes aristogiton</i> (70-90 mm), c. <i>Charaxes kahruba</i> (65-70 mm), d. <i>Charaxes marmax</i> (90-120 mm), e. <i>Charaxes solon</i> (70-80 mm), f. <i>Discophora sondaica</i> (80-90 mm), g. <i>Thaumantis diores</i> (95-115 mm), h. <i>Melanitis leda</i> (60-80 mm), i. <i>Melanitis leda</i> (60-80 mm), j. <i>Lethe europa</i> (65-75 mm)	191
5.22	Photographs of Nymphalid butterflies found in West Bengal. a. <i>Lethe confusa</i> (50-60 mm), b. <i>Lethe verma</i> (55-60 mm), c. <i>Elymnias hypermnestra undularis</i> (60-80 mm), d. <i>Elymnias vasudeva</i> (80-90 mm), e. <i>Elymnias malelas</i> (75-85 mm), f. <i>Mycalesis anaxias</i> (48-55 mm), g. <i>Mycalesis perseus</i> (38-55 mm), h. <i>Mycalesis mineus</i> (40-50 mm), i. <i>Mycalesis visala</i> (45-55 mm), j. <i>Mycalesis malsara</i> (40-45 mm)	192

Figure	Title of the figure	Page no.
5.23	Photographs of Nymphalid butterflies found in West Bengal. a. <i>Orsotriaena medus</i> (45-55 mm), b. <i>Ypthima sakra</i> (45-55 mm), c. <i>Ypthima nareda</i> (40-50 mm), d. <i>Ypthima nareda</i> (40-50 mm), e. <i>Ypthima huebneri</i> (30-40 mm), f. <i>Ypthima baldus</i> (32-48 mm), g. <i>Acraea violae</i> (50-65 mm), h. <i>Acraea violae</i> (50-65 mm), i. <i>Acraea issoria</i> (45-85 mm), j. <i>Cethosia biblis</i> (65-90 mm)	193
5.24	Photographs of Nymphalid butterflies found in West Bengal. a. <i>Cethosia biblis</i> (65-90 mm), b. <i>Cethosia cyane</i> (80-95 mm), c. <i>Cethosia cyane</i> (80-95 mm), d. <i>Vindula erota</i> (72-110 mm), e. <i>Vindula erota</i> female (72-110 mm), f. <i>Vindula erota</i> male (72-110 mm), g. <i>Cirrochroa aoris</i> female (80-90 mm), h. <i>Cirrochroa aoris</i> male (80-90 mm), i. <i>Cirrochroa tyche</i> (65-75 mm), j. <i>Vagrans egista</i> (55-95 mm)	194
5.25	Photographs of Nymphalid butterflies found in West Bengal. a. <i>Phalanta phalantha</i> (50-60 mm), b. <i>Moduza procris</i> (60-75 mm), c. <i>Auzakia danava</i> (80-85 mm), d. <i>Athyma perius</i> (60-70 mm), e. <i>Athyma ranga</i> (60-70 mm), f. <i>Athyma selenophora</i> female (55-75 mm), g. <i>Athyma selenophora</i> male (55-75 mm), h. <i>Athyma cama</i> female (60-75 mm), i. <i>Athyma cama</i> male (60-75 mm), j. <i>Athyma nefte</i> female (55-70 mm)	195
5.26	Photographs of Nymphalid butterflies found in West Bengal. a. <i>Athyma nefte</i> male (55-70 mm), b. <i>Pantoporia hordonia</i> (45-50 mm), c. <i>Lasippa viraja</i> (55-75 mm), d. <i>Neptis miah</i> (45-60 mm), e. <i>Neptis ananta</i> (50-70 mm), f. <i>Neptis hylas</i> (50-60 mm), g. <i>Neptis jumbah</i> (60-70 mm), h. <i>Phaedyra columella</i> (60-70 mm), i. <i>Neptis clinia</i> (45-60 mm), j. <i>Neptis soma</i> (50-70 mm)	196
5.27	Photographs of Nymphalid butterflies found in West Bengal. a. <i>Neptis magadha</i> (55-60 mm), b. <i>Neptis yerburyi</i> (58-70 mm), c. <i>Neptis nashona</i> (65-70 mm), d. <i>Neptis vikasi</i> (60-70 mm), e. <i>Neptis harita</i> (55-60 mm), f. <i>Lebadea martha</i> (62-75 mm), g. <i>Euthalia monina kesava</i> (50-80 mm), h. <i>Euthalia phemius</i> female (65-70 mm), i. <i>Euthalia phemius</i> male (65-70 mm), j. <i>Euthalia aconthea</i> (55-80 mm)	197
5.28	Photographs of Nymphalid butterflies found in West Bengal. a. <i>Euthalia lubentina</i> (60-80 mm), b. <i>Euthalia nais</i> (60-70 mm), c. <i>Tanaecia lepidea</i> (65-80 mm), d. <i>Tanaecia julii</i> (65-85 mm), e. <i>Tanaecia jahnu</i> (65-80 mm), f. <i>Cyrestis thyodamas</i> (50-60 mm), g. <i>Chersonesia risa</i> (40-45 mm), h. <i>Pseudergolis wedah</i> (55-65 mm), i. <i>Dichorhagia nesimachus</i> (65-85 mm), j. <i>Stibochiona nicea</i> (60-80 mm)	198
5.29	Photographs of Nymphalid butterflies found in West Bengal. a. <i>Ariadne ariadne</i> (45-60 mm), b. <i>Ariadne merione</i> (45-60 mm), c. <i>Apatura chevana</i> (75-80 mm), d. <i>Apatura ambica</i> (65-90 mm), e. <i>Apatura ambica</i> (65-90 mm), f. <i>Rohana parisatis</i> female (40-50 mm), g. <i>Rohana parisatis</i> male (40-50 mm), h. <i>Euripus nyctelius</i> (65-85 mm), i. <i>Hestina nama</i> (95-105 mm), j. <i>Sephisa chandra</i> (75-90 mm)	199
5.30	Photographs of Nymphalid butterflies found in West Bengal. a. <i>Symbrenthia hippoclus</i> (45-55 mm), b. <i>Vanessa indica</i> (55-65 mm), c. <i>Vanessa cardui</i> (55-70 mm), d. <i>Nymphalis caschmirensis</i> (55-65 mm), e. <i>Junonia orithya</i> (45-60 mm), f. <i>Junonia hierta</i> female (45-60 mm), g. <i>Junonia hierta</i> male (45-60 mm), h. <i>Junonia iphita</i> (55-80 mm), i. <i>Junonia lemonias</i> (40-60 mm), j. <i>Junonia almana</i> (60-65 mm)	200

Figure	Title of the figure	Page no.
5.31	Photographs of Nymphalid butterflies found in West Bengal. a. <i>Junonia atlites</i> (55-65 mm), b. <i>Hypolimnas bolina</i> female (70-110 mm), c. <i>Hypolimnas bolina</i> male (70-110 mm), d. <i>Hypolimnas misippus</i> female (70-85 mm), e. <i>Hypolimnas misippus</i> male (70-85 mm), f. <i>Kallima inachus</i> (85-110 mm), g. <i>Kallima inachus</i> (85-110 mm), h. <i>Doleschallia bisaltide</i> (75-85 mm), i. <i>Doleschallia bisaltide</i> (75-85 mm), j. <i>Calinaga buddha</i> (90-105 mm)	201
5.32	Photographs of Hesperid butterflies found in West Bengal. a. <i>Bibasis jaina</i> (60-70 mm), b. <i>Bibasis sena</i> (42-50 mm), c. <i>Hasora anura</i> (50-55 mm), d. <i>Hasora chromus</i> (45-50 mm), e. <i>Hasora badra</i> (50-55 mm), f. <i>Badamia exclamationis</i> (50-55 mm), g. <i>Choaspes benjaminii</i> (50-60 mm), h. <i>Celaenorrhinus leucocera</i> (45-55 mm), i. <i>Spialia galba</i> (20-27 mm), j. <i>Spialia galba</i> (20-27 mm)	225
5.33	Photographs of Hesperid butterflies found in West Bengal. a. <i>Sarangesa dasahara</i> (26-35 mm), b. <i>Sarangesa dasahara</i> (26-35 mm), c. <i>Pseudocoladenia dan</i> (40-46 mm), d. <i>Coladenia indrani</i> (40-46 mm), e. <i>Coladenia agni</i> (40 mm), f. <i>Coladenia agni</i> (40 mm), g. <i>Gerosis bhagava</i> (35-45 mm), h. <i>Gerosis phisara</i> (35-45 mm), i. <i>Tagiades gana</i> (45-50 mm), j. <i>Tagiades japetus ravi</i> (45-50 mm)	226
5.34	Photographs of Hesperid butterflies found in West Bengal. a. <i>Tagiades litigiosa litigiosa</i> (37-44 mm), b. <i>Tagiades menaka menaka</i> (37-44 mm), c. <i>Odontoptilum angulata</i> (40-45 mm), d. <i>Odontoptilum angulata</i> (40-45 mm), e. <i>Caprona ransonnetii</i> (35-45 mm), f. <i>Caprona ransonnetii</i> (35-45 mm), g. <i>Ctenoptilum vasava</i> (35-40 mm), h. <i>Mooreana trichoneura</i> (40-45 mm), i. <i>Mooreana trichoneura</i> (40-45 mm), j. <i>Taractrocera maevis</i> (22-28 mm)	227
5.35	Photographs of Hesperid butterflies found in West Bengal. a. <i>Oriens goloides</i> (22-28 mm), b. <i>Telicota bambusae</i> (33-36 mm), c. <i>Telicota colon</i> (32-36 mm), d. <i>Cephrenes chrysozona oceanica</i> (46 mm), e. <i>Parnara guttatus</i> (32-38 mm), f. <i>Borbo cinnara</i> (30-36 mm), g. <i>Pelopidas mathias</i> (32-38 mm), h. <i>Pelopidas subochracea</i> (38-42 mm), i. <i>Pelopidas assamensis</i> (42-60 mm), j. <i>Pelopidas conjuncta</i> (45-52 mm)	228
5.36	Photographs of Hesperid butterflies found in West Bengal. a. <i>Polytremis eltola</i> (35-45 mm), b. <i>Baoris farri</i> (43-48 mm), c. <i>Caltoris kumara</i> (42-46 mm), d. <i>Sebastomyia dolopia</i> (32-34 mm), e. <i>Halpe zema</i> (35 mm), f. <i>Halpe porus</i> (32 mm), g. <i>Halpe porus</i> (32 mm), h. <i>Arnetta atkinsoni</i> (25-32 mm), i. <i>Suastus gremius</i> (32-45 mm), j. <i>Iambrix salsala</i> (26-30 mm)	229
5.37	Photographs of Hesperid butterflies found in West Bengal. a. <i>Iambrix salsala</i> (26-30 mm), b. <i>Hyarotis adrastus praba</i> (38-48 mm), c. <i>Zographetus ogygia</i> (30-32 mm), d. <i>Gangara thyrsis</i> (70-76 mm), e. <i>Gangara thyrsis</i> (70-76 mm), f. <i>Matapa aria</i> (40-45 mm), g. <i>Erionota thrax</i> (70-76 mm), h. <i>Baracus vittatus</i> (30-34 mm), i. <i>Astictopterus jama olivascens</i> (38-40 mm), j. <i>Ochus subvittatus</i> (22-25 mm)	230
5.38	Photographs of Hesperid butterflies found in West Bengal. a. <i>Ampittia discorides</i> (22-28 mm), b. <i>Ampittia discorides</i> (22-28 mm), c. <i>Psolos fuligo</i> (30-40 mm), d. <i>Ancistroides nigrita</i> (48-60 mm), e. <i>Notocrypta curvifascia</i> (38-50 mm), f. <i>Udaspes folus</i> (40-48 mm), g. <i>Udaspes folus</i> (40-48 mm), h. <i>Aeromachus pygmaeus</i> (20-22 mm), i. <i>Aeromachus pygmaeus</i> (20-22 mm), j. <i>Lotongus sarala</i> (42-44 mm)	231

Figure	Title of the figure	Page no.
5.39	Map of Gorumara National Park, Jalpaiguri district, West Bengal	235
5.40	Variation in the abundance (Mean \pm S.E.) of different species of butterflies across pre-monsoon (A), monsoon (B), post-monsoon (C), and winter (D) seasons in varied habitat types observed in Gorumara National Park during the study period (January 2010 - December 2011). Species present in the respective seasons and habitats are only represented.	261
6.1	Life stages of <i>Appias libythea</i> . a. Adult male, b. Adult female, c. Egg d. Egg immediately before hatching, e. Instar I, f. Instar II, g. Instar III, h. Instar IV, i. Instar V, j. Pupa, k. Pupa before emergence	271
6.2	Life stages of <i>Hyarotis adrastus praba</i> . a. Adult under-wing, b. Adult upper-wing, c. Egg d. Egg immediately before hatching, e. Instar I, f. Instar II, g. Instar III, h. Instar IV, i. Instar V, j. Pupa, k. Pupa before emergence	276
7.1	Natural predators affecting various life-cycle stage of butterflies. a. Infected larva (parasitoid), b. Larva infected by parasitoid (fly) cocoon with lava, c. Ants feeding on larva, d. Spider feeding on larva, e. Parasitoid (fly) emerging from pupa, f. Rubber fly feeding on adult butterfly, g. Praying mantis feeding on adult butterfly.	281