

REFERENCES

1. Abdel-Fattah G. M. and Al-Amri S. M. (2012). Induced systemic resistance in tomato plants against *Fusarium oxysporum* f. sp. *lycopersici* by different kinds of compost. *African Journal of Biotechnology*. 11:12454-12463.
2. Abdel-Fattah G.M., Al-Amri S.M.(2012). Induced systemic resistance in tomato plants against *Fusarium oxysporum* f.sp. *lycopersici* by different kinds of compost. *African Journal of Biotechnology*. 11(61):12454-12463.
3. Aboellil A.H. (2007). Trilogy, a product of neem (*Azadirachta indica*) induces resistance in Cucumber against *Podosphaera xanthii*. *Research Journal of Microbiology* .2:402-414.
4. Abo-Elyousr K.A., Hussein M.A.M., Allam A.D.A., Hassan M.H.(2009). Salicylic acid induced systemic resistance on onion plants against *Stemphylium vesicarium*. *Archives of Phytopathology and Plant Protection* . 42(11): 1042-1050.
5. Adam A., Ongena M., Duby F., Dommes J., Thonart P.(2008). Systemic resistance and lipoxygenase-related defence response induced in tomato by *Pseudomonas putida* strain BTP1. *BMC Plant Biology*. 8:113. doi:10.1186/1471-2229-8-113.
6. Agostini J.P., Timmer L.W. (1992). Selective isolation procedures for differentiation of two strains of *Colletotrichum gloeosporioides* from Citrus. *The American Phytopathological Society, Plant Disease*; 76(11): 1176-1178.
7. Agrios G.N. (2005). *Plant pathology*. Fifth edition. Elsevier Academic press, London.
8. Akimoto K., Kitagawa Y., Akamatsu T., Hirose N., Sugano M., Shimizu S., and Yamada H. (1993). Protective effects of sesamin against liver damage caused by alcohol or carbon tetrachloride in rodents. *Annals of nutrition and metabolism* 37(4):218-224.
9. Akinbode O. A. and Ikotun T. (2011). Potential of two *Tricoderma* species as antagonistic agents against *Colletotrichum destructivum* of cowpea. *African Journal of Microbiology Research*. 5(5): 551-554.
10. Akram W. and Anjum T. (2011). Quantitative Changes in Defense System of Tomato Induced by two Strains of *Bacillus* against *Fusarium* wilt. *Indian Horticulture Journal*. 1(1):14-18.
11. Akter S., Kadir J., Juraimi A. S., Saud H. M. and Elmahdi S. (2014). Isolation and identification of antagonistic bacteria from phylloplane of rice as biocontrol agents for sheath blight, *Journal of Environmental Biology*. 35(6): 1095.

REFERENCES

12. Alam S.S., Sakamoto K., Amemiya Y., Inubushi K. (2010). Biocontrol of soil-borne Fusarium wilts of tomato and cabbage with a root colonizing fungus, *Penicillium* sp. EU0013. Paper presented at: 19th World Congress of Soil Science, Soil Solutions for a Changing World; Brisbane, Australia
13. Alamri S.A. (2008). Epiphytic Microflora on the leaves of *Juniperus procera* from aseer region, Saudi Arabia. *Journal of Biological Sciences*. 8(5): 24-27.
14. Alemu F. and Alemu T. (2013). Antifungal activity of secondary metabolites of *Pseudomonas fluorescens* isolates as a biocontrol agent of chocolate spot disease (*Botrytis fabae*) of faba bean in Ethiopia. *African Journal of Microbiology*. 7 (47): 5364-5373.
15. Alhubaishi A.A., Abdel-Kader M.I. (1991). Phyllosphere and phylloplane fungi of qat in Sana'a, Yemen Arab Republic. *Journal of Basic Microbiology*. 31(2):83-9.
16. Amme S., Rutten T., Melzer M., Sonsmann G., Vissers J.P., Schlesier B. and Mock, H.P.(2005). A proteome approach defines protective functions of tobacco leaf trichomes. *Proteomics*.5: 2508–2518.
17. Anahosur K.H., Hegde R.K., Patil S.H. (1985). Role of sugars and phenols in the charcoal rot resistance of sorghum. *Phytopathologische Zeitschrift*. 113:30-35.
18. Andrews J. H. and Hirano S. S. (Eds.). (2012). *Microbial ecology of leaves*. Springer Science & Business Media.
19. Andrews J. H. and Harris R. F. (2000). The ecology and biogeography of microorganisms on plant surfaces. *Annual review of phytopathology*. 38(1): 145-180.
20. Anfoka G., Buchenauer H. (1997). Systemic acquired resistance in tomato against *Phytophthora infestans* by pre-inoculation with tobacco necrosis virus. *Physiological and Molecular Plant Pathology*. 50(2):85-101.
21. Arnold A. E. and Herre E. A. (2003). Canopy cover and leaf age affect colonization by tropical fungal endophytes: ecological pattern and process in *Theobroma cacao* (Malvaceae). *Mycologia*. 95(3): 388-398.
22. Arun K, Mali PC, Manga VK. (2010). Changes of some phenolic compounds and enzyme activities on infected pearl millet caused by *Sclerospora graminicola*. *International Journal of Plant Physiology and Biochemistry*. 2(1):6-10.
23. Ascensão L., Marques N., Pais M.S.(1995). Glandular trichomes on vegetative and reproductive organs of *Leonotis leonurus* (Lamiaceae) *Annals of Botany*.75:619–626.

24. Asha A.N., Kannabiran B.(2001). Effect of *Datura metel* leaf extract on the enzymatic and nucleic acid changes in the chilli seedlings infected with *Colletotrichumcapsici*. Indian Phytopathology. 54:373-375.
25. Austin B., Dickinson C. H. and Goodfellow M. (1977). Antagonistic interactions of phylloplane bacteria with *Drechslera dictyoides* (Drechsler) Shoemaker. Canadian journal of microbiology. 23(6):710-715.
26. Baker C. J., Mock N. M., Whitaker B. D., Hammond R. W., Nemchinov L., Roberts D. P. and Aver'yanov A. A. (2014). Characterization of apoplast phenolics: in vitro oxidation of acetosyringone results in a rapid and prolonged increase in the redox potential. Physiological and Molecular Plant Pathology. 86: 57-63.
27. Balestra G.M., Heydari A., Ceccarelli D., Ovidi E., Quattrucci A. (2009). Antibacterial effect of *Allium sativum* and *Ficus carica* extracts on tomato bacterial pathogens. Crop Protection. 28(10):807-811.
28. Barak J. D., Kramer L. C. and Hao L. Y. (2011). Colonization of tomato plants by *Salmonella enterica* is cultivar dependent, and type 1 trichomes are preferred colonization sites. Applied and environmental microbiology. 77(2): 498-504.
29. Baruzzi F., Quintieri L., Morea M. and Caputo L. (2011). Antimicrobial compounds produced by *Bacillus* spp. and applications in food. Science against microbial pathogens: communicating current research and technological advances. 2:1102-1111.
30. Bashan Y. and de-Bashan L. E. (2002). Protection of tomato seedlings against infection by *Pseudomonas syringae* pv. tomato by using the plant growth-promoting bacterium *Azospirillum brasilense*. Applied and Environmental Microbiology. 68(6): 2637-2643.
31. Beattie G. A., and Lindow S. E. (1999). Bacterial colonization of leaves: a spectrum of strategies. Phytopathology 89:353-359
32. Bednarek P., and Osbourn A. (2009). Plant-Microbe interactions: chemical diversity in plant defense. Science. 324: 746–748.
33. Besser K., Harper A., Welsby N., Schauvinhold I., Slocombe S., Li Y., ... and Broun P. (2009). Divergent regulation of terpenoid metabolism in the trichomes of wild and cultivated tomato species. Plant Physiology. 149(1): 499-514.
34. Bhatia I.S., Uppal D.S. and Bajaj K.C. (1972). Study of phenolic contents of resistant and susceptible varieties of tomato *Lycopersicum esculantum* in relation to early blight disease. Indian Phytopathology. 25:231-235.
35. Bhattacharya K. and Warde W. B. (1986). Expression of gene-specific and age-related resistance and the accumulation of glyceollin in soybean leaves infected with

REFERENCES

- Phytophthora megasperma* f. sp. *glycinea*. Physiological and Molecular Plant Pathology. 29: 105 – 113
36. Bhattacharyya M.K., Warde W.B. (1988). Phenylalanine ammonia lyase activity in soybean hypocotyls and leaves following infection with *Phytophthora megasperma* f. sp. *glycinea*. Canadian Journal of Botany. 66:18-23.
 37. Bhullar B.S., Bajaj K.L. and Bhatia .I.S. (1972). Studies on phenols of the resistant and susceptible varieties of chillies in relation to anthracnose disease. Phytopathology Zeitschrift. 75: 236-240.
 38. Blakeman J. P. and Fokkema N. J. (1982). Potential for biological control of plant diseases on the phylloplane. Annual review of phytopathology. 20(1): 167-190.
 39. Blakeman J. P. & Atkinson P. (1981) Antimicrobial substances associated with the aerial surfaces of plants. In: Microbial Ecology of the Phylloplane (Ed. by J. P.Blakeman), Academic Press, London.pp .245–263.
 40. Blakeman J. P. (1985) Ecological succession of leaf surface microorganisms in relation to biological control. In: Biological Control of the Phylloplane (Ed. by C. E.Windels & S. E.Lindow). The American Phytopathological Society, Minnesota . pp. 6–30.
 41. Blakeman J.P. and Fokkema N.J. (1982). Potential for biological control of plant diseases on the phylloplane, Ann Rev Phytopathol. 58: 464-466.
 42. Blount J.W., Dixon R.A., Paiva N.L. (1992).Stress responses in alfalfa (*Medicago sativa* L.) XVI. Antifungal activity of medicarpin and its biosynthetic precursors; implications for the genetic manipulation of stress metabolites. Physiol. Mol. Plant Pathol, 41: 333–349.
 43. Bodenhausen N., Horton M. W. and Bergelson J. (2013). Bacterial communities associated with the leaves and the roots of *Arabidopsis thaliana*. *PLoS One*. 8(2): e56329.
 44. Boller T.(1989). Primary signals and second messengers in the reaction of plants to pathogens. Boss WF, Morre DJ, Second messengers in plant growth and development, (eds) Alan R Liss, New York, pp: 227-255
 45. Boonchitsirikul C., Wasano K., Nose A.(1998). Activities of phenylalanine ammonia lyase (PAL) and tyrosine ammonia lyase (TAL) in young rice panicles inoculated with *Pyricularia grisea*. Japanese Journal of Tropical Agriculture. 42(1):39-45.
 46. Brandl M. T., Quinones B. and Lindow S. E. (2001). Heterogeneous transcription of an indoleacetic acid biosynthetic gene in *Erwinia herbicola* on plant surfaces. Proc. Natl. Acad. Sci. USA. 98:3454-3459.

47. Brandl M.T., Amundson R. (2008). Leaf Age as a Risk Factor in Contamination of Lettuce with *Escherichia coli* O157:H7 and *Salmonella enterica*, *Appl. Environ. Microbiol.* 74(8):2298-2306.
48. Carder P (2010). Microbial communities of spinach at various stages of plant growth from seed to maturity. *Cassava (Manihot esculenta) suspension cells and leaves.* *Ann Bot.* 94(1): 87-97.
49. Cavalcanti F. R., Resende M. L. V., Ribeiro P. M., Pereira R. B., & Oliveira J. T. A. (2008). Induction of resistance against *Verticillium dahliae* in cacao by a *Crinipellis perniciososa* suspension. *Journal of Plant Pathology.* 273-280.
50. Chagnot C., Zorgani M. A., Astruc T. and Desvaux M. (2013). Proteinaceous determinants of surface colonization in bacteria: bacterial adhesion and biofilm formation from a protein secretion perspective. *Front Microbiol.* 4(303.10): 3389.
51. Chakrabarty P.K., Mukewar P.M., Raj S., Sravan Kumar V. (2002). Biochemical factors governing resistance in diploid cotton against grey mildew. *Indian Phytopathology.* 55(2):140-146.
52. Chauhan D. and Swami D. A. (2014). Studies On Phylloplane Microflora Of Dhak. In *Journal of Emerging Technologies and Innovative Research.*1(7).
53. Chérif M., Arfaoui A. and Rhaïem A.(2007). Phenolic compounds and their role in biocontrol and resistance of chickpea to fungal pathogenic attacks, *Tunisian Journal of Plant Protection.* 2(1): 7-21.
54. Chernin L., Ismailov Z., Haran S., and Chet I. (1995). Chitinolytic *Enterobacter agglomerans* antagonistic to fungal plant pathogens. *Appl Environ Microbiol.* 61(5): 1720-1726.
55. Collinge D. B., Kragh K. M., Mikkelsen J. D., Nielsen K. K., Rasmussen U. and Vad K. (1993). Plant chitinases. *The Plant Journal.* 3(1): 31-40.
56. Colson M., Pupier R. and Perrin A. (1993). Biomathematical analysis of the number of glandular trichomes on leaves of *Mentha× piperita* (Translated from French). *Can J Bot.* 71: 1202-1211.
57. Comolli L. R. (2014). Intra-and inter-species interactions in microbial communities. *Frontiers in microbiology*, 5.
58. Compant,S., Duffy B., Nowak J., Clément C. and Barka E. A. (2005). Use of plant growth-promoting bacteria for biocontrol of plant diseases: principles, mechanisms of action, and future prospects. *Applied and environmental microbiology.*71(9): 4951-4959.
59. Conrath U., Pieterse C. M. and Mauch-Mani B. (2002). Priming in plant–pathogen interactions. *Trends in plant science.* 7(5): 210-216.

REFERENCES

60. Constabel CP and Barbehenn R. (2008). Defensive Roles of Polyphenol Oxidase in Plants. Chapter 12. In: Schaller A (Ed.). Induced Plant Resistance to Herbivory. New York: Springer Verlag. 253-269.
61. Constabel C.P., Bergey D.R., Ryan C.A. (1995). Systemin activates synthesis of wound inducible tomato leaf polyphenol oxidase via the octadecanoid defense signaling pathway. Proceedings of the National Academy of Sciences United States of America. 92:407-411.
62. Cook D. W., Long P. G., Ganesh S., and Cheah L. H. (1997). Attachment microbes antagonistic against *Botrytis cinerea*—biological control and scanning electron microscope studies in vivo. Annals of applied biology. 131(3): 503-518.
63. Cottenye N. (2010) Antimicrobial surfaces based on self-assembled nanoreactors: from block copolymer synthesis to bacterial adhesion study. [Ph.D Thesis] University of Basel, Switzerland.
64. Croteau R., Felton M., Karp F. and Kjonaas R. (1981). Relationship of camphor biosynthesis to leaf development in sage (*Salvia officinalis*). Plant Physiology. 67(4): 820-824.
65. Cruz J., Ortiz C., Guzman F., Fernández-Lafuente R. and Torres R. (2014). Antimicrobial peptides: promising compounds against pathogenic microorganisms. Current medicinal chemistry. 21(20): 2299-2321.
66. Dai G.H., Nicole M., Andary C., Martinez, C.; Bresson, E.; Boher, B.; Daniel, J.F.; Geiger, J.P. Flavonoids accumulate in cell walls, middle lamellae and callose-rich papillae during an incompatible interaction between *Xanthomonas campestris* pv. malvacearum and cotton. Physiol. Mol. Plant Pathol. 49: 285–306.
67. Das S., Aggarwal R., Singh D. V. (2003). Differential induction of defense related enzymes involved in lignin biosynthesis in wheat in response to spot blotch infection. Indian Phytopathology. 56(2): 129-133.
68. De Jager E.S., Wehner F.C., Korsten L. (2001). Microbial ecology of the mango phylloplane. Microbial Ecology .42: 201–207.
69. de Lima Favaro L. C., de Souza Sebastianes F. L. and Araújo W. L. (2012). *Epicoccum nigrum* P16, a sugarcane endophyte, produces antifungal compounds and induces root growth. PLoS One. 7(6): e36826.
70. De Vleeschauwer D., Djavaheri M., Bakker P. A. and Höfte M. (2008). *Pseudomonas fluorescens* WCS374r-induced systemic resistance in rice against *Magnaporthe oryzae* is based on pseudobactin-mediated priming for a salicylic acid-repressible multifaceted defense response. Plant Physiology. 148(4): 1996-2012.

71. De Wit P. J. and Spikman G. (1982). Evidence for the occurrence of race and cultivar-specific elicitors of necrosis in intercellular fluids of compatible interactions of *Cladosporium fulvum* and tomato. *Physiological Plant Pathology*, 21(1): 1-11.
72. De Wit P J., Buurlage M. B., and Hammond K. E. (1986). The occurrence of host-, pathogen-and interaction-specific proteins in the apoplast of *Cladosporium fulvum* (syn. *Fulvia fulva*) infected tomato leaves. *Physiological and Molecular Plant Pathology*. 29(2): 159-172.
73. DeBruyne T., Pieters L., Deelstra H. and Vlietinck A. (1999). Condensed vegetable tannins: biodiversity in structure and biology activities. *Biochemical Systematics and Ecology*. 27:445–459
74. Delaunois B., Jeandet P., Clément C., Baillieul F., Dorey S. and Cordelier S. (2014). Uncovering plant-pathogen crosstalk through apoplastic proteomic studies. *Plant cell wall in pathogenesis, parasitism and symbiosis*, 133.
75. Dias A.C.F., Taketani R.G., Andreote F.D., Luvizotto D.M., da Silva J.L., Nascimento R.D.S., de Melo I.S. (2012). Interspecific variation of the bacterial community structure in the phyllosphere of the three major plant components of mangrove forests, *Braz. J. Microbiol* 43(2):653-660.
76. Dickinson C.H. (1983). Fungal colonization of *Pisum* leaves, *Can. J. Bot.* 45(6): 915-927.
77. Dixon R.A., Paiva N.L. (1985), Stress-induced phenylpropanoid metabolism. *Plant Cell* 7:1085-1097.
78. Doehlemann G. and Hemetsberger C. (2013). Apoplastic immunity and its suppression by filamentous plant pathogens. *New Phytologist*. 198(4): 1001-1016.
79. Dolar F. S. (1997). Determination of the races of *Fusarium oxysporum* f. sp. *ciceris* in Ankara province, Turkey. *Journal of Turkish Phytopathology*. 26(1): 11-15.
80. Dong J., Chen C., Chen Z. (2003). Expression profiles of the *Arabidopsis* WRKY gene superfamily during plant defense response. *Plant Molecular Biology*; 51:21-37.
81. Dutta S., Mishra A. K. and Kumar B. D. (2008). Induction of systemic resistance against fusarial wilt in pigeon pea through interaction of plant growth promoting rhizobacteria and rhizobia. *Soil Biology And Biochemistry*. 40(2): 452-461.
82. Ebel J. (1986). Phytoalexin synthesis: The biochemical analysis of the induction process. *Annual Review of Phytopathology*. 24: 235-264.
83. Eberl H. J. and Demaret L. (2007). A finite difference scheme for a degenerated diffusion equation arising in microbial ecology. *El J Diff Equs CS*. 15: 77-95.

REFERENCES

84. El-Said A. H. (2001). Phyllosphere and phylloplane fungi of banana cultivated in Upper Egypt and their cellulolytic ability. *Mycobiology*.29(4): 210-217.
85. El-Zahaby H.M.(2008). Role of *Pseudomonas fluorescence* in induced systemic resistance for tomato vascular wilt disease caused by *Fusarium oxysporum* f. sp. *lycopersici*. *Journal of Agricultural Science, Mansoura University*; 33:4158.
86. Enya J., Shinohara H., Yoshida S., Tsukiboshi T., Negishi H., Suyama K., and Tsushima S., (2006). Culturable leaf-associated bacteria on Tomato plants and their potential as biological control agents. *Microbial Ecology*. 53 (4):524–536.
87. Ercolani, G. L.(1991). Distribution of epiphytic bacteria on olive leaves and the influence of leaf age and sampling time, *Microbial Ecology*. 21:35-48.
88. Fahmy G.M., Ouf S.A.(1999). Significance of microclimate on phylloplane mycoflora of green and senescing leaves of *Zygophyllum album* L, *Journal of Arid Environments*. 41(3):257-276.
89. Farag M. A., Huhman D. V., Dixon R. A. and Sumner L. W. (2008). Metabolomics reveals novel pathways and differential mechanistic and elicitor-specific responses in phenylpropanoid and isoflavonoid biosynthesis in *Medicago truncatula* cell cultures. *Plant Physiology*. 146: 387–402.
90. Farkas G.L.and Kiraly Z.(1962). Role of phenolic compounds in the physiology of plant disease resistance. *Phytopathology*. 44: 105-150.
91. Fattahi S., Zabihi E., Abedian Z., Pourbagher R., Ardekani A. M., Mostafazadeh A., & Akhavan-Niaki H. (2014). Total phenolic and flavonoid contents of aqueous extract of stinging nettle and in vitro antiproliferative effect on hela and BT-474 Cell lines. *International journal of molecular and cellular medicine*, 3(2), 102.
92. Faust K. and Raes, J. (2012). Microbial interactions: from networks to models. *Nature Reviews Microbiology*. 10(8): 538-550.
93. Favilli F., Messini A. (1990). Nitrogen Fixation at Phyllospheric level in Coniferous Plants in Italy, *Plant and Soil*. 128:91-95.
94. Fayzalla E.A., Abdel-Fattah G.M., Ibrahim A.S., El-Sharkawy H.H.A.(2009). Induction of resistance in tomato plants against *Fusarium oxysporum* f.sp. *lycopersici* by vesicular arbuscular mycorrhizae (VAM). *Journal of Agricultural Science Mansoura University*. 34:9787-9799.
95. Fernández L., Agaras B., Zalba P., Wall L.G., and Valverde C., (2012). *Pseudomonas* spp. isolates with high phosphate-mobilizing potential and root colonization properties from agricultural bulk soils under no-till management. *Biology and Fertility of Soils*.48 (7): 763-773.

-
96. Ferraz H. G. M., Resende R. S., Silveira P. R., Andrade C. C. L., Milagres E. A., Oliveira J. R. and Rodrigues F. D. Á. (2014). Rhizobacteria induces resistance against *Fusarium* wilt of tomato by increasing the activity of defense enzymes. *Bragantia*. 73(3): 274-283.
 97. Feucht W. and Treutter. D. (1999). The role of flavan-3-ols and proanthocyanidins in plant defence. In: Inderjit S, Dakshini KMM, Foy CL (eds), Principles and practices in plant ecology. CRC Press, Boca Raton. pp 307–338.
 98. Fletcher J., Leach J.E., Eversole K., Tauxe R.(2013). Human pathogens on plants: designing a multidisciplinary strategy for research, *Phytopathology*. 103(4):306-315.
 99. Frändberg E. (1997). Antifungal compounds of chitinolytic bacteria from grain ecosystems. 1401-6249.
 100. Fravel D. R., Connick Jr W. J. and Lewis J. A. (1998). Formulation of microorganisms to control plant diseases. In Formulation of microbial biopesticides (187-202). Springer Netherlands.
 101. Freiberg E.(1998). Microclimatic parameters influencing nitrogen fixation in the phyllosphere in a Costa Rican premontane rain forest, *Oecologia*. 117(1-2):9-18.
 102. Fu F. and Wang Q. (2011). Removal of heavy metal ions from wastewaters: a review. *Journal of environmental management*. 92(3): 407-418.
 103. Gall L. S. (1970). Significance of microbial interactions in control of microbial ecosystems, *Biotechnology and bioengineering*:12(3):333-340.
 104. Gandía-Herrero F., Escribano J. and García-Carmona F. (2005). Betaxanthins as substrates for tyrosinase. An approach to the role of tyrosinase in the biosynthetic pathway of betalains. *Plant physiology*. 138(1): 421-432.
 105. Gaspar T., Penel C., Hagege D., Greppin H.(1991). Peroxidase in Plant Growth, differentiation and Developmental Processes. Chapter 4. In: Biochemical, Molecular and Physiological Aspects of Plant Peroxidases. Lobarzewski J., Greppin H., Pennel C., Gaspar T. (Eds.), Lublin (Poland) and Geneva (Switzerland). pp. 249–280.
 106. Geetha N. P., Amruthesh K. N., Sharathchandra R. G. and Shetty H. S. (2005). Resistance to downy mildew in pearl millet is associated with increased phenylalanine ammonia lyase activity. *Functional Plant Biology*. 32(3): 267-275.
 107. Gertzen R. and Escobar M. A. (2014). Assays of polyphenol oxidase activity in walnut leaf tissue. *Bio-protocol*, 4, e1213.
 108. Gherbawy Y., Elhariry H., Altalhi A., El-Deeb B. and Khiralla G. (2012). Molecular screening of *Streptomyces* isolates for antifungal activity and family 19 chitinase enzymes. *Journal of Microbiology*. 50(3): 459-468.

REFERENCES

109. Girish N. and Umesha S. (2005). Effect of plant growth promoting rhizobacteria on bacterial canker of tomato. *Archives of Phytopathology and Plant protection*. 38(3): 235-243.
110. Gonzáles W. L., Negritto M. A., Suárez L. H. and Gianoli E. (2008). Induction of glandular and non-glandular trichomes by damage in leaves of *Madia sativa* under contrasting water regimes. *Acta oecologica*. 33(1): 128-132.
111. Gómez-Vásquez R., Day R., Buschmann H., Randles S., Beeching J.R., Cooper R.M. (2004). Phenylpropanoids, phenylalanine ammonia lyase and peroxidases in elicitor-challenged cassava (*Manihot esculenta*) suspension cells and leaves. *Annals of Botany*. 94(1): 87-97.
112. Gonzalez M., Pujol M., Metraux J.P., Gonzalez-Garcia V. I. C. E. N. T. E., Bolton M.D. and Borrás-Hidalgo O. R. L. A. N. D. O. (2011). Tobacco leaf spot and root rot caused by *Rhizoctonia solani* Kühn. *Molecular Plant Pathology*. 12 (3): 209-216.
113. Gonzalez-Aguilar G.A., Tizuado-Hernandez M.E., Zavaleeta-Gatica R., Martinez-Tellez M.A. (2004). Methyl jasmonate treatments reduce chilling injury and activate the defense response of guava fruits. *Biochemical and Biophysical Research Communication*. 313:694–701.
114. Gould K.S. and Lister C.(2006). Flavonoid functions in plants. In: Andersen, Ø.M. and Markham, K.R (Eds), *Flavonoids chemistry, biochemistry and applications*. Boca Raton: CRC. pp. 397–442.
115. Guleria S. and Kumar A. (2006). Antifungal activity of some Himalayan medicinal plants using direct bioautography. *Journal of cell and molecular Biology*. 5: 95-98
116. Gupta S.K., Kaushik C.D.(2002). Metabolic changes in mustard leaf and siliqua wall due to the infection of *Alternaria* blight (*Alternaria brassicae*). *Cruciferae Newsletter*. 24:85-86.
117. Hahlbrock K. and Scheel D. (1989). Physiology and molecular biology of phenylpropanoid metabolism. *Annual review of plant biology*. 40(1): 347-369.
118. Halfeld-Vieira B. D. A., Vieira Júnior J. R., Romeiro R. D. S., Silva H. S. A. and Baracat-Pereira M. C. (2006). Induction of systemic resistance in tomato by the autochthonous phylloplane resident *Bacillus cereus*, *Pesquisa Agropecuária Brasileira*. 41(8): 1247-1252.
119. Halfeld-Vieira B.D.A., Vieira Júnior J.R., Romeiro R.D.S., Silva H.S.A., and Baracat-Pereira M.C. (2008). Induction of systemic resistance in tomato by the autochthonous phylloplane resident *Bacillus cereus*. *Pesquisa Agropecuaria Brasileira*. 41 (8):1247-1252.

-
120. Hamid R., Khan M. A., Ahmad M., Ahmad M. M., Abdin M. Z., Musarrat J. and Javed S. (2013). Chitinases: an update. *Journal of Pharmacy and Bioallied Sciences*. 5(1): 21.
 121. Harborne J.B., Williams C.A.(2000). Advances in flavonoid research since 1992. *Phytochemistry*, 55, 481–504.
 122. Harish S., Kavino M., Kumar N., Balasubramanian P. and Samiyappan R. (2009). Induction of defense-related proteins by mixtures of plant growth promoting endophytic bacteria against Banana bunchy top virus. *Biological Control*. 51(1):16-25.
 123. Harman GE, Howell CR, Viterbo A, Chet I, Lorito M. (2004). *Trichoderma* species -opportunistic, avirulent plant symbionts. *Nature Reviews Microbiology*. 2:43-56.
 124. Herger G., Klingauf F. (1990). Control of powdery mildew fungi with extracts of the giant knotweed, *Reynoutria sachalinensis* (Polygonaceae). *Mededelingen van de Faculteit landbouwwetenschappen, Rijksuniversiteit-Gent*. 55(3a):1007-1014.
 125. Hindumathy C. K.(2012) .The defense activator from yeast for rapid induction of resistance in susceptible pearl millet hybrid against downy mildew disease. *International Journal of Agriculture Sciences*. 4(2):196-201.
 126. Hiraga S., Sasaki K., Ito H., Ohashi Y. and Matsui H. (2001). A large family of class III plant peroxidases. *Plant and Cell Physiology*. 42(5): 462-468.
 127. Hirano S., Upper C., (2000). Bacteria in the Leaf Ecosystem with Emphasis on *Pseudomonas syringae*—a Pathogen, Ice Nucleus, and Epiphyte. *Microbiology and Molecular Biology Reviews*. 64(3): 624–653.
 128. Ibrahim Y.E.(2012). Activities of antioxidants enzymes in Salicylic acid treated tomato against *Xanthomonas vesicatoria*. *African Journal of Microbiology Research*. 6(27): 5678.
 129. Ilori M. O., Amund O. O., Ezeani C. J., Omoijiahina S., and Adebusoye S. A. (2006). Occurrence and growth potentials of hydrocarbon degrading bacteria on the phylloplane of some tropical plants. *African Journal of biotechnology*. 5(7): 542.
 130. Jackson C.R., and Denney W.C. (2011). Annual and seasonal variation in the phyllosphere bacterial community associated with leaves of the southern magnolia (*Magnolia grandiflora*). *Microbial ecology*. 61 (1): 113-122.
 131. Jackson C.R, Randolph K.C, Osborn S.L, Tyler H.T (2013). Culture dependent and independent analysis of bacterial communities associated with commercial salad leaf vegetables, *BMC Microbiology*. 13:274.

REFERENCES

132. Jacques M. A. (1996). The effect of leaf age and position on the dynamics of microbial populations on aerial plant surfaces. In *Aerial Plant Surface Microbiology* (pp. 233-248). Springer US.
133. Jasiński M., Kachlicki P., Rodziewicz P., Figlerowicz M. and Stobiecki, M. (2009). Changes in the profile of flavonoid accumulation in *Medicago truncatula* leaves during infection with fungal pathogen *Phoma medicaginis*. *Plant Physiology and Biochemistry*. 47: 847–853.
134. Jayaswal R.K., Fernandez M.A., and Schroeder R.G. (1990). Isolation and characterization of a *Pseudomonas* strain that restricts growth of various phytopathogenic fungi. *Applied and Environmental Microbiology*. 56 (4): 1053-1058.
135. Kagale S., Marimuthu T., Kagale J., Thayumanavan B., Samiyappan R. (2011). Induction of systemic resistance in rice by leaf extracts of *Zizyphus jujuba* and *Ipomoea carnea* against *Rhizoctonia solani*. *Plant Signaling & Behavior*. 6(7):919-923.
136. Kalappanavar I.K., Hiremath R.V. (2000). Biochemical factors for multiple resistance to foliar diseases of sorghum. *Madras Agricultural Journal*. 87:66-70.
137. Kale M.C., Choudhary A.D. (2001). Induction of phenylalanine ammonia-lyase in groundnut cultivars in response to biotic and abiotic stress. *Indian Phytopathology*. 54(3):288-292.
138. Kalita P., Bora L.C., Bhagabati K.N.(1996). Phylloplane microflora of citrus and their role in management of citrus canker. *Indian Phytopathology*. 49(3):234-237.
139. Kang J. H., Liu G., Shi F., Jones A. D., Beaudry R. M. and Howe, G. A. (2010). The tomato odorless-2 mutant is defective in trichome-based production of diverse specialized metabolites and broad-spectrum resistance to insect herbivores. *Plant physiology*. 154(1):262-272.
140. Karabourniotis G., Kotsabassidis D., Manetas Y. (1995). Trichome density and its protective potential against ultraviolet-B radiation damage during leaf development. *Canadian Journal of Botany*. 73:376-383.
141. Karthikeyan M., Radhika K., Mathiyazhagan S., Bhaskaran R., Samiyappan R., Velazhahan R.(2006). Induction of phenolics and defense-related enzymes in coconut (*Cocos nucifera* L.) roots treated with biocontrol agents, *Brazilian Journal of Plant Physiology*. 18(3):367-377.
142. Kerr J.R. (1999). Bacterial inhibition of fungal growth and pathogenicity. *Microbial ecology in health and disease*. 11 (3):129-142.

-
143. Kim D. S. and Hwang B. K. (2014). An important role of the pepper phenylalanine ammonia-lyase gene (PAL1) in salicylic acid-dependent signalling of the defence response to microbial pathogens. *Journal of experimental botany*, eru109.
 144. Kinkel L. L. (1997) Microbial population dynamics on leaves. *Annual Review of Phytopathology*. 35:327–347.
 145. Kishore G.K., Pande S., Podile A.R. (2005). Phylloplane bacteria increase seedling emergence, growth and yield of field-grown groundnut (*Arachis hypogaea L.*), *Letters in Applied Microbiology*. 40:260-268.
 146. Kishore G.K., Pande S., Podile A.R. (2005). Phylloplane bacteria increase seedling emergence, growth and yield of field-grown groundnut (*Arachis hypogaea L.*). *Letters in Applied Microbiology*. 40:260-268.
 147. Klett P.F. (2011). Bacterial-fungal interactions: Hyphens between agricultural, clinical, environmental, and food microbiologists. *Microbiology and Molecular Biology Reviews*. 75(4):583-609.
 148. Knoll D., Schreiber L. (2000). Plant–Microbe Interactions: Wetting of Ivy (*Hedera helix L.*) Leaf Surfaces in Relation to Colonization by Epiphytic Microorganisms, *Microbial Ecology* 41:33–42.
 149. Koenig R. L., Morris R. O. and Polacco J. C. (2002). tRNA is the source of low-level trans-zeatin production in *Methylobacterium* spp. *Journal of bacteriology*. 184(7):1832-1842.
 150. Kojima M., Takeuchi W. (1989). Detection and characterization of p-coumaric Acid hydroxylase in Mungbean, *Vigna mungo*, seedlings, *Journal of Biochemistry*. 105:265-270.
 151. Kolb D. and Müller M. (2003). Different trichome types on the leaves of Styrian oil pumpkin. *Phyton*. 43(2): 365-379.
 152. Kong G.A., Kochman J.K., and Brown J.F. (1997). Phylloplane bacteria antagonistic to the sunflower pathogen *Alternaria helianthi*. *Australasian Plant Pathology*. 26(2):85-97.
 153. Korsten L and Jager ESD, (1995). Mode of action of *Bacillus subtilis* for control of avocado postharvest pathogens. *South African Avocado Growers' Association Yearbook*. Vol. 18, pp. 124-130.
 154. Korsten L., Sanders G.M., Kotze J. (1992). Morphology of the Avocado Phylloplane as Related to Microbial Colonization. In *Proc. of Second World Avocado Congress*. p.145.
 155. Kosuge T. (1969). The role of phenolics in host response to infection. *Annual Review of Phytopathology*. 7(1): 195-222.

REFERENCES

156. Kotireddy M.(1971). Certain chemical constituents of rice plants in relation to resistance to blast disease. *Annals of Annamalai University of Agricultural Research*. 3:106.
157. Kowalski S. P., Eannetta N. T., Hirzel A. T. and Steffens J. C. (1992). Purification and characterization of polyphenol oxidase from glandular trichomes of *Solanum berthaultii*. *Plant Physiology*. 100(2): 677-684.
158. Krimm U., Abanda-Nkpwatt D., Schwab W. and Schreiber, L. (2005). Epiphytic microorganisms on strawberry plants (*Fragaria ananassa* cv. Elsanta): identification of bacterial isolates and analysis of their interaction with leaf surfaces. *FEMS Microbiology Ecology*: 53(3), 483-492.
159. Kuberan T., Vidhyapallavi R.S., Balamurugan A., Nepolean P., Jayanthi R., PremkumarR. (2012). Isolation and biocontrol potential of phylloplane *Trichoderma* against *Glomerella cingulata* in tea. *Journal of Agriculture Technology*. 8:1039-1050.
160. Kuc J. *et al.* (1956). Chlorogenic and caffeic acids as fungistatic agents produced by potatoes in response to inoculation with *Helminthosporium carbonum*. *Journal of the American Chemical Society*. 1.: 2123 - 2125.
161. Kurti P.B. *et al*, (2010). Maize Leaf Epiphytic Bacteria Diversity Patterns Are Genetically Correlated with Resistance to Fungal Pathogen Infection, *Molecular Plant-Microbe Interactions*. 23(4):473-484.
162. Kyndt J. A., Meyer T. E., Cusanovich M. A. and Van Beeumen J. J. (2002). Characterization of a bacterial tyrosine ammonia lyase, a biosynthetic enzyme for the photoactive yellow protein. *FEBS letters*. 512(1-3): 240-244.
163. Lagrimini L.M., Burkhart W., Moyer M. and Rothstein S. (1987). Molecular cloning of complementary DNA encoding the lignin-forming peroxidase from tobacco: Molecular analysis and tissue-specific expression. *Proceedings of the National Academy of Sciences of the USA*.84: 7542-7546.
164. Lanna-Filho R., Romeiro R. D. S. and Alves, E. (2010). Bacterial spot and early blight biocontrol by epiphytic bacteria in tomato plants. *Pesquisa Agropecuária Brasileira*. 45(12): 1381-1387.
165. Last F.T. (1955). Seasonal incidence of *Sporobolomyces* on cereal leaves. *Transactions of the British Mycological Society*. 38:221–239.
166. Last F.T. (1970). Factors associated with the distribution of some phylloplane microbes. *European Journal of Plant Pathology*. 76(3), 140-143.
167. Lattanzio V., Lattanzio V.M.T. and Cardinali A.(2006). Role of phenolics in the resistance mechanisms of plants against fungal pathogens and insects. In Filippo

- Imperato, Phytochemistry: Advances in Research. Kerala: Research Signpost Publishers.23-67.
168. Lauge R., Joosten M., Van den Ackerveken G., Van den Broek H., de Wit P. (1997). The in planta-produced extracellular proteins ECP1 and ECP2 of *Cladosporium fulvum* are virulence factors. *Molecular Plant–Microbe Interactions*. 10: 725–734.
169. Lavania M., Chauhan P. S., Chauhan S. V. S., Singh H. B. and Nautiyal C. S. (2006). Induction of plant defense enzymes and phenolics by treatment with plant growth–promoting rhizobacteria *Serratia marcescens* NBRI1213. *Current microbiology*. 52(5): 363-368.
170. Legard D. E., McQuilken M. P., Whipps J. M., Fenlon J. S., Fermor T. R., Thompson I. P., ... and Lynch J. M. (1994). Studies of seasonal changes in the microbial populations on the phyllosphere of spring wheat as a prelude to the release of a genetically modified microorganism. *Agriculture, Ecosystems & Environment*. 50(2): 87-101.
171. Libbert E., Wichner S., Schiewer U., Risch H., Kaiser W.(1966). The Influence of Epiphytic Bacteria on Auxin Metabolism. *Planta*. 68(4):327-334.
172. Lim H.S., Kim Y.S., Kim S.D.(1991). *Pseudomonas stutzeri* YPL-1 genetic transformation and antifungal mechanism against *Fusarium solani*, an agent of plant root rot. *Applied and Environmental Microbiology*. 57: 510-516.
173. Limtong S., Kaewwichian R., Yongmanitchai W., Kawasaki H.(2014). Diversity of culturable yeasts in phylloplane of sugarcane in Thailand and their capability to produce Indole-3-acetic acid, *World Journal of Microbiology and Biotechnology*.
174. Lin S.Y.H., Trumble J.T. and Kumamoto J. (1987). Activity of volatile compounds in glandular trichomes of *Lycopersicon* species against two insect herbivores. *Journal of Chemical Ecology*. 13:837–850.
175. Lindow S.E, Brandl M.T. (2003). Microbiology of the Phyllosphere. *Applied and Environmental Microbiology*. 69(4):1875-1883.
176. Lindow S.E., Desurmont C., Elkins R., McGourty G., Clark E., Brandl M.T.(1998). Occurrence of Indole-3-Acetic acid- Producing Bacteria on Pear Trees and Their Association with Fruit Russet. *Phytopathology*. 88(11):1149-1157.
177. Lozovaya V. V., Lygin A. V., Zernova O. V., Li S. X., Hartman G. L. and Widholm J. M. (2004). Isoflavonoid accumulation in soybean hairy roots upon treatment with *Fusarium solani*, *Plant Physiology and Biochemistry*. 42: 671–679.
178. Luckwill L. C. (1943). The genus *Lycopersicon*. *Aberdeen Univ. Studies*, (120).

REFERENCES

179. Macarasin D., Patel J., Bauchan G., Giron J. A. and Ravishankar S. (2013). Effect of spinach cultivar and bacterial adherence factors on survival of *Escherichia coli* O157: H7 on spinach leaves. *Journal of Food Protection*®. 76(11): 1829-1837.
180. Madi L., and Katan J. (1998). *Penicillium janczewskii* and its metabolites, applied to leaves, elicit systemic acquired resistance to stem rot caused by *Rhizoctonia solani*. *Physiological and Molecular Plant Pathology*. 53(3): 163-175.
181. Maffei M., Chialva F. and Sacco T. (1989). Glandular trichomes and essential oils in developing peppermint leaves. *New Phytologist*. 111(4): 707-716.
182. Maffei F. H. D. A., Magaldi C., Pinho S. Z., Lastoria S., Pinho W., Yoshida W. B. and Rollo H. A. (1986). Varicose veins and chronic venous insufficiency in Brazil: prevalence among 1755 inhabitants of a country town. *International journal of epidemiology*. 15(2): 210-217.
183. Maffei M., Scannerini S. (2000). UV-B effect on photomorphogenesis and essential oil composition in peppermint (*Mentha piperita* L.). *Journal of Essential Oil Research*. 12:523–529.
184. Magnusson J., Ström K., Roos S., Sjögren J., and Schnürer J. (2003). Broad and complex antifungal activity among environmental isolates of lactic acid bacteria. *FEMS Microbiology Letters*. 219 (1): 129-135.
185. Mandal S.(2010). Induction of phenolics, lignin and key defense enzymes in eggplant (*Solanum melongena* L.) roots in response to elicitors. *African Journal of Biotechnology*. 9(47):8038-8047
186. Mandavia M.K., Parameshwaran M. (1993). Changes in amino acids and phenols in lima bean (*Phaseolus lunatus*) varieties resistant and susceptible to stem rot disease. *Gujarat Agricultural University Research Journal*. 18(2):19-23.
187. Mandrell, R., Gorski, L., & Brandl, M. (2006). Attachment of microorganisms to fresh produce. *Microbiology of fruits and vegetables*. CRC, Taylor and Francis, Boca Raton, FL, 34-61.
188. Marks M.D., Tian L., Wenger J.P., Omburo S.N., Soto-Fuentes W., He, J., Gang D.R., Weiblen G.D. and Dixon R.A. (2009) Identification of candidate genes affecting delta(9)-tetrahydrocannabinol biosynthesis in *Cannabis sativa*. *Journal of Experimental Botany*. 60 :3715–3726.
189. May R. B., Völksch, and Kampmann G. (1997). Antagonistic activities of epiphytic bacteria from Soybean leaves against *Pseudomonas syringae* pv. *glycinea* *in vitro* and *in planta*. *Microbial Ecology*. 34 (2): 118-124.
190. Mayer A.M., Harel E and Shaul R. B.(1965). Assay of catechol oxidase a critical comparison of methods. *Phytochemistry*, 5: 783-789.

-
191. Mayer A.M., Harel E. (1979). Polyphenol oxidases in plants. *Phytochemistry*. 18:193-215.
 192. Meyer K.M., and Johan H.J Leveau. (2012). Microbiology of the phyllosphere: a playground for testing ecological concepts. *Oecologia* 168.3: 621-629.
 193. McCormack P. J., Wildman H. G. and Jeffries P. (1994). Production of antibacterial compounds by phylloplane-inhabiting yeasts and yeastlike fungi. *Applied and environmental microbiology*, 60(3): 927-931.
 194. Meena B., Radha Jayalaskhmi R., Vidhyasekharan P. and Velazhakan P.(2000). Effect of foliar application *Pseudomonas fluorescence* on activities of phenylalanine ammonia-lyase, chitinase and 1, 3-gluconase and accumulation of phenolics in rice, *Acta-Phytopathology Entomol. Hung.* 34:307-315.
 195. Meena K. K., Kumar M. Kalyuzhnaya M. G., Yandigeri M. S., Singh D. P., Saxena A. K. and Arora D. K. (2012). Epiphytic pink-pigmented methylotrophic bacteria enhance germination and seedling growth of wheat (*Triticum aestivum*) by producing phytohormone. *Antonie Van Leeuwenhoek*.101(4): 777-786.
 196. Mercier J. and Lindow S.E. (2000). Role of leaf surface sugars in colonization of plants by bacterial epiphytes. *Applied Environmental Microbiology*. 66 (1): 369-374.
 197. Michereff S. J., Silveira N. S. S., Reis A., and Mariano R. L. R. (1994). Epiphytic bacteria antagonistic to *Curvularia* leaf spot of yam. *Microbial ecology*. 28(1): 101-110.
 198. Milillo, S.R.; Badamo, J.M.; Boor, K.J.; Wiedmann, M. (2008) Growth and persistence of *Listeria monocytogenes* isolates on the plant model *Arabidopsis thaliana*. *Food Microbiology*. 25, 698-704.
 199. Mishra S., Singh A., Keswani C., Saxena A., Sarma B. K and Singh, H. B. (2015). Harnessing plant-microbe interactions for enhanced protection against phytopathogens. In *Plant Microbes Symbiosis: Applied Facets* (pp. 111-125). Springer India.
 200. Mitra R., Cuesta-Alonso E., Wayadande A., Talley J., Gilliland S., Fletcher J. (2009). Effect of route of introduction and host cultivar on the colonization, internalization, and movement of the human pathogen *Escherichia coli* O157:H7 in spinach. *Journal of Food Protection*. 72, 1521–1530.
 201. Mohamed A. and Sater A. (2001). Antagonistic interactions between fungal pathogen and leaf surface fungi of Onion (*Allium cepa*), *Pak J Biol Sci*. 4 (7): 838-842.
 202. Monaghan J., Zipfel C.(2012). Plant pattern recognition receptor complexes at the plasma membrane. *Current Opinion in Plant Biology*. 15: 349–357.

REFERENCES

203. Monier J. M. and Lindow S. E. (2004). Frequency, size, and localization of bacterial aggregates on bean leaf surfaces. *Applied and environmental microbiology*. 70(1): 346-355.
204. Morris C. E. and Rouse D. I. (1985) Role of nutrients in regulating epiphytic bacterial populations. In: *Biological control of the Phylloplane* (Ed. by C. E. Windels & S. E. Lindow), The American Phytopathological Society, Minnesota. pp. 63–82.
205. Morrison T.A., Buxton D.R. (1993). Activity of phenylalanine ammonia-lyase, tyrosine ammonia-lyase, and cinnamyl alcohol dehydrogenase in the maize stalk. *Crop Science*. 33:1264-1268.
206. Naranasamy P., (2013). *Biological Management of Diseases of Crops*. Springer. 673.
207. Newell S.Y. *et al.*, (1989). Decomposition and microbial dynamics for standing, naturally positioned leaves of the salt-marsh grass *Spartina alterniflora*, *Marine Biology*. 101(4):471-481.
208. Nicholson R.L., Hammerschmidt R. (1992). Phenolic compounds and their role in disease resistance. *Annual Review of Phytopathology*; 30:369-389.
209. Nicot P.C. (2011). *Classical and Augmentative Biological Control Against Diseases and Pests: Critical Status Analysis and Review of Factors Influencing Their Success*. International Organization for Biological and Integrated Control of Noxious Animals and Plants, West Palaearctic Regional Section (IOBC/WPRS), Europe
210. Ojha S., Chatterjee N.C. (2012). Induction of resistance in tomato plants against *Fusarium oxysporum* f.sp. *lycopersici* mediated through salicylic acid and *Trichoderma harzianum*. *Journal of Plant Protection Research*. 52(2):220-225.
211. O'Leary B. M., Rico A., McCraw S., Fones H. N. and Preston G. M. (2014). The infiltration-centrifugation technique for extraction of apoplastic fluid from plant leaves using *Phaseolus vulgaris* as an example. *Journal of visualized experiments: JoVE*. (94).
212. Oriani M. A. D. G., Vendramim J. D. and Vasconcelos C. J. (2011). No-choice ovipositional nonpreference of *Bemisia tabaci* (Gennadius) B biotype on tomato genotypes. *Scientia Agricola*, 68(2): 147-153.
213. Owens Jr W. A. (1953). Age and mental abilities: a longitudinal study. *Genetic Psychology Monographs*.
214. Pandey A., Palni L. M. S. and Coulomb N. (1997). Antifungal activity of bacteria isolated from the rhizosphere of established tea bushes, *Microbiological Research*. 152(1):105-112.

-
215. Pandey R. R., Arora D. K. and Dubey R. C. (1993). Antagonistic interactions between fungal pathogens and phylloplane fungi of guava. *Mycopathologia*. 124(1): 31-39.
216. Parent J. G. and Asselin, A. (1984). Detection of pathogenesis-related proteins (PR or b) and of other proteins in the intercellular fluid of hypersensitive plants infected with tobacco mosaic virus. *Canadian Journal of Botany*, 62(3), 564-569.
217. Parr A.J. and Bolwell G.P. (2000). Phenols in the plant and in man. The potential for possible nutritional enhancement of the diet by modifying the phenols content or profile. *Journal of the Science of Food and Agriculture*. 80: 985–1012.
218. Pati B.R., Sengupta S., Chandra A.K.(1995). Impact of selected phyllospheric diazotrophs on the growth of wheat seedlings and assay of the growth substances produced by diazotrophs, *Microbiological Research*. 150(2):121-127.
219. Patil P.V. and Kachapur M.R.(1998). Role of phyllosphere micro-organisms in the control of Sunflower rust caused by *Puccinia helianthi* Schw. *Karnataka Journal of Agricultural Sciences*. 11 (2): 410-414.
220. Petatán-Sagahón I., Anducho-Reyes M.A., Silva-Rojas H.V., Arana-Cuenca A., Tellez-Jurado A., Cárdenas-Álvarez I.O., and Mercado-Flores Y. (2011). Isolation of bacteria with antifungal activity against the phytopathogenic fungi *Stenocarpella maydis* and *Stenocarpella macrospora*. *International Journal of Molecular Sciences*.12(9): 5522–5537.
221. Phelan V. V., Liu W. T., Pogliano K. and Dorrestein P. C. (2012). Microbial metabolic exchange [mdash] the chemotype-to-phenotype link. *Nature chemical biology*. 8(1): 26-35.
222. Pimentel I. C., Glienke-Blanco C., Gabardo J., Stuart R. M. and Azevedo J. L. (2006). Identification and colonization of endophytic fungi from soybean (*Glycine max* (L.) Merrill) under different environmental conditions. *Brazilian archives of biology and technology*, 49(5):705-711.
223. Podile R., Lami V.D.V.(1998). Seed bacterization with *Bacillus subtilis*, AF1 increase PAL and reduce the incidence of *Fusarium* wilt in Pigeon pea. *Journal of Phytopathology*. 146:255-259.
224. Prabakaran.M., Merinal, S. and Panneerselvam, A. (2011). Investigation of phylloplane mycoflora from some medicinal plants. *European Journal of Experimental Biology*, 1 (2):219-225.
225. Pradeep T., Jambhale N.D.(2002) .Relationship between phenolics, polyphenol oxidase and peroxidase, and resistance to powdery mildew in *Zizyphus*. *Indian Phytopathology*. 55(2):195-196.

REFERENCES

226. Prasanna N.D., Vijayalakshmi K., Seshagirirao K. and Shaheen S.K. (2014). Characterization of antifungal compounds produced by *Pseudomonas stutzeri* EGB3 isolated from the gut of earthworm (*Eisenia foetida*). *Journal of Microbiology and Antimicrobials*. 6 (3): 57-65.
227. Prasannath.K. (2013). Pathogenecity and Virulence Factors of Phytobacteria. *Scholars Academic Journal of Biosciences*. 1(1):24-33.
228. Pruvost, O., Boher, B., Brocherieux, C., Nicole, M., and Chiroleu, F. (2002). Survival of *Xanthomonas axonopodis* pv. *citri* in leaf lesions under tropical environmental conditions and simulated splash dispersal of inoculum. *Phytopathology*, 92(4): 336-346.
229. Rajendiran R., Jegadeeshkumar D., Sureshkumar B. T. and Nisha T. (2010). In vitro assessment of antagonistic activity of *Trichoderma viride* against post harvest pathogens. *Journal of Agricultural Technology*. 6(1): 31-35.
230. Raju S., Jayalakshmi S. K. and Sreeramulu K. (2008). Comparative study on the induction of defense related enzymes in two different cultivars of chickpea (*Cicer arietinum* L) genotypes by salicylic acid, spermine and *Fusarium oxysporum* f. sp. *ciceri*. *Australian Journal of Crop Science*. 2(3): 121-140.
231. Ramos-Solano B., Algar E., Gutierrez-Mañero F. J., Bonilla A., Lucas J. A. and García-Seco D. (2015). Bacterial bioeffectors delay postharvest fungal growth and modify total phenolics, flavonoids and anthocyanins in blackberries. *LWT-Food Science and Technology*. 61(2): 437-443.
232. Ran L., Gu W. Z. and Wu G. J. (2004). Role of Salicylic Acid in Induction of Resistance against Bacterial Wilt in *Eucalyptus urophylla* and Changes of Peroxidase and Polyphenol Oxidase [J]. *Forest Research*, 1, 003.
233. Rao R.P., Hunter A., Kashpur O., Normanly J.(2010). Aberrant Synthesis of Indole-3-Acetic acid in *Saccharomyces cerevisiae* Triggers Morphogenic Transition, A Virulence Trait of Pathogenic Fungi. *Genetics*. 185(1):211-220.
234. Redford A. J. and Fierer N. (2009). Bacterial succession on the leaf surface: a novel system for studying successional dynamics. *Microbial ecology*,58(1):189-198.
235. Rerie W.G., Feldmann K.A., Marks M.D.(1994). The GLABRA2 gene encodes a homeo domain protein required for normal trichome development in *Arabidopsis* *Genes Dev*.813881399.
236. Reuveni M., and Cohen Y. (1978). Growth retardation and changes in phenolic compounds, with special reference to scopoletin, in mildewed and ethylene-treated tobacco plants. *Physiological Plant Pathology*. 12(2), 179IN11183-182189.

237. Rini C.R. and Sulochana K.K.(2007). Usefulness of Trichoderma and Pseudomonas against Rhizoctonia solani and Fusarium oxysporum infecting tomato. Journal of Tropical Agriculture.45 (1-2): 21–28.
238. Romero D., Pérez-García A., Rivera M. E., Cazorla F. M. and De Vicente A. (2004). Isolation and evaluation of antagonistic bacteria towards the cucurbit powdery mildew fungus Podosphaera fusca. Applied microbiology and biotechnology. 64(2): 263-269.
239. Rosler J, Krekel F, Amerhein N, Schmid J.(1997) Maize phenylalanine ammonia-lyase has tyrosine ammonia-lyase activity. Plant Physiology. 113:175-179.
240. Ruelas C., Tiznado-Hernández M. E., Sánchez-Estrada A., Robles-Burgueño M. R., and Troncoso-Rojas R. (2006). Changes in phenolic acid content during Alternaria alternata infection in tomato fruit. Journal of phytopathology. 154(4): 236-244.
241. Ruinen J. (1956). Occurrence of Beijerinckia species in the phyllosphere. Nature 177:220-221.
242. Ryals J.A., Neuenschwander U.H., Willits M.G., Molina A., Steiner H.Y., Hunt M.D.(1996). Systematic acquired resistance. Plant cell. 8: 1809-1819.
243. Saha A.K., Ray A., Das P. (2013). Fungal colonization of phylloplane of Psidium guineense sw. growing in Suryamaninagar, Tripura, Northeast India, International Journal of Basic and Applied Chemical Sciences. 3 (1):62-67.
244. Saravanan T., Bhaskaran R., Muthusamy M.(2004). Pseudomonas fluorescens induced enzymological changes in banana roots (Cv. Rasthali) against Fusarium wilt disease. Plant pathology Journal .3:72-80.
245. Sarkar K.R., Joshi R.D.(1977). Peroxidase and catalase activity in MLO infected brinjal. Indian Phytopathology. 30:518-521.
246. Sathiyathan S., Vidhyasekaran P. (1981). Role of phenolics in brown spot disease resistance in rice. Indian Phytopathology. 34:225-231.
247. Savary S. (1987). Decrease by plant development and leaf age of susceptibility of groundnut to rust (Puccinia arachidis) in a susceptible cultivar. Netherlands Journal of Plant Pathology, 93(1): 25-31.
248. Sawale A., Kadam T.A., Karale M.A., and Kadam O.A., (2014). Antimicrobial Activity of Secondary Metabolites from Halophilic Bacillus pumilus sp. International Journal of Current Microbiology and Applied Sciences. 3(6): 506-512.
249. Scagel C.F., Lee J.,(2012). Phenolic composition of basil plants is differentially altered by plant nutrient status and inoculation with mycorrhizal fungi. HortScience, 47(5): 660-671.

REFERENCES

250. Schillmiller A. L., Miner D. P., Larson M., McDowell E., Gang D. R., Wilkerson C. and Last, R. L. (2010). Studies of a Biochemical Factory: Tomato Trichome Deep Expressed Sequence Tag Sequencing and Proteomics. *Plant Physiology*. 153(3), 1212–1223.
251. Schliemann W., Ammer C., and Strack D. (2008). Metabolite profiling of mycorrhizal roots of *Medicago truncatula*. *Phytochemistry*. 69: 112–146
252. Schnürer J. and Magnusson J. (2005). Antifungal lactic acid bacteria as biopreservatives. *Trends in Food Science & Technology*. 16 (1–3): 70–78.
253. Scholze H. and Boch J. (2011). TAL effectors are remote controls for gene activation. *Current opinion in microbiology*. 14(1):47-53.
254. Schrey S.D., Eric Erkenbrack E., Früh E., Fengler S., Hommel K., Horlacher N., Schulz D., Ecke M., Kulik A., Fiedler H.P., Hampp R. and Tarkka M.T.(2012). Production of fungal and bacterial growth modulating secondary metabolites is widespread among mycorrhiza-associated streptomycetes, *BMC Microbiology*. 12 (1): 164.
255. Sen K., Sengupta C. and Saha J. Received: 4 th Aug-2014 Revised: 20 th Aug-2014 Accepted: 21 st Aug-2014 Research article PGPR Consortium In Alleviating Downy Mildew Of Cucumber.
256. Sharma I.M.(2003). Relationship between phenylpropanoid metabolism and resistance to *Pythium ultimum* Throw. in apple root stocks. *Journal of Mycology and Plant Pathology*. 33(1):114-118.
257. Sharma J.R., Mishra B., Krishna Jha.(1992). Biochemical relationship in resistant and susceptible cultivars with Turicum leaf blight disease in maize. *Indian Phytopathology*. 45:241-243.
258. Sharma P.D. (1984). Ecology of phylloplane and litter fungi of triticale, *Nordic Journal of Botany*.: 707-71.
259. Sheldrake A. R. (1973). The production of hormones in higher plants. *Biological Reviews*. 48(4):509-559.
260. Shenton M. R., Berberich T., Kamo M., Yamashita T., Taira H. and Terauchi R. (2012). Use of intercellular washing fluid to investigate the secreted proteome of the rice–Magnaporthe interaction. *Journal of plant research*. 125(2): 311-316.
261. Sherf B.A., Bajar A.M., Kolattukudy P.E.(1993). Abolition of an Inducible Highly Anionic Peroxidase Activity in Transgenic Tomato. *Plant Physiology*. 101(1):201-208.
262. Shulaev V., Silverman P., Raskin I. (1997). Airborne signalling by methyl salicylate in plant pathogen resistance. *Nature*, 385: 718-721.

-
263. Sivakumar G., Sharma R.C. (2003). Induced biochemical changes due to seed bacterization by *Pseudomonas fluorescens* in maize plants. *Indian Phytopathology* .56:134-137.
264. Skadhauge B. Thomsen K. von Wettstein D.(1997). The role of barley testa layer and its flavonoid content in resistance to *Fusarium* infections. *Hereditas* . 126: 147–160.
265. Smid E. J. and Lacroix C. (2013). Microbe–microbe interactions in mixed culture food fermentations. *Current opinion in biotechnology*. 24(2):148-154.
266. Smith K., Goodman.R, (1999). Host variation for Interactions with Beneficial Plant-Associated Microbes. *Annual Review of Phytopathology*. 37: 473-491.
267. Solanki M.K., Robert A.S., Singh R.K., Kumar S., Pandey A.K., Srivastava A.K., Arora D.K. .(2012).Characterization of mycolytic enzymes of *Bacillus* strains and their bio-protection role against *Rhizoctonia solani* in tomato. *Current Microbiology* . 65(3):330-336.
268. Sowndhararajan K., Marimuthu S. and Manian S. (2013). Biocontrol potential of phylloplane bacterium *Ochrobactrum anthropi* BMO-111 against blister blight disease of tea. *Journal of applied microbiology*. 114(1): 209-218.
269. Sowndhararajan K., Marimuthu S. and Manian S. (2013). Biocontrol potential of phylloplane bacterium *Ochrobactrum anthropi* BMO-111 against blister blight disease of tea. *Journal of applied microbiology*. 114(1): 209-218.
270. Srivastava R. and Shalini. (2008). Antifungal activity of *Pseudomonas fluorescens* against different plant pathogenic fungi, *The Internet Journal of Microbiology*. 7(2): 2789-2796.
271. Stockwell V. O., McLaughlin R. J., Henkels M. D., Loper J. E., Sugar D. and Roberts R. G.(1999). Epiphytic colonization of pear stigmas and hypanthia by bacteria during primary bloom. *Phytopathology*. 89:1162-1168.
272. Suárez-Estrella F., Vargas-Garcia C., Lopez M. J., Capel C. and Moreno J. (2007). Antagonistic activity of bacteria and fungi from horticultural compost against *Fusarium oxysporum* f. sp. *melonis*. *Crop Protection*. 26(1) :46-53.
273. Swadling I.R. and Jeffries P., (2010). Antagonistic properties of two bacterial biocontrol agents of Grey Mould disease. *Biocontrol Science and Technology*. 8 (3): 439-448.
274. Taggar G.K.; Gill R.S.; Gupta A.K.; Sandhu J.S.(2012) *Plant Signaling & Behavior*. 7(10), 1321-1329
275. Tanti A., Bhattacharyya P., Dutta P., Sarmah S. R., Madhab M., Saikia D., Kachari A, and Berceoyjy R. (2016). Diversity of phylloplane microflora in certain tea

REFERENCES

- cultivars of Assam, North-East India. *European Journal of Biological Research*, 6(4), 287-292.
276. Taware P. B., Dhumal K. N., Oulkar D. P., Patil S. H. and Banerjee K. A. U. S. H. I. K. (2010). Phenolic alterations in grape leaves, berries and wines due to foliar and cluster powdery mildew infections. *International Journal of Pharma and Bio Sciences*. 1: 1-14.
277. Telfer A., Bollman K. M., Poethig R. S. (1997). Phase change and the regulation of trichome distribution in *Arabidopsis thaliana*. *Development* .124: 645-654.
278. Thakur S. and Harsh N. S. K. (2014). Efficacy of volatile metabolites of phylloplane fungi of *Rauwolfia serpentina* against *Alternaria alternata*. *Forest Pathology*, 1, 2.
279. Thangavelu R., Palaniswami A., Doraiswamy S. and Velazhahan R. (2003). The effect of *Pseudomonas fluorescens* and *Fusarium oxysporum* f. sp. *cubense* on induction of defense enzymes and phenolics in banana. *Biologia Plantarum*. 46(1): 107-112.
280. Thipyapong P., Hunt M.D., Steffens J.C. (2004). Antisense downregulation of polyphenol oxidase results in enhanced disease susceptibility. *Planta* . 220:105-117.
281. Thipyapong P., Steffens J.C. (1997). Tomato Polyphenol Oxidase: Differential Response of the Polyphenol Oxidase F Promoter to Injuries and Wound Signals. *Plant Physiology*. 115:409-418.
282. Thomashow, L. S., & Weller, D. M. (1996). Current concepts in the use of introduced bacteria for biological disease control: mechanisms and antifungal metabolites. In *Plant-microbe interactions* (pp. 187-235). Springer US.
283. Thompson I. P., Bailey M. J., Fenlon J.S., Fermor T. R., Lilley A. K., Lynch J. M., McCormack P. J., McQuilken M. P. and Purdy K. J. (1993). Quantitative and qualitative seasonal changes in the microbial community from the phyllosphere of sugar beet (*Beta vulgaris*). *Plant Soil*.150:177–191.
284. Tokin B.P (1960). Phytoncides, The destroyer of microbes. *Moscow Izdet. Sov. Rossiya*, p. 198.
285. Treutter D. (2005). Significance of flavonoids in plant resistance and enhancement of their biosynthesis. *Plant Biology*. 7: 581–591.
286. Treutter D. (2006). Significance of flavonoids in plant resistance: a review. *Environmental Chemistry Letters*. 4(3): 147-157.
287. Tudzynski B. (1997). Fungal Phytohormones in Pathogenic and Mutualistic Associations. *The Mycota*. 5:167-184.

-
288. Tyagi V. K., and Chauhan S. K. (1982). The effect of leaf exudates on the spore germination of phylloplane mycoflora of chilli (*Capsicum annum* L.) cultivars. *Plant and Soil*. 65(2): 249-256.
289. Uknes S., Mauch-Mani B., Moyer M., Potter S., Williams S., Dincher S., Chandler D., Slusarenko A., Ward E., Ryals J. (1992). Acquired resistance in *Arabidopsis*. *Plant Cell* 4: 645–656
290. Upadhyay R. K. (1981). Effect of volatile phylloplane fungal metabolites on the growth of a foliar pathogen. *Experientia*. 37(7): 707-708.
291. Vacheron J., Desbrosses G., Bouffaud M. L., Touraine B. and Prigent-Combaret C. (2014). Plant growth-promoting rhizobacteria and root system functioning. *Ecophysiology of root systems-environment interaction*, 166.
292. Valkama E., Salminen J. P., Koricheva J. and Pihlaja K. (2003). Comparative analysis of leaf trichome structure and composition of epicuticular flavonoids in Finnish birch species. *Annals of Botany*. 91(6): 643-655.
293. Van Huystee R.B.(1987). Some molecular aspects of plant peroxidase biosynthetic studies. *Annual Review of Plant Physiology*. 38: 205-219.
294. van Loon L.C., Rep M., Pieterse C.M.J. (2006). Significance of inducible defense-related proteins in infected plants. *Annual Review of Phytopathology*. 44: 135–162.
295. Vandenberg P.A., Gonzalez C.F., Wright A.M., and Kunka B.S., (1983). Iron-Chelating Compounds Produced by Soil Pseudomonads: Correlation with Fungal Growth Inhibition, *Applied and Environmental Microbiology*. 46(1): 128-132.
296. Vastakaite V., Buzaitė O.(2011). Plant growth promoting properties of culturable phylloepiphytic pine bacteria, *Biologija*. 57(1):33-36.
297. Velasco G.L., Welbaum G.E., Falkinham J.O., Ponder M.A.(2011). Phylloplane Bacterial Community Structure of Spinach (*Spinacia oleracea*) as Affected by Cultivar and Environmental Conditions at Time of Harvest, *Diversity* 3:721-738.
298. Venkateshvarulu B., Sirohi C.E.(1976). Photoperiods studies on disease intensity, sugars, starch and phenol content of brown rust of wheat. *Indian Journal of Agricultural Sciences*. 44:549-554.
299. Verhagen B.W., Trotel-Aziz P., Couderchet M., Hofte M., Aziz A.(2010) . *Pseudomonas* spp.-induced systemic resistance to *Botrytis cinerea* is associated with induction and priming of defence responses in grapevine. *Journal of Experimental Botany*. 61:249-260.
300. Vidhyasekaran P. (1974). Possible role of action of phenolics in inducing disease resistance in ragi plants against *Helminthosporium tetramera*. *Indian Phytopathology*. 22:587-595.

REFERENCES

301. Volksch B., Nußske J. and May R. (1996). Characterization of two epiphytic bacteria from soybean leaves with antagonistic activities against *Pseudomonas syringae* pv. *glycinea*. *Journall of Basic Microbiology*. 36:453–462.
302. Vorholt J.A.,(2012). Microbial life in the phyllosphere, *Nature Reviews Microbiology*.10:828-840.
303. Walker H. L. (1981). Granular formulation of *Alternaria macrospora* for control of spurred anoda (*Anoda cristata*). *Weed Science*. 342-345.
304. Wang Y., Kim S.G., Wu J., Huh H.H., Lee S.J., Rakwal R., *et al.* (2013). Secretome analysis of the rice bacterium *Xanthomonas oryzae* (Xoo) using in vitro and in planta systems. *Proteomics*.13:1901–1912. doi: 10.1002/pmic.201200454
305. Warmink J.A. and Elsas J.D. (2009). Migratory Response of Soil Bacteria to *Lyophyllum* sp. Strain Karsten in Soil Microcosms. *Applied and Environmental Microbiology*.75(9): 2820-2830.
306. Watts K. T., Mijts B. N., Lee P. C., Manning A. J. and Schmidt-Dannert C. (2006). Discovery of a substrate selectivity switch in tyrosine ammonia-lyase, a member of the aromatic amino acid lyase family. *Chemistry & biology*. 13(12): 1317-1326.
307. Weiskopf L.(2013). The potential of bacterial volatiles for crop protection against phytopathogenic fungi.
308. Welinder K.G., Justesen A.F., Kjaersgård I.V., Jensen R.B., Rasmussen S.K., Jespersen H.M. and Duroux, L. (2002). Structural diversity and transcription of class III peroxidases from *Arabidopsis thaliana*. *European Journal of Biochemistry*. 269(24): 6063-6081
309. Werker E., Putievsky E., Ravid U., Dudai N., Katzir I.(1993) Glandular hairs and essential oil in developing leaves of *Ocimum basilicum* L. (Lamiaceae). *Annals of Botany*.71:43–50.
310. Weryszko-Chmielewska E. and Chernetsky M. (2005). Structure of trichomes from the surface of leaves of some species of *Kalanchoë* Adans. *Acta Biologica Cracoviensia Series Botanica*. 47(2): 15-22.
311. Whipps J.M., Hand P., Pink D., Bending G.D. (2008). Phyllosphere microbiology with special reference to diversity and plant genotype, *Journal of Applied Microbiology*. 105(6):1744-1755.
312. Whipps J.M., Hand P., Pink D.A., Bending G.D. (2008). Human pathogens and the phyllosphere, *Advances in applied microbiology*. 64:183-221.
313. Wilkens R. T., Shea G. O., Halbreich S. and Stamp N. E. (1996). Resource availability and the trichome defenses of tomato plants. *Oecologia*. 106(2): 181-191.

-
314. Williams T.R. and Marco M.L.(2014). Phyllosphere microbiota composition and microbial community transplantation on lettuce plants grown indoors. *MBio* 5:e01564-14.doi:10.1128/mBio.01564-14.
315. Wojakowska A., Muth D., Narożna D., Mądrzak C., Stobiecki M. and Kachlicki P. (2013). Changes of phenolic secondary metabolite profiles in the reaction of narrow leaf lupin (*Lupinus angustifolius*) plants to infections with *Colletotrichum lupini* fungus or treatment with its toxin. *Metabolomics*. 9(3): 575-589.
316. Wolin M. J., Miller T. L. and Stewart C. S. (1997). Microbe-microbe interactions. In *The rumen microbial ecosystem* (467-491). Springer Netherlands.
317. Wood K.R., Barbara D.J. (1971). Virus multiplication and peroxidase activity in leaves of cucumber (*Cucumis sativus* L.) cultivars systemically infected with W strain of *Cucumber mosaic virus*. *Physiological Plant Pathology*.1:73-81.
318. Yadav R. K. P., Halley J. M., Karamanoli K., Constantinidou,H. I. and Vokou, D. (2004). Bacterial populations on the leaves of Mediterranean plants: quantitative features and testing of distribution models, *Environmental and experimental botany*, 52(1): 63-77.
319. Yadav R.K.P., Karamanoli K., Vokou D.(2011). Bacterial populations on the phyllosphere of Mediterranean plants: influence of leaf age and leaf surface, *Frontiers of Agriculture in China* 5(1):60-63.
320. Yang C. H. and Crowley D. E. (2000). Rhizosphere microbial community structure in relation to root location and plant iron nutritional status. *Applied and environmental microbiology*. 66(1): 345-351.
321. Yao Y. Y. and Wong S. K. M. (1995). Generalization of rough sets using relationships between attribute values. In *Proceedings of the 2nd annual Joint Conference on Information sciences* :30-33.
322. Zacheo G., Orlando C., Zacheo T.B.(1993). Characterization of anionic peroxidases in tomato isolines infected by *Meloidogyne incognita*. *Journal of Nematology*. 25(2):249-56.
323. Zhang L., Tian L. H., Zhao J. F., Song Y., Zhang C. J. and Guo Y. (2009). Identification of an apoplastic protein involved in the initial phase of salt stress response in rice root by two-dimensional electrophoresis. *Plant Physiology*. 149(2): 916-928.
324. Živkovic S., Stojanović S., Ivanović Ž., Gavrilović V., Popović T., and Balaž J. (2010). Screening of antagonistic activity of microorganisms against *Colletotrichum acutatum* and *Colletotrichum gloeosporioides*. *Archives of Biological Sciences*. 62 (3): 611-623.