

LIST OF FIGURES

Figure no.	Description
2.1	Leading causes of children death worldwide
2.2	Incidences of diarrheal deaths in children below 5 years worldwide
2.3	A: Number of cases of infections due to various microbial pathogens B: Number of deaths due to microbial infection worldwide in 2000
2.4	<i>S. dysenteriae</i> epidemic episodes worldwide
2.5	Evolution of invasive pathogenic <i>Shigella</i> from non-invasive non-pathogenic <i>E. coli</i> to whom it is closely related
2.6	TTSA needle like complex of <i>Shigella</i> to release effector molecules in host cells
2.7	Pathogenesis of <i>Shigella</i>
2.8	Caspase 1 dependent apoptosis of macrophage engulfing by <i>Shigella</i> via release of IpaB and IpaC
2.9	Cytokine profile during <i>Shigella</i> infection in intestinal tract
2.10	Additional probiotic selection criteria for screening commercializing probiotic strain
2.11	Probiotics mediated immune system activation
2.12	Effect of probiotic lactobacilli on liver and intestinal cells that play crucial role in prevention of infection
2.13	Mechanisms behind the probiotic action in intestinal tract
2.14	Antimicrobial substances produced by lactobacilli
2.15	Mode of bacteriocin action targeting Gram positive and Gram negative bacteria
2.16	Probiotic mediated modulation of host immune system
2.17	Safety assessment of probiotic microorganisms
4.1	Growth of acid producing lactobacilli on MRS-BCP agar (A) control: uninoculated MRS-BCP agar plate, (B) Colonies of microorganisms isolated from stool sample after 24 hours of incubation on MRS-BCP agar plates
4.2	Growth of acid producing colony (putative <i>Lactobacillus</i>) on MRS agar plates after 24 hours of incubation
4.3	Microscopic image (100X magnification) of Gram positive rod like bacteria, A) iso RT4-54; B) isolate RT5-6; C) isolate RT9-10; D) isolate RT16-2; E) isolate RT26-6; F) isolate RT27-17

4.4	Photographic images of test tubes of biochemical tests; 1) Simmons citrate agar; 2) TSI agar; 3) HL agar; 4) peptone water with sugar and Andrade's reagent; C-negative control and T-test sample (putative <i>Lactobacillus</i> isolate)
4.5	Agarose (1%) gel showing PCR-DNA
4.6	TGE agar plates showing zone of inhibition against <i>S. dysenteriae</i> lawn
4.7	16 SrRNA gene sequences of RT4-54, RT5-2, RT9-10, RT16-2, RT26-6 and RT27-17
4.8	A phylogenetic tree of isolated probiotic lactobacilli based 16 SrRNA gene sequencing using neighbour-joining method in MEGA 4 using Kimura 2 parameter model.
4.9	Adherence of probiotic lactobacilli to Caco-2 cells observed under oil immersion microscope (100X magnification) after staining with giemsa stain
4.10	Coaggregation of bacterial pathogens by lactobacilli
4.11	Lactobacilli mediated competitive inhibition of bacterial pathogen adhesion on Caco-2 cells
4.12	MRS-agar plates exhibiting biocompatibility amongst the isolates, a- <i>Lactobacillus pentosus</i> , b- <i>L. paraplantarum</i> , c- <i>L. rhamnosus</i>
4.13	Phagocytic killing of <i>S. dysenteriae</i> in rat peritoneal macrophages under lactobacilli treatment groups; A) lactobacilli cells; B) LCFS
4.14	Estimation of lipid peroxidation (MDA): outcome on the lipid-peroxidation levels (in terms of MDA levels) of rat peritoneal macrophages infected with <i>S. dysenteriae</i> type 1 in presence of a) lactobacilli cells b) LCFS of probiotic lactobacilli.
4.15	Estimation of nitrite levels: effect of a) lactobacilli cells b) LCFS of probiotic lactobacilli, on nitrite levels produced in macrophage infected with <i>S. dysenteriae</i> type 1.
4.16	Estimation of levels of SOD: effect of a) probiotic lactobacilli cell suspension, b) LCFS of probiotic lactobacilli on SOD levels of macrophage.
4.17	Levels of reduced glutathione (GSH): effect of LCFS of probiotic lactobacilli on GSH levels in macrophages.
4.18	Adherence and internalization of <i>S. dysenteriae</i> in presence of probiotic lactobacilli treatment groups under three treatment sets; A) Percentage <i>Shigella</i> adherence to Caco-2 cells; B) Percentage <i>Shigella</i> internalization in Caco-2 cells.

4.19	Adherence and internalization of <i>S. dysenteriae</i> in presence of probiotic LCFS treatment groups under three treatment sets; A) Percentage <i>Shigella</i> adherence to Caco-2 cells; B) Percentage <i>Shigella</i> internalization in Caco-2 cells
4.20	Percentage survivability of Caco-2 cells under lactobacilli treatment groups
4.21	Agarose (1.5%) gel profile showing RT-PCR amplified products
4.22	Agarose (1.5%) gel profile showing RT-PCR amplified products
4.23	Gene Expression analysis of genes of toll like receptors (<i>TLR2</i> , <i>TLR4</i> , <i>TLR9</i>)
4.24	Gene Expression analysis of genes of nuclear transcriptional factor and pro-inflammatory genes
4.25	Gene Expression analysis of genes of anti-inflammatory cytokines (<i>IL-10</i> , <i>TGF-β</i>)
4.26	Gene Expression analysis of genes of interleukin (<i>IL-6</i> , <i>IL-12</i> , <i>IL18</i> , <i>IL-21</i>)
4.27	96 well plate image depicting MIC of LCFS against <i>S. dysenteriae</i> growth
4.28	Time dependent killing of <i>S. dysenteriae</i> by LCFS of each lactobacilli taken individually and in cocktail
4.29	Lactic acid production by each lactobacilli isolates A) LP4; B) LP1; C) LP6
4.30	Hydrogen peroxide production by lactobacilli isolates A) LP4; B) LP1; C) LP6