

List of Figures

1.1	Schematic of lower and middle atmosphere showing thermal and dynamical structure [Adapted from Meriwether and Gerrard, 2004].	5
1.2	Vertical distribution of heating rates due to absorption of solar short wave radiation [Adapted from <i>London (1980)</i>].	7
1.3	Zonal mean temperature from SABER retrievals and from WACCM [Adapted from <i>Smith (2012a)</i>].	9
1.4	Zonal mean zonal wind from the URAP climatology and simulated by WACCM [Adapted from <i>Smith (2012a)</i>].	11
1.5	A Summary of atmospheric waves [Adapted from <i>Lin (2007)</i>].	13
1.6	A surrealistic representation of atmospheric gravity waves [Adapted from <i>Hines (1974)</i>].	16
1.7	Vertical distribution of the amplitude of the MQBO, MSAO, SSAO, SQBO, and annual component at the equator [Adapted from <i>Baldwin et al. (2001)</i>].	27
1.8	Temperature and zonal wind for 2013 SSW event	29
1.9	Schematic of upper and lower mesospheric inversion layers [Adapted from <i>Meriwether and Gerrard (2004)</i>].	32
1.10	Red, green, and yellow airglow emission from the Earth's upper atmosphere photographed by the International Space Station. (http://auroranightglow.blogspot.in/2012/05/night-glow.html)	37

1.11	Schematic of the peak emission altitudes for the various nightglow emissions in the MLT region.	38
1.12	Schematic of the simplified energy level diagram along with various emissions of the atomic oxygen.	40
2.1	(a) Optical layout of Near InfraRed Imaging Spectrograph (NIRIS) [<i>Singh and Pallamraju (2017a)</i>].	51
2.2	Schematic of a reflection grating.	53
2.3	NIRIS obtained spectral images	56
2.4	(a) View angles calibration of NIRIS.	58
2.5	A nocturnal spectral image and spectral lines obtained by NIRIS	60
2.6	Nocturnal variation of three different rotational lines of OH(6-2) Meinel band and O ₂ (0-1) atmospheric band emissions	61
2.7	Nocturnal variations in O ₂ and OH intensities and rotational temperatures on the night of 16 April 2013 and 04 May 2013	66
2.8	Vertical profile of SABER and NIRIS derived temperatures	68
2.9	Optical configuration and photograph of CMAP	70
2.10	The filter wheel assembly and bipolar temperature controller along with Peltier elements to control the temperature	73
2.11	Typical temperature stability achieved in our in-house developed temperature controlled chamber used in CMAP	74
2.12	Nocturnal variations in the OI(630.0), OI(557.7 nm), and sodium (589.0 and 589.6 nm) nightglow emission intensities as measured by CMAP at Gurushikhar, Mount Abu on the night of 24-25 October 2014	76
2.13	Real part (solid) and imaginary part (dashed) for the Morlet wavelet in the time domain.	87
2.14	Periods that are present in nocturnal variation in O ₂ and OH emission intensities	90

2.15	Nocturnal variation in I(O ₂) and I(OH) intensities and Lomb-Scargle periodograms of the residuals	91
2.16	Residuals of the nightglow emission intensities for both O ₂ and OH emissions showing downwards phase propagation which indicate an upwards propagation of gravity wave	92
2.17	(Variations in the O ₂ and OH intensities along N-S meridian and statistically significant meridional wavelengths for O ₂ and OH intensity variation throughout the night of 4 May 2013	94
2.18	The wavelet transform of the residuals of I(O ₂) and I(OH) showing occurrence of the same periodicities as seen from Lomb-Scargle method along with their temporal occurrences	96
2.19	A schematic of the propagation of gravity wave in which the phase is shown to be progressing downwards (upward propagating wave)	97
3.1	Details of the duration of observations obtained from the Near InfraRed Imaging Spectrograph (NIRIS) at Gurushikhar, Mount Abu (24.6°N, 72.8°E), in India.	103
3.2	Variations in nightly mean O ₂ and OH intensities, daily mean sunspot numbers and F10.7 cm radio flux, and corresponding O ₂ and OH temperatures	106
3.3	Lomb-Scargle periodograms of the O ₂ and OH intensities and corresponding temperatures and solar parameters for different durations	108
3.4	Four-day running averages of O ₂ and OH intensities and SSN and correlation coefficients in between these parameters	111
3.5	The procedure used to derive gravity wave periodicities from the nocturnal variations of the nightglow intensity data is shown for a sample night of 10 June 2015	116

3.6	Gravity wave periodicities obtained from the O ₂ and OH emission intensities for all the 437 nights of observations are shown which are folded into a one year interval	118
3.7	Same as in Figure 3.6 but for the temperatures obtained from the O ₂ (0-1) atmospheric and OH(6-2) band emissions intensities.	120
3.8	Statistical results of the GW occurrence frequency observed by all sky camera [Adapted from <i>Wu and Killeen (1996)</i>].	123
3.9	Same as in Figures 3.6 and 3.7 (a, b, d, e) but for the periods that have been derived from only those nights of observation for which the total duration of observation was greater than 6 h . . .	124
4.1	Time sequence of difference OH images in geographic coordinates from 0340 to 0500 UT on 11 May 2004 [Adapted from <i>Yue et al. (2009)</i>]	132
4.2	Outgoing longwave radiation (OLR) on 26 October 2014 as derived from VHRM on-board Indian meteorological satellite (Kalpana-1)	136
4.3	Nocturnal variations in the OI (557.7 nm), O ₂ (0-1) atmospheric band, Na (589.3 nm), and OH(6-2) Meinel band nightglow emission intensities as measured by CMAP and NIRIS at Gurushikhar, Mount Abu	138
4.4	Nocturnal variation in OI, O ₂ , Na, and OH nightglow emission intensities on the night of 26 October 2014. Variation of OLR values at the location ‘P’ for 24-hour duration	141
4.5	Lomb-Scargle periodograms of the zero mean unit variance for the parameters shown in Figures 4.4 for the night of 26 October 2014.	143
4.6	Periods obtained using Lomb-Scargle analyses for OI, O ₂ , Na, and OH nightglow emission intensities and in OLR(P) data obtained for each night during 16–27 October 2014.	145

4.7	Normalized wavelet spectra of the corresponding zero mean unit variances of the O ₂ , Na, and OH nightglow emission intensities as shown in Figure 4.4 and OLR(CR1) showing temporal evolution of the periods seen in Figure 4.5	148
4.8	Schematic showing the wave propagation direction and distance from the source region to the observational location.	150
4.9	Nocturnal variation in meridional and zonal wind at 94 km altitude as obtained from HWM14 and in vertical wavelength calculated using GW linear dispersion relation with the HWM14 winds	152
4.10	TIDI (TIMED Doppler Interferometer) winds showing the presence of shear in the zonal/ meridional winds.	154
4.11	The difference in T(O ₂) and T(OH) on the night of (a) 16 April 2013, and (b) 6 May 2013	159
4.12	The five rows in the different subplots represent the difference in T(O ₂) and T(OH), I(O ₂) and I(OH), residuals of O ₂ and OH emission intensities, and the periodicities present in the O ₂ and OH emission intensities, respectively.	162
5.1	NIRIS derived O ₂ and OH rotational temperatures over Gurushikhar, Mount Abu for a few selected nights during 2013 SSW period.	170
5.2	Temperatures in the polar cap poleward of 60°N and longitudinally averaged zonal wind at 60°N at 10 hPa pressure level obtained from MERRA dataset for eleven SSW events which occurred during 2004–2013	173
5.3	Longitudinally averaged mesospheric temperatures derived from the SABER, SOFIE, and OSIRIS dataset at 10° latitude interval for 2009 and 2013 SSW events	176
5.4	Similar to Figure 5.3 but for 2006 major SSW event.	177

5.5	Similar to Figure 5.3 but for 2010 major SSW event.	178
5.6	Similar to Figure 5.3 but for 2011 minor SSW event.	179
5.7	Similar to Figure 5.3 but for 2012 minor SSW event.	180
5.8	Similar to Figure 5.3 but for 2008 SSW event.	181
5.9	Differences in SSW temperatures (ΔT_S) over 60° – 90° N at 10 hPa pressure level vs difference in mesospheric temperatures (ΔT_M) for six selected latitude ranges for 2013 and 2009 SSW event as derived from SABER datasets	183
5.10	Difference in SSW temperatures (ΔT_S) over 60° – 90° N at 10 hPa pressure level vs difference in mesospheric temperatures (ΔT_M) at 10° latitude intervals as derived from SABER dataset from both the hemispheres for three major and five minor SSW events as shown in Figure 5.2	184
5.11	Same as in Figure 5.9 but for SOFIE data at two latitudinal ranges in NH and SH for three major SSW events of 2009, 2010, and 2013 combined together and for OSIRIS data for two major SSW events of 2009 and 2010	187
5.12	Slopes ($\Delta T_M/\Delta T_S$) derived from analyses for each individual major SSW event using SABER data with respect to the geographic latitude for all major and minor SSW events	189
5.13	Peak stratospheric temperature for all the eleven selected SSW events with slope reversal latitudes in the poleward direction in both hemispheres. Latitudinal difference in peak slopes of the “double-humped” structure with respect to the peak stratospheric temperatures are also shown	191
5.14	Daily mean mesospheric temperatures derived from SABER data for all the eleven SSW events with respect to geographic equator and CIRA-86 derived temperatures. The differences in the SABER measured and CIRA-86 estimated mesospheric temperature are also shown	193