

LIST OF FIGURES

Figure	Title	Page No
1.1	Structure of polyether sulfone	3
1.2	a) Sterical exclusion; b) Donnan exclusion	11
2.1	Polyelectrolytes used for coating	48
2.2	Scheme of layer-by-layer adsorption of polyelectrolytes on porous supporting membrane	49
2.3	JEOL JSM-6390 scanning electron microscope.....	51
2.4	Amicon 8050 ultrafiltration cell	53
3.1	Polyelectrolytes used for coating	65
3.2	SEM images of supor and 7 bilayer CHI/PSS membranes.....	66
3.3	FT-IR of supor membrane and CHI/PSS 0, 3,5,6,7 bilayers at pH 1.7 (sulfonate region, 1033cm^{-1})	67
3.4	Percentage transmission of BSA as a function of number of CHI/PSS bilayers when the multilayers are fabricated at two different pHs 1.72 and 2.12	70
3.5	BSA solution flux as a function of number of CHI/PSS bilayers when the multilayers are fabricated at two different pHs 1.72 and 2.12.....	70
3.6	Area under the sulfonate peak (1033 cm^{-1}) in the FT-IR spectrum as a function of the number of deposited layers at two different pHs 1.72 and 2.12	72
3.7	Percentage transmission of BSA as a function of number of CHI/PSS bilayers when the multilayers are fabricated in the presence and the absence of salt.....	75
3.8	BSA solution flux as a function of number of CHI/PSS bilayers when the multilayers are fabricated in the presence and the absence of salt.....	76

3.9	The peak area at 1033 cm^{-1} in the FT-IR spectrum as a function of the number of bilayers when multilayers are fabricated from CHI (medium MW)/PSS and CHI (high MW) /PSS.....	79
3.10	Percentage transmission of BSA as a function of the number of bilayers for BSA solutions of different pHs., 8.8, 4.8 and 3.7	83
3.11	BSA solution flux as a function of the number of bilayers for BSA solutions of different pHs., 8.8, 4.8 and 3.7	83
3.12	Percentage transmission of ovalbumin as a function of the number of bilayers for ovalbumin solutions of different pHs, 8.8 and 4.5	86
3.13	Ovalbumin solution flux as a function of the number of bilayers for ovalbumin solutions of different pHs, 8.8 and 4.5	87
3.14	Mass balance of solute on the membrane boundary layer	88
3.15	Percentage transmission of lysozyme as a function of the number of bilayers for lysozyme solutions of different pHs., 10.6, 8.8 and 4.5	94
3.16	Lysozyme solution flux as a function of the number of bilayers for lysozyme solutions of different pHs., 10.6, 8.8 and 4.5	94
3.17	Single protein transmission profile of BSA, ovalbumin and lysozyme at pH 8.8	96
3.18	Effect of ionic strength on the percentage transmission of BSA (0.5 mg/ml, pH 8.8), ovalbumin (0.5 mg/ml, pH 8.8) and lysozyme (0.25 mg/ml, pH 8.8) through CHI/PSS multilayer membranes	101
3.19	Effect of ionic strength on the solution flux of BSA (0.5 mg/ml, pH 8.8), ovalbumin (0.5 mg/ml, pH 8.8) and lysozyme (0.25 mg/ml, pH 8.8) through CHI/PSS multilayer membranes	102
3.20	Percentage transmission Vs NaCl concentration for BSA and ovalbumin with 9 bl CHI/PSS membrane	107

3.21	Solution flux Vs NaCl concentration for BSA and ovalbumin with 9 bl CHI/PSS membrane	108
3.22	Comparison of percentage transmission and NaCl concentration for ovalbumin through 5 bl and 9 bl CHI/PSS membranes.....	108
3.23	Comparison of flux and NaCl concentration for ovalbumin through 5 bl and 9 bl CHI/PSS membranes	109
3.24	Percentage transmission of BSA Vs Concentration of BSA (CHI/ PSS 8 th , 9 th , 10 th bilayer).....	116
3.25	BSA flux Vs Concentration of BSA (CHI/PSS 8 th , 9 th , 10 th bilayer)....	116
3.26	Transmission profile of BSA of different concentration through CHI/PSS multilayers	117
3.27	Structure of amino acids used for permeation study.....	121
3.28	Transmission profile of L-histidine through CHI/PSS multilayer membrane.....	124
3.29	Transmission profile of glycine through CHI/ PSS multilayer membrane	126
3.30	Transmission profile of L-lysine through CHI/PSS multilayer membrane.....	127
3.31	FT-IR spectrum of lysine adsorbed 5 bl CHI/PSS membranes	129
3.32	Transmission profile of L-aspartic acid through CHI/PSS multilayer membrane.....	131
3.33	Transmission profile of lysine, aspartic acid and histidine through CHI/PSS multilayer membrane at pH 5	132
4.1	pH dependent loading of lysozyme (0.25 mg/ml) to 5 bl CHI/PSS under UF conditions. Amide I band in the FT-IR spectrum is shown	148
4.2	Schematic representation of the mechanism of protein immobilization onto supor membrane <i>via</i> an intermediate adsorbed CHI/PSS multilayers.....	150

4.3	FT-IR spectrum of lysozyme adsorbed (UF) at pH 10.6 on 5 bl CHI/PSS membrane showing amide I, amide II and amide II' bands.....	152
4.4	SEM pictures of lysozyme adsorbed 5 bl CHI/PSS membranes under UF conditions	154
4.5	AFM images of lysozyme adsorbed CHI/PSS 5 bl membranes (UF) conditions.....	155
4.6	pH dependent loading of lysozyme (0.25 mg/ml) to CHI/PSS multilayer under simple adsorption conditions. Amide I band (1650 cm^{-1}) in the FT-IR spectrum is shown.....	156
4.7	FT-IR spectrum showing the amide I band in time dependent loading of lysozyme (0.25 mg/ml; pH 5) to CHI/PSS multilayer (simple adsorption).....	159
4.8	Time dependent loading of lysozyme (0.25 mg/ml; pH 5) to CHI/PSS multilayer. The amide I band peak area in the FT-IR spectrum at 1650 cm^{-1} is plotted as a function of time	160
4.9	Time dependent loading of lysozyme (0.25 mg/ml; pH 5) to CHI/PSS multilayer. The percentage absorption of lysozyme solution at 280 nm is plotted as a function of time.....	160
4.10	SEM pictures of lysozyme (pH 5) adsorbed membranes (under simple adsorption conditions)	162
4.11	BSA (pH 4.8) adsorbed membranes with varying number of bilayers (simple adsorption). Amide I band (1650 cm^{-1}) in the FT-IR spectrum is shown.....	163
4.12	BSA (pH 4.8) adsorbed membranes with varying number of bilayers under UF conditions. Amide I band (1650 cm^{-1}) in the FT-IR spectrum is shown.....	164
4.13	SEM pictures of BSA (pH 4.8) adsorbed membranes under UF conditions	165
4.14	pH dependent loading of BSA to 5 bl CHI/PSS membrane under simple adsorption conditions. Amide I band in the FT-IR spectrum is shown	166

4.15	Time dependent loading of BSA pH 4.8 to 5 bl CHI/PSS membrane (simple adsorption). Amide I band in the FT-IR spectrum is shown	168
4.16	Time dependent loading of BSA (0.5 mg/ml; pH 4.8) to 5 bl CHI/PSS membrane. The amide I band (1650 cm ⁻¹) peak area in the FT-IR spectrum is plotted as a function of time.....	168
4.17	Time dependent adsorption of BSA (0.5 mg/ml; pH 4.8) to 5 bl CHI/PSS membrane. The percentage absorbance of BSA at 280 nm obtained from UV absorbance of protein solution before and after adsorption is plotted as a function of time	169
4.18	Time-dependent desorption of lysozyme (pH 8.8) from 5 bl membrane. The UV absorbance of the desorbate at 280 nm is recorded at different time intervals	172
4.19	Time-dependent desorption of lysozyme (pH10.6) from 5 bl membrane. The UV absorbance of the desorbate at 280 nm is recorded at different time intervals	172
4.20	Desorption of lysozyme (pH 8.8) from adsorbed 5 bl membrane by salt of varying concentrations.....	174
4.21	Desorption of lysozyme (pH10.6) from adsorbed 5 bl membrane by salt of varying concentration	175
4.22	UV absorbance of lysozyme (pH 10.6) adsorbed and desorbed membranes	176
4.23	SEM images of lysozyme desorbed membranes with 0.5M NaCl a: lysozyme pH 8.8, b: lysozyme pH 10.6	177
4.24	Ion-exchange-type displacement, where salt counter ions replace adsorbed proteins in high ionic strength buffers.....	177
4.25	Time-dependent desorption of lysozyme pH (8.8) from 5 bl CHI/PSS membrane with 0.5 M NaCl	179
4.26	Time dependent desorption of lysozyme (0.25 mg/ml; pH 8.8) from 5 bl CHI/PSS membrane with 0.5 M NaCl	179
4.27	Time-dependent desorption of lysozyme pH (10.6) from 5 bl membrane with 0.5 M NaCl.....	180

4.28	Fluorescence spectra of lysozyme adsorbed (UF) 5bl CHI/PSS membrane at pH 10.6, pH 8.8 and pH 5.....	187
4.29	Fluorescence spectra of lysozyme (pH 10.6) adsorbed (UF) 5bl CHI/PSS membranes with increasing concentration of iodide ion	188
4.30	Variation of fluorescence intensity of lysozyme (pH 10.6) adsorbed (UF) 5bl CHI/PSS membranes with quencher concentration	188
4.31	Perrin plot for lysozyme (pH 10.6) adsorbed (UF) 5bl CHI/PSS membranes	191
4.32	Perrin plot for lysozyme (pH 5) adsorbed (UF) 5bl CHI/PSS membranes.....	192
4.33	Perrin plot for lysozyme (pH 5) adsorbed (simple adsorption) 5bl CHI/PSS membranes.....	192