

List of Tables

3.1	Energy of the first excited state relative to the ground state (in MeV) for oxygen isotopes for different interactions.	33
3.2	Energy difference between the first excited state and the ground state (in MeV) for four different interactions in fluorine.	33
3.3	$B(E2; 2_1^+ \rightarrow 0_{gs}^+)$ in $e^2 \text{ fm}^4$: for different interaction in oxygen isotopes.	34
3.4	The proton single-particle energies for different interactions.	36
4.1	Dimensions of matrices involved for $^{62-66}\text{Fe}$ in the m scheme for fp shell	50
4.2	Main configurations in the wave functions of the ground state and the first excited state for $^{62-66}\text{Fe}$ calculated with GXPF1A interaction.	54
4.3	Excitation energies, quadrupole moments and $B(E2)$ for $^{62-66}\text{Fe}$ isotopes.	54
4.4	Main configurations in the wave functions of the ground state and the first excited state for $^{61-65}\text{Fe}$ calculated with GXPF1A interaction.	57
5.1	Single particle energies used in GXPF1A, KB3G and fpg interactions in MeV.	60
5.2	Experimental and theoretical low-lying states (up to 2 MeV) of ^{58}Mn . Energies are in MeV. The results of GXPF1 interaction are taken from Ref. [Lid06].	63
5.3	Experimental and theoretical low-lying states (up to 1.6 MeV) of ^{60}Mn . Energies are in MeV. The results of GXPF1 interaction taken from Ref. [Lid06] [NNDC].	66
5.4	Experimental and theoretical low-lying states (up to 1 MeV) of ^{62}Mn . Energies are in MeV.	66
5.5	Main configurations in the wave functions of the ground state and the first excited state for ^{58}Mn for different interactions.	67

5.6	Main configurations in the wave functions of the ground state and the first excited state for ^{60}Mn for different interactions.	67
5.7	Main configurations in the wave functions of the ground state and the first excited state for ^{62}Mn for different interactions.	67
7.1	M-scheme dimensions as a function of jumps for $^{69-79}\text{Cu}$ isotopes. . .	94
7.2	M-scheme dimensions as a function of jumps for $^{68-78}\text{Ni}$ isotopes. . .	95
7.3	M-scheme dimensions as a function of jumps for $^{70-80}\text{Zn}$ isotopes. . .	96
7.4	B(E2) value for $^{69-79}\text{Cu}$ isotopes . The results in the bracket from earlier work of Stefanescu <i>et al.</i> [Ste08].	97