

LIST OF TABLES

- Table 3.1** : Name of the primers with sequences used in this study
- Table 3.2** : Segregation type codes for population type CP and genotype codes for a CP population, depending on the locus segregation type
- Table 3.3** : Genes selected for the real time quantification studies and primer sequences for gene specific amplification
- Table 4.1.1** : Parental difference of caffeine and flavanols in the three populations at 99% confidence level
- Table 4.1.2** : Estimates of variance (eigenvalues) and accumulated variances of parental characters
- Table 4.1.3** : Available putative triploid populations of tea and their parental combinations
- Table 4.2.1** : Possible genomic combination of triploid parent (29/2) of the allo-tetraploid 398/2.A and B of triploid 29/2 denotes separate genome. The Diploid gamete “A” shows allele size similarity with “Genome A” of 29/2. The diploid gamete “a” shows allele size similarity with “Genome B” of 29/2.
- Table 4.2.2** : Segregation pattern of the loci heterozygous in the diploid parent
- Table 4.2.3** : Segregation pattern of the loci homozygous in the diploid parent
- Table 4.2.4** : Segregation pattern of locus heterozygous in both the parent: Two Alleles are segregating
- Table 4.2.5** : Segregation pattern of locus heterozygous in both the parents: Three alleles are segregating
- Table 4.2.6** : Pearson’s correlation table showing the relationship between the GD (Jaccard’s) between parents and progenies and the biochemical content of the progenies.
- Table 4.2.7** : The coefficient of correlation and corresponding probability values (-10 LogP) of each of the amplicons in Stock 615.
- Table 4.2.8** : The coefficient of correlation and corresponding probability values (-10 LogP) of each of the amplicons in Stock 616
- Table 4.4.1** : Some of the important sequences selected for RT-PCR analysis