

REFERENCES

- [1] Kostanecki, SV & Tambor, J 1899, ‘Mmonoxychalkone’, Journal of Chemischen Berichte, vol. 34, no. 3, pp.1921-1926.
- [2] Rupe, H & Wasser Zug, D, ‘Notes on Chromophore grouping’, Journal of Chemischen Berichte, vol. 32, no. 2, pp.3527-3531.
- [3] Rao, SR & Ravishankar, GA 2002, ‘Plant cell cultures: Chemical factories of secondary metabolites’, Biotechnology Advances, vol. 20, no. 2, pp. 101-153.
- [4] Cowan, MM 1999, ‘Plant products as antimicrobial agents’, Clinical Microbiology Review, vol. 12, no. 4, pp. 564-582.
- [5] Basics of Scientific Literature Review, Available from: <https://en.wikipedia.org/wiki/Literature_review>.
- [6] Parlak, C, Ramasami, P, Sadolalu, C, Chidan Kumar, CS, Mahir Tursun, Ching Kheng Quah, Lydia Rhyman, Metin Bilge, Hoong-Kun Fun & Siddegowda Chandraju 2015, ‘(2E)-1-(5-Chlorothiophen-2-yl)-3-{4-[{(E)-2-phenylethenyl]} phenyl prop-2-en-1-one’, Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy, vol. 149, pp. 385-395.
- [7] Chidan Kumar, CS, Govindarasu, K, Hoong-Kun Fun, Kavitha, E, Siddegowda Chandraju & Ching Kheng Quah 2015, ‘Synthesis, molecular structure, spectroscopic characterization and quantum chemical calculation studies of (2E)-1-(5-Chlorothiopene-2-yl)-3-(2, 3, 4-trimethoxyphenyl) prop-2-en-1 one’, Journal of Molecular Structure, vol. 1085, pp. 63- 77.
- [8] Rajesh, K, Amit, K, Vipin, D, Archana, G, Poonam, T, Patil, PS, Prathmesh, D, Deepika, C & Watve JG 2017, ‘Molecular structure, spectroscopic (FT-IR, FT-Raman, UV, NMR and THz) investigation and hyperpolarizability studies of 3 (2-Chloro-6-fluorophenyl)-1- (2-thienyl) prop-2-en-1-one’, Journal of Molecular Structure, vol. 1129, no. 1, pp 292-304.
- [9] Subhendu, C, Joydeep, C, Manash, G & Talapatra, GB 2012, ‘Genesis of enhanced Raman bands in SERS spectra of 2-mercaptoimidazole: FTIR, Raman, DFT, and SERS’, Journal of Physical Chemistry, vol. 116, no. 2, pp. 10934-10947.

- [10] Ghomrasni, S, Aribi, I, Ayachi, S, Haj Said, A & Alimi, K 2015, 'Molecular structure – optical property relationships of 1, 3-bis (4-methoxyphenyl) prop-2 en-1-one: A DFT and TD-DFT investigation', Journal of Physics and Chemistry and Solids, vol. 83, pp. 157-168.
- [11] Kieninger,M, Cachau, E, Oberhammer, H & Oscar, N 2007, 'Comparison of large basis set DFT and MP2 calculations in the study of the barrier for internal rotation of 2,3,5,6 tetrafluoroanisole', International Journal of Quantum Chemistry, vol. 107, no.2 , pp. 403-417.
- [12] Siva Priya, M, AsenathBenitta, T & James C 2011, 'Vibrational spectroscopic investigation and normal coordinate analysis of the fibrate hypolipidemic agent 5-(2, 5-dimethylphenoxy)-2, 2-dimethylpentanoic acid (Gemfibrozil)', Journal of Molecular Structure, vol. 990, no. 1, pp. 253-262.
- [13] Suhana, A, Renjith, RP, DuanAlwani, Z, NuridayantiChe, K, Ibrahim, AR, Stevan, A, Sanja, JA, Yohannan, C & Van Alsenoy, C 2017, 'Synthesis, crystal structure, Hirshfeld surface analysis, spectroscopic characterization, reactivity study by DFT and MD approaches and molecular docking study of a novel chalcone derivative', Journal of Molecular Structure, vol.1135, pp. 234-246.
- [14] Abdel Latif Samir, A & Mohamed Adel, A 2016, 'Synthesis, structure, spectroscopic properties and DFT studies on some 7-hydroxy 4-methyl-8 (arylazo)-2H-1-benzopyran-2-one and their complexes with some divalent transition metal ions', Journal of Molecular Structure, vol. 1134, pp. 307-318.
- [15] Chidan Kumar, CS, Balachandran, V, Hoong-Kun Fun, Siddegowda, C & Ching Kheng Quah 2015, 'Synthesis, crystal growth, single crystal X-ray analysis and vibrational spectral studies of (2E) -3-(2-chloro-4-fluorophenyl)-1-(3,4 dimethoxyphenyl) prop-2-en-1-one: A combined DFT study', Journal of Molecular Structure, vol. 1100, no. 1, pp. 299-310.
- [16] Jose, C, Alejandro, A, Norma, R, Guadalupe, V, Daniel Glassmon, M & Luz, M 2013, 'Experimental and quantum mechanical studies of a novel synthetic prenylated chalcone', Chemistry Central Journal, vol. 7, pp. 17-28.
- [17] McKinnon, JJ, Jayatilaka, D & Spackman, MA 2007, 'Towards quantitative analysis of intermolecular interactions with Hirshfeld surfaces', Chemical Communications, vol. 37, no. 1, pp. 3814-3816.

- [18] Rohl, AL, Moret, M, Kaminsky, W, Claborn, K, McKinnon, JJ & Kahr, B 2008, ‘Hirshfeld surfaces identify inadequacies in computations of intermolecular interactions in crystals: pentamorphic 1, 8-dihydroxyanthraquinone’, *Crystal Growth and Design*, vol. 8, no.12, pp. 4517-4519.
- [19] Luo, YH, Zhang, CG, Xu, B & Sun, BW 2012, ‘A cocrystal strategy for the precipitation of liquid 2,3-dimethyl pyrazine with hydroxyl substituted benzoic acid and a Hirshfeld surfaces analysis of them’, *Crystal EngineeringCommunication*, vol. 14, no. 20, pp. 6860-6864.
- [20] Venkatesan, P, Thamotharan, S, Ilangovan, A, Hongze Liang& Tom Sundius 2015, ‘Crystal structure, Hirshfeld surfaces and DFT computation of NLO active (2E)-2(ethoxycarbonyl)-3-[(1-methoxy-1-oxo-3-phenylpropan-2-yl) amino] prop-2-enoic acid’, *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, vol. 153, pp. 625-36.
- [21] Brancale, A & Silvestri, R 2007, ‘Core nucleus for protein inhibitor of tubulin polymerization’, *Journal of Medicinal Research Review*, vol. 27, no. 2, pp. 209-238.
- [22] Yohannan Panicker, C, Hema Tresa, V, Prakash, S, Narayana, B, Sarojini, BK, Fun, HK, Javeed, A, Srivastava, SK & Van Alsenoy, C 2015, ‘Infrared spectrum, NBO, HOMO–LUMO, MEP and molecular docking studies (2E)-3-(3-nitrophenyl)-1-[4-piperidin-1-yl]prop-2-en-1-one’, *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, vol. 148, pp. 18-28.
- [23] Cemal Parlak, Ramasami, P, Chidan Kumar, CS, Mahir, T, Ching Kheng Quah, Lydia, R, Metin Bilge, Hoong-Kun Fun & Chandrapu, S 2015, ‘FT-IR spectrum of (2E)-3-(3-nitrophenyl)-1-[4-piperidin-1-yl]prop-2-en-1-one’, *Spectrochimica Acta Part A: Molecular and Bio molecular Spectroscopy*, vol.149, no.1, pp. 385-395.
- [24] Lynnette, J, Sajan, D, Shettigar, V, Chaitanya, K, Neeraj Misra, Sundius, T & Nemec, I 2013, ‘Density Functional Study on the Structural Conformations and Intramolecular Charge Transfer From the Vibrational Spectra of 1-(4- aminophenyl)3-(3,4-dimethoxyphenyl)-prop-2-en-1-one’, *Materials Chemistry and Physics*, vol. 141, no. 1, pp. 248-262.
- [25] Jasinski, JP, Pek, AE, Narayana, B, Kamath, PK & Yathirajan, HS 2010, ‘(E)-1-(4 Bromophenyl)-3-(2-methoxyphenyl) prop-2-en-1-one’, *Acta*

Crystallographica section E: Structure Reports Online, vol. 66, no. 8, pp. 1991-1996.

- [26] Jasinski, JP, Butcher, RJ, Narayana, B, Veena, K&Yathirajan, HS 2010, '(2E)-1 (3-Chlorophenyl) 3-(4-chlorophenyl) prop-2-en-1-one', Acta Crystallographica Section E: Structure Reports Online, vol. 66, pp. 1638-1639.
- [27] Brunck, TK & Weinhold, F 1979, 'Quantum-mechanical studies on the origin of barriers to internal rotation about single bonds', Journal of American Chemical Society, vol. 101, no.7, pp. 1700 -1709.
- [28] Baran, J, Trzebiatowska, M & Ratajczak, H 2004, 'Studies in the Vibrational Spectroscopy of molecular crystals and liquid crystals', Journal of Molecular Structure, vol. 708, no. 1-3, pp. 127-144.
- [29] Pijun Liu, Liyue Liu & Yafei Zhang 2003, 'Alignment characterization of single wall carbon nanotubes by Raman scattering', Physics Letters A, vol. 313, pp. 302-306.
- [30] Jeremy Jones, W, Thomas, DK, Thomas, DW & Williams, G 2004, 'On the determination of order parameter for homogeneous and twisted nematicliquid crystals from Raman spectroscopy', Journal of Molecular Structure, vol. 708, no. 1, pp. 145-163.
- [31] Spiro, TG & Strekas, TC 1972, 'Resonance Raman spectra of hemoglobin and Cytochrome C: Inverse polarization and vibronic scattering', Proceedings of the National Academy of Sciences of the United States of America, vol. 69, no. 9, pp. 2622-2626.
- [32] Walrafen, GE 2005, 'Dispersion of the Raman depolarization ratio of HDO in water and heavy water from 295 to 368 K, and from concentrated (NaClO₄22O)-O-/D-O-O/H', Journal of Chemical Physics, vol. 122, no.17, pp. 174502-174509.
- [33] Rafailor, PM, Hadjiev, VG, Jantoljak, H & Thomsen, C 1999, 'Raman depolarization ratio of vibrational modes in solid C₆₀', Solid State Communications, vol. 112, no. 9, pp. 517- 520.

- [34] Gi Xue, 1997, ‘Fourier transform Raman spectroscopy and its application for the analysis of polymeric materials’, Progress in Polymer Science, vol. 22, no. 2, pp. 313-406.
- [35] Gurvinder singh, B & Rakesh Mohan, S 2016, ‘Raman Spectroscopy-Basic principles, instrumentation and selected applications for the characterization of drugs of abuse’, Egyptian Journal of Forensic Sciences, vol. 6, no. 3, pp. 209-215.
- [36] Tony Owen 1996, Fundamentals of UV-Visible Spectroscopy, Hewlett Packard Company, Germany.
- [37] Clark, BJ, Frost, T & Russel, MA 1993, UV spectroscopy: techniques, instrumentation and data handling, Chapman & Hall, London.
- [38] Akul Mehta, 2011, Ultraviolet-visible (UV-Vis) spectroscopy – principle. Available from : <<https://pharmaxchange.info/2011/12/ultraviolet-visible-uv-vis-spectroscopy-principle/>>
- [39] David Young 2001, Computational chemistry, John Wiley & Sons, Inc., New York.
- [40] Frank Jensen 2007, Introduction to computational chemistry, Wiley & Sons Ltd., Denmark.
- [41] Weinhold, F & Landis, CR, Valency and bonding: A natural bond orbital donor-acceptor perspective, Cambridge University Press, Cambridge.
- [42] Montejo, M, Navarro, A, Kealey, GJ, Vazquez, J & Gonzalez, JJL 2004, ‘Inter molecular hydrogen bonding and charge transfer in solid furan’, Journal of American Chemical Society, vol. 126, no. 46, pp. 15087-15095.
- [43] Reed, AE, Curtiss, LA & Weinhold, F 1988, ‘Inter molecular interactions froma natural bond orbital, donor-acceptor view point’, Chemical Reviews, vol. 88, no. 6, pp. 899-926.

- [44] Sheena Mary, Y, Nasser El-Brollosy, R, Ali El-Emam, A, Omar Al- Deeb, A, Jojo, PJ, YohannanPanicker, C & Christian Van Alsenoy 2014, ‘Vibrational spectra, NBO analysis, HOMO–LUMO and first hyperpolarizability of 2-{[(2-Methylprop-2-en-1-yl)oxy]methyl}-6 phenyl-2,3,4,5-tetrahydro-1,2,4-triazine-3,5-dione,a potential chemotherapeutic agent based on density functional theory calculations’, SpectrochimicaActa Part A: Molecular and Biomolecular Spectroscopy, vol. 133, no. 14, pp. 449-456.
- [45] Schachtschneider, JH & Snyder, RG 1963, ‘Vibrational analysis of the n paraffins—I: Assignments of infrared bands in the spectra of C₃H₈ through n-C₁₉H₄₀’, Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy, vol. 19, no.1, pp. 85-116
- [46] Gwinn, WD 1971, ‘Normal Coordinates: General Theory, Redundant Coordinates, and General Analysis Using Electronic Computers’, Journal of chemical physics, vol. 55, no. 1 , pp. 477- 481.
- [47] Wilson, EB, Decius, JC & Cross, PC 1955, Molecular vibrations, Mc Graw Hill, New York.
- [48] Sundius, T 1980, ‘A new damped least-squares method for the calculation of molecular force fields’, Journal of Molecular Spectroscopy, vol. 82, no. 1, pp. 138-151.
- [49] Sundius, T 2002, ‘Scaling of ab initio force fields by MOLVIB’, Vibrational Spectroscopy, vol. 29, no. 1, pp. 89-95.
- [50] Fukui, K 1982, ‘Role of Frontier Orbitals in Chemical Reactions’, Science, vol. 218, no. 4, pp.747-754.
- [51] Ghatak, KL 2014, Text book of organic chemistry and problem analysis, PHI Learning Pvt. Ltd., Uitgewer.

- [52] McKinnon, JJ, Spackman, MA & Mitchell, AS 2004, ‘Novel tool for visualizing and exploring intermolecular interactions in molecular crystals’, *Acta Crystallographic B*, vol. 60, no. 6, pp. 627-668.
- [53] Hirschfeld, FL 1977, ‘Bonded-atom fragments for describing molecular charge densities’, *Theoretica Chimica Acta*, vol. 44, no. 2, pp. 129-138.
- [54] McKinnon, JJ, Jayatilaka, D & Spackman, MA 2007, ‘Towards quantitative analysis of intermolecular interactions with Hirshfeld surfaces’, *Chemical Communication*, vol. 37, no. 2, pp. 3814-3816.
- [55] Spackman, MA & McKinnon, JJ 2002, ‘Finger printing intermolecular interactions in molecular crystals’, *Journal of Crystal Engineering Communication*, vol. 4, no. 66, pp. 378-392.
- [56] Morris, GM, Huey, R, Lindstrom, W, Sanner, MF, Belew, RK, Goodsell, DS & Olson, AJ 2009, ‘AutoDock4 and AutoDockTools4: Automated docking with selective receptor flexibility’, *Journal of Computational Chemistry*, vol. 30, no. 16, pp. 2785-2791.
- [57] Trott, O & Olson, AJ 2010, ‘Auto Dock Vina: improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading’, *Journal of Computational Chemistry*, vol. 31, no. 2, pp. 455-461.
- [58] Gatti, C, Saunders, V & Roetti, C 1994, ‘Crystal field effects on the topological properties of the electron density in molecular crystals: the case of urea’, *Journal of Chemical Physics*, vol. 101, no. 12, pp. 10686-10696.
- [59] Grabowski, SJ 2000, ‘High-level ab initio calculations of di-hydrogen bonded complexes’, *Journal of Chemical Physics*, vol. 104, no. 23, pp. 5551-5557.
- [60] Espinosa, E, Molins, E & Lecomte, C 1998, ‘Hydrogen bond strengths revealed by topological Analyses of experimentally observed electron densities’, *Chemical Physics Letter*, vol. 285, no. 3, pp. 170-173.

- [61] Nantasenamat, C, Isarankura-Na-Ayudha, C & Pracayasittikul, V 2010, ‘Advances in computational methods to predict the biological activity of compounds’, Expert Opinion on Drug Discovery, vol. 5, no.7, pp. 633-654.
- [62] Pan, MH & Ho, CT 2008, ‘Chemopreventive effects of natural dietary compounds on cancer development’, Chemical Society Reviews, vol. 37, no. 11, pp. 2558-74.
- [63] Dhar, DN 1981, Chemistry of chalcones and related compounds, Wiley, New York.
- [64] Elias, DW, Beazely, MA & Kandepu, NM 1999, ‘Bioactivities of chalcones’, Current Medicinal Chemistry, vol. 6, no. 12, pp. 1125-49.
- [65] Torigge, T, Arisawa, M, Itoh, S, Fujii, M & Maruyama, HB 1983, ‘Antimutagenic chalcones: antagonizing the mutagenicity of benzo (a) pyrene on salmonellatyphimurium’, Biochemical and Biophysical Research Communications, vol.112, no. 3, pp. 833-842.
- [66] Mager, L, Melzer, C, Barzoukas, M, Fort, A, Mery, S & Nicoud, JF 1997, ‘High net gain at 514 nm in a photorefractive polymer doped with a chalcone derivative’, Applied Physics Letters, vol. 71, no. 16, pp. 2248-2250.
- [67] Kumar, PR, Ravindrachary, V, Janardhana, K & Poojary, B 2012, ‘Structural and optical properties of a new chalcone single crystal’, Journal of Crystal Growth, vol. 354, no. 1, pp. 182-187.
- [68] Linke, HA & Eveleigh, DE 1975, ‘Dihydrochalcones and chalcones of some flavanone glycosides: qualitative and quantitative determinations’, Zeitschriftfur Naturforschung B, vol. 30, no. 11-12, pp. 940-942.
- [69] Jasinski, JP, Pek, AE, Narayana, B, Kamath, PK & Yathirajan, HS 2010, ‘(E)-1-(4Bromophenyl)-3-(2-methoxyphenyl) prop-2-en-1-one’, Acta Crystallographica Section E: Structure Reports Online, vol. 66, no. 8, pp. 1995-1996.

- [70] Ward, FE, Garling, DL, Buckler, RT, Lawler, DM & Cummings, DP 1981, ‘Antimicrobial 3-methylene flavanones’, *Journal of Medicinal Chemistry*, vol. 24, no. 9, pp. 1073-1077.
- [71] Frisch, MJ, Trucks, GW, Schlegel, HB, Scuseria, GE, Robb, MA, Cheeseman, JR, Scalmani, G, Barone, V, Mennucci, B, Petersson, GA, Nakatsuji, H, Caricato, M, Li, X, Hratchian, HP, Izmaylov, AF, Bloino, J, Zheng, G, Sonnenberg, JL, Hada, M, Ehara, M, Toyota, K, Fukuda, R, Hasegawa, J, Ishida, M, Nakajima, T, Honda, Y, Kitao, O, Nakai, H, Vreven, T, Montgomery, JA, Jr., Peralta, JE, Ogliaro, F, Bearpark, M, Heyd, JJ, Brothers, E, Kudin, KN, Staroverov, VN, Keith, T, Kobayashi, R, Normand, J, Raghavachari, K, Rendell, A, Burant, JC, Iyengar, SS, Tomasi, J, Cossi, M, Rega, N, Millam, JM, Klene, M, Knox, JE, Cross, JB, Bakken, V, Adamo, C, Jaramillo, J, Gomperts, R, Stratmann, RE, Yazyev, O, Austin, AJ, Cammi, R, Pomelli, C, Ochterski, JW, Martin, RL, Morokuma, K, Zakrzewski, VG, Voth, GA, Salvador, P, Dannenberg, JJ, Dapprich, S, Daniels, AD, Farkas, O, Foresman, JB, Ortiz, JV, Cioslowski, J & Fox, DJ 2010 Gaussian 09, Revision B.01, Gaussian, Inc., Wallingford CT.
- [72] Sundius, T 1990, ‘Molvib-A flexible program for force field calculations’, *Journal of Molecular Structure*, vol. 218, pp. 321-326.
- [73] Pulay, P, Fogarasi, G, Pongor, G, Boggs, JE & Vargha, A 1983, ‘Combination of theoretical ab initio and experimental information to obtain reliable harmonic force constants. Scaled quantum mechanical (QM) force fields for glyoxal, acrolein, butadiene, formaldehyde, and ethylene’, *Journal of the American Chemical Society*, vol. 105, no. 24, pp. 7037-47.
- [74] Kleinman, DA 1962, ‘Nonlinear dielectric polarization in optical media’, *Physical Review*, vol. 126, no. 6, pp. 1977.
- [75] Morris, GM, Goodsell, DS, Halliday, RS, Huey, R, Hart, WE, Belew, RK & Olson, AJ 1998, ‘Automated docking using a Lamarckian genetic algorithm and an empirical binding free energy function. Journal of Computational Chemistry’, vol. 19, no. 14, pp. 1639-1662.
- [76] Trott, O & Olson, AJ 2010, ‘Auto Dock Vina: improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading’, *Journal of Computational Chemistry*, vol. 31, no. 2, pp. 455-461.

- [77] Espinoza-Hicks, JC, Camacho-Dávila, AA, Flores-Holguín, NR, Nevárez-Moorillón, GV, Glossman-Mitnik, D & Rodríguez-Valdez, LM 2013, ‘Experimental and quantum chemical studies of a novel synthetic prenylated chalcone’, *Chemistry Central Journal*, vol. 7, no. 1, pp. 17-28.
- [78] Varsanyi, G, 1969, *Vibrational spectra of benzene derivatives*, Academic Press, New York.
- [79] Tommasini, M, Castiglioni, C, Del Zoppo, M & Zerbi, G 1999, ‘Relationship between infrared and Raman intensities in molecules with polarized π electrons’, *Journal of Molecular Structure*, vol. 480-481, pp. 179-188.
- [80] Smith, BC 1998, *Infrared spectral interpretation: a systematic approach*, CRC press. London.
- [81] Gussoni, M & Castiglioni, C 2000, ‘Infrared intensities. Use of the CH-stretching band intensity as a tool for evaluating the acidity of hydrogen atoms in hydrocarbons’, *Journal of Molecular Structure*, vol. 521, no. 1-3, pp. 1-8.
- [82] Palomar, J, De Paz, JL & Catalan, J 1999, ‘Vibrational study of intramolecular hydrogen bonding in o-hydroxybenzoyl compounds’, *Chemical Physics*, vol.246, no. 1-3, pp. 167-208.
- [83] Rumi, M & Zerbi, G 1999, ‘Conformational dependence of vibrational and molecular nonlinear optical properties in substituted benzenes: the role of π electron conjugation and back-donation’, *Journal of Molecular Structure*, vol. 509, no. 1-3, pp. 11-28.
- [84] Socrates, G 1990, *Infrared characteristic frequencies*, Wiley-Inter Science Publication, New York.
- [85] Allen, SD, Almond, MJ, Bruneel, JL, Gilbert, A, Hollins, P & Mascetti, J 2000, ‘The phototwo moleculesisation of trans-cinnamic acid and its derivatives: a study by vibrational microspectroscopy’, *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, vol. 56, no. 12, pp. 2423-2430.

- [86] Sanchez-Cortes, S & Garcia-Ramos, JV 1999, ‘Photoinduced coupling and adsorption of caffeic acid on silver surface studied by surface-enhanced Raman spectroscopy’, Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy, vol. 55, no. 14, pp. 2935-2941.
- [87] Castro, JL, Lopez Ramirez, MR, Lopez Tocon, I & Otero JC2002, ‘Surface enhanced Raman scattering of 3-phenylpropionic acid (hydrocinnamic acid)’, Journal of Raman Spectroscopy, vol. 33, no. 6, pp. 455-9.
- [88] Dimmock, JR, Elias, DW, Beazely, MA & Kandepu, NM 1999, ‘Bioactivities of chalcones’, Current Medicinal Chemistry, vol. 6, pp. 1125-1149.
- [89] Tomar, V, Bhattacharjee, G, Kamaluddin & Ashok Kumar 2007, ‘Synthesis and antimicrobial evaluation of new chalcones containing piperazine or 2,5vdichlorothiophene moiety’, Bioorganic Medicinal Chemistry Letter, vol. 17, no. 19, pp. 5321-5324.
- [90] Katib, S, Nerya, O, Musa, R, Shmuel, M, Tamir, S & Vaya, J 2005, ‘Chalcones as potent tyrosinaseinhibitors: the importance of a 2, 4-substituted resorcinol moiety’, Bioorganic Medicinal Chemistry, vol. 13, no. 2, pp. 433-441.
- [91] Avila, HP, Smania Ede, F, Monache, FD & Smania, A Jr 2008, ‘Structure-activity relationship of antibacterial chalcones’, Bioorganic Medicinal Chemistry, vol. 16, no. 22, pp. 9790-9794.
- [92] Basavaraju, YB & Devaraju, 2002, ‘1-(3, 4-Dimethoxyphenyl)-3-phenyl-prop-2-en1-one’, Indian Journal of Hetrocyclic Chemistry, vol. 11, no. 3, pp. 229-232.
- [93] Jasinski, JP, Butcher, RJ, MusthafaKhaleel, V, Sarojini, BK & Narayana, B 2011, ‘(2E)-3-(3-4-Dimethoxyphenyl)-1-(4-hydroxyphenyl) prop-2-en-1-

one’, Acta Crystallographica section E: Structure Reports Online, E67, pp. 813-814.

- [94] Glendening, ED, Reed, AE, Carpenter, JE & Weinhold, F 1998, ‘NBO Version 3.1’, TCI, University of Wisconsin, Madison.
- [95] Todd A. Keith 2016, AIMAll (Version 18.08.17), TK Gristmill Software, Overland Park KS, USA.
- [96] LU T & Chen, F 2012, ‘Multiwfn, a multifunctional wave function analyzer’, Journal of Computational Chemistry, vol. 33, no. 5, pp. 580-592.
- [97] Wolff, SK, Grimwodd, DJ, Mc Kinnon, JJ, Turner, MJ, Jayathilaka, D & Spackman, MA 2012, Crystal Explorer (Version 3.1) Univrsity of Western Australia.
- [98] Gatti, C, Saunders, V & Roetti, C 1994, ‘Crystal field effects on the topological properties of the electron density in molecular crystals: The case of urea’, Journal of Chemical Physics, vol. 101, no.12, pp. 10686-10696.
- [99] Grabowski, SJ 2000, ‘High-level ab initio calculations of di-hydrogen bonded complexes’, Journal of Physical Chemistry A, vol. 104, no. 23, pp. 5551-5557.
- [100] Espinosa, E, Molins, E & Lecomte, C 1998, ‘Hydrogen bond strengths revealed bytopological Analyses of experimentally observed electron densities’, Chemical Physics Letter, vol. 285, no. 3-4, pp. 170-173.
- [101] Socrates, G 1980, Infrared characteristic group of wave numbers, Wiley Interscience, New York.

- [102] Socrates, G 2001, Infrared and Raman characteristic group frequencies, table and charts, Wiley, Chichester.
- [103] Brain Smith, C 1999, Infrared spectral interpretation: a systematic approach, CRC press, USA.
- [104] Sajan, D, Hubert Joe, I, Jayakumar, VS & Zaleski, J 2006, ‘Structural and electronic contributions to hyperpolarizability in methyl p-hydroxy benzoate’, Journal of Molecular Structure, vol. 785, no. 1-3, pp. 43-53.
- [105] Roeges, NPG 1994, Guide to complete interpretation of IR spectra of organic compounds, Wiley, New York.
- [106] Clothup, NB, Daly, LH & Wiberly, SE 1990, Introduction to IR and Raman Spectroscopy, Academic Press, New York.
- [107] Troeberg, L, Chen, X, Flaherty, TM, Morty, RE, Cheng, M, Springer, HC, McKerrow, JH, Kenyon, GL, Lonsdale-Eccles, JD, Coetzer, THT & Cohen, FE 2000, ‘Chalcone, acylhydrazide, and related amides kill cultured Trypanosomabrucei’, Molecular Medicine, vol. 6, no. 8, pp. 660–669.
- [108] Ram, VJ, Saxena, AS, Srivastava, S & Chandra, S 2000, ‘Oxygenated chalcones and bischalcones as potential antimalarial agents’, Bioorganic Medical Chemistry Letter, vol. 10, no. 19, pp. 2159–2161.
- [109] Dimmock, JR, Elias, DW, Beazely, MA & Kandepu, NM 1999, ‘Bioactivities of chalcones’, Current Medical Chemistry, vol. 6, no. 12, pp. 1125–1149.
- [110] Rohl, AL, Moret, M, Kaminsky, W, Claborn, K, McKinnon, JJ & Kahr B 2008, ‘Hirshfeld surfaces identify inadequacies in computations of intermolecular interactions in crystals: pentamorphic 1, 8-dihydroxyanthraquinone’, Crystal Growth and Design, vol. 8, no. 12, pp. 4517-4525.

- [111] Luo, YH, Zhang, CG, Xu, B & Sun, BW 2012, ‘A cocrystal strategy for the precipitation of liquid 2,3-dimethyl pyrazine with hydroxyl substituted benzoic acid and a Hirshfeld surfaces analysis of them’, Crystal Engineering Communication, vol. 14, no.20, pp. 6860-6868.
- [112] Jasinski, JP, Butcher, RJ, Narayana, B, Veena, K & Yathirajan, HS 2010, ‘(2E)-1 (3-Chlorophenyl) 3-(4-chlorophenyl) prop-2-en-1-one’, Acta Crystallographica, Section E: Structure Reports Online, vol. 66, pp.1638-1639.
- [113] Karabacak M, Karagoz D & Kurt M 2009, ‘FT-IR, FT-Raman vibrational spectra and molecular structure investigation of 2-chloro-4-methylaniline: a combined experimental and theoretical study’, Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy, vol.72, no. 5, pp. 1076–1083.
- [114] Kurt, M, Yurdakul, M & Yurdakul, S 2004,‘Molecular structure and vibrational spectra of 3 chloro-4-methyl aniline by density functional theory and ab initio Hartree-Fock calculations’, Journal of Molecular Structure (Theochem.), vol. 711, no. 1-3, pp. 25-32.
- [115] Vein, DL, Colthup, NB, Fateley, WG & Grasseli JG 1991, The handbook of infrared and Raman characteristic frequencies of organic molecules, Academic Press, New York.
- [116] Sundaraganesan, N, Meganathan, C, Joshua, BD, Mani, P & Jayaprakash, A 2008, ‘FT IR, FT Raman spectra and ab initio HF, DFT vibrational analysis of 2, 3-difluoro phenol’, Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy, vol.68, no. 3, pp. 561-566.
- [117] Jacquemin, D, Preat, J & Perpete, EA 2005, ‘A TD-DFT study of the absorptionspectra of fast dye salts’, Chemical Physics Letter, vol. 410, no.4-6, pp. 254-259.

- [118] Soga, S, Shirai, H, Kobori, M & Hirayama, N 2008, ‘Use of amino acid composition to predict ligand-binding sites’, *Journal of Chemical Information and Modeling*, vol. 47, no. 2, pp. 400-406.
- [119] Shaik, B, Zafar, T & Agarwal, VK 2013, ‘Estimation of anti-HIV activity of HEPT analogues, using MLR, ANN, and SVM techniques’, *International Journal of Medicinal Chemistry*. vol. 17, pp. 1661–1677.
- [120] Lipinski, CA, Lombardo, F, Dominy, BW & Feene, PJ 2001, ‘Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings’, *Advanced Drug Delivery Reviews*, vol. 46, no.1-3, pp. 3-26.
- [121] Gramatica, P & Sangion, AA 2016, ‘Historical excursus on the statistical validation parameters for QSAR models: A clarification concerning metrics and terminology’, *Journal of Chemical Information and Modeling*, vol. 56, no. 6, pp. 1127-1131.
- [122] Prashar, H, Chawla, A, Sharma, A & Kharb, R, 2012, ‘Chalcone as versatile moiety for diverse Pharmacological activities’, *International Journal of Pharmaceutical science and Research*, vol. 3, no.7, pp.1913-1927.
- [123] Hilton, S, Naud, S, Caldwell, JJ, Boxall, K, Burns, S, Anderson, VE, Antoni, L, Allen, CE, Pearl, LH, Oliver, AW, Wynne, AG, Garrett, MD & Collins, I, 2010, ‘Identification and characterization of 2-aminopyridine inhibitors of checkpoint kinase 2’, *Journal of Medicinal Chemistry*, vol. 18, no.1, 707-718.
- [124] Ji, H, Delker, SL, Li, H, Martasek, P, Roman, LJ, Poulos, TL & Silverman, RB 2010, ‘Exploration of the active site of neuronal nitric oxide synthase by the design and synthesis of pyrrolidinomethyl 2-aminopyridine derivatives’, *Journal of Medicinal Chemistry*, vol. 53, no. 21, pp. 7804–7824.
- [125] Krayushkin, MM, Sedishev, IP, Yarovenko, VN, Zavarzin, IV, Kotovskaya, SK, Kozhenikov, DN, Charushin, VN & Russ 2008, ‘Synthesis of pyridines

from 1, 2, 4-triazines under high pressure’, Journal of Organic Chemistry, vol. 44, no. 3, 407–411.

- [126] Danagulyan, GG, Saakyan, LG & Panosyan, GA 2000, ‘Synthesis of fused systems containing an angular nitrogen atom from substituted 2-benzyl-4-hydrazine pyridimines’, Chemistry of Heterocyclic Compounds, vol.36, no.2, pp. 185-189.
- [127] Xiao-Dong, Y & Yu-Ye, Y 2008, ‘Synthesis, crystal structure and fungicidal activity of 1-(4-chlorophenyl)-2-(5-((3, 5-dimethyl-1H-pyrazol-1-yl) methyl)-4 phenyl-4H-1, 2, 4-triazol-3-ylthio) ethanone’, Structural Chemistry, vol. 19, no.4, pp. 693–696.
- [128] Jerry Jasinski, P, Ray Butcher, J, Samshuddin, S, Narayana, B & Yathiraja, HS 2011, ‘(2E)-3 (3-Bromo-4-methoxyphenyl)-1- (pyridin-2-yl) prop-2-en-one’, Acta Crystallographica section E: Structure Reports Online, vol. E67, pp. 352–353.
- [129] Kohn, W & Sham, LJ 1965, ‘Self-consistent equations including exchange and correlation effects’, Physical Review, vol. 140, no. 4A, pp. 1133–1138.
- [130] Parr, RG & Yang, W 1989, Density-functional theory of atoms and molecules, Oxford University Press, Oxford
- [131] Guo, D & Goodman, L 1996 ‘Nature of barrier forces in acetaldehyde. Nature of barrier forces in acetaldehyde’, Journal of Physical Chemistry, vol. 100, no.30, pp. 12540–12545.
- [132] Alver, O, Parlak, C & Senyel, M 2007, ‘FT-IR and NMR investigation of 1 phenylpiperazine: A combined experimental and theoretical study’, Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy, vol. 67A, no. 3-4, pp. 793-801.

- [133] Scheibye, S, El-Barbary, AA, Lawesson, SO, Fritz, H & Rihs, G 1982, 'Studies on organophosphorus compounds—XLI: Formation of 3-pyrazoline-5-thione disulfides from 3, 5-pyrazolidinediones. C-alkylation of 3, 5 pyrazolidinediones', *Tetrahedron*, vol. 38, no. 24, pp. 3753–3760.
- [134] EI-Shahawy, AS, Ahmed, SM & Sayed, NK 2007, 'INDO/SCF-CI calculations and structural spectroscopic studies of some complexes of 4 hydroxyacetanilide', *SpectrochimicaActa Part A: Molecular and Biomolecular Spectroscopy*, vol. 66, no. 1, pp. 143–152.
- [135] Jones, RA & Moritz, AG, 1965, 'Pyrrole studies—VI, the N...H stretching frequencies of substituted pyrroles: methyl and carbethoxy substituents', *SpectrochimicaActa Part A: Molecular and Biomolecular Spectroscopy*, vol. 21, no. 2, pp. 295–299.
- [136] Gyomore, A, Kovacs, Z, Nagy, T, Kudar, V, Szabo, A & Csampai, A 2008, 'DNMR, DFT and preparative study on the conformation of (Z)-4, 5, 6, 7 tetrahydropyrazolo [1, 5-e] benzo [g][1, 5] diazonin-8-ones and (Z)-4, 5 dihydropyrazolo [1, 5-d] ...,' *Tetrahedron*, vol. 64, no. 48, pp.10837–10848.
- [137] Huang, MH, Wu, YY, Feick, H, Tran, N, Weber, E & Yang, PD 2001, 'Catalytic growth of zinc oxide nanowires by vapor transport', *Advanced Materials Communications*, vol.13, no. 2, pp. 113-116.
- [138] Tauc, J & Abeles, F (Ed.) 1970, *Optical properties of solids*, Amsterdam North Holland.
- [139] Kurtaran, R, Odabasioglu, S, Azizoglu, A, Kara, H & Atakol, O 2007, 'Experimental and computational study on [2, 6-bis (3, 5-dimethyl-N-pyrazolyl) pyridine]- (dithiocyanato) mercury (II)', *Polyhedron*, vol. 26, no. 17, pp. 5069–5074.
- [140] Di Carlo, G, Mascolo, N, Izzo, AA & Francesco, C 1999, 'Flavonoids: old and new aspects of a class of natural therapeutic drugs', *Journal of Life Sciences*, vol. 65, no. 4, pp. 337-353.

- [141] Ferlay, J, Shin, HR & Bray, F 2008, 'International Agency for Research on Cancer; Lyon, France: 2010. GLOBOCAN 2008, Cancer Incidence and Mortality Worldwide: IARC Cancer Base No. 10.
- [142] Kim, YS, Kumar, V, Lee, S, Iwai, A, Neckers, L, Malhotra, SV & Trepel, JB 2012, 'Methoxychalcone inhibitors of androgen receptor translocation and function, Bioorganic and Medicinal Chemistry Letters, vol. , no. 2, pp. 2105-2109.
- [143] Yathirajan, HS, Mayekar, AN, Narayana, B, Sarojini, BK& Michael, B 2007, '(2E)-1-(2,4-Dichlorophenyl)-3-(4,5-dimethoxy-2-nitrophenyl)-2-propen-1-one', Acta Crystallographica Section E, vol. 63, no. 1, 2200-2201.
- [144] Kevin, E, Bryan, T & Kenneth, M 2007, 'Critical assessment of the performance of density functional methods for several atomic and molecular properties', Journal of Chemical Theory Computation, vol.3, no.2 , pp. 407-433
- [145] Alan Reed, E, Larry Curtiss, A& Frank, W 1988, 'Intermolecular interactions from a natural bond orbital, donor-acceptor view point', Journal of Chemical Review, vol.88,no.6, pp. 899-926.
- [146] Chandra, S, Saleem, H, Sundaraganesan, N & Sebastian, S 2009, 'The spectroscopic FT-IR gas phase, FT-IR, FT-Raman, polarizabilities analysis of Naphthoic acid by density functional methods', Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy, vol. 74, no. 3, pp. 704-713.