

LIST OF TABLES

TABLE NO	TITLE	PAGE NO
4.1	Optimized bond length bond angle of BRMP by B3LYP/6-31G** in comparison with the XRD data	57
4.2	Optimized dihedral angle of BRMP by B3LYP/6-31G** in comparison with the XRD data	58
4.3	Second order perturbation theory analysis of fock matrix in Natural bond orbital basis	61
4.4	Natural Population Analysis	63
4.5	Definition of internal coordinates of 1-4-Bromophenyl -3-(2-methoxyphenyl) prop-2-en-1-one	67
4.6	Definition of local symmetry coordinates (much like the natural symmetry coordinates) and the corresponding force constant (mdyne /Å) with scale factors used.	69
4.7	Calculated Vibrational wave numbers, observed IR and Raman frequencies and their assignments with Potential Energy Distribution% for 1-4-Bromophenyl -3-(2-methoxyphenyl) prop-2-en-1-one.	71
4.8	The dipole moments components μ (D), static polarizability components α (a. u), the average polarizability α_0 (esu), the anisotropy of the polarizability $\Delta\alpha$ (esu), and the first hyper polarizability components β (a. u), the first hyper polarizability β_0 ($\times 10^{-30}$ esu) of BRMP1-4-Bromophenyl -3-(2-methoxyphenyl) prop-2-en-1-one.	75
4.9	Bond distances and type between receptor and (E)-1-(4-Bromophenyl)-3-(2-methoxyphenyl) prop-2-en-1-one (ligand)	81
5.1	Optimized geometrical parameters of DIM monomer and dimer by B3LYP/6-31G** in comparison with XRD data	88
5.2	Second order perturbation theory analysis of Fock matrix in	91

	NBO basis	
5.3	Topological parameters for bonds of interacting atoms of one molecule and two molecules: electron density $\rho(r)$, Laplacian of electron density $\nabla^2 \rho(r)$, electron kinetic energy density (G), electron potential energy density (V), total electron energy density (H) at bond critical point	95
5.4	Definition of internal coordinates of DIM	100
5.5	Definition of local symmetry coordinates (much like the natural symmetry coordinates) and the corresponding force constant (mdyne /Å) with scale factors used.	102
5.6	Calculated vibrational wave numbers, observed IR and Raman frequencies, IR and Raman intensities and their assignments with PED% for DIM	104
6.1	Optimized geometrical parameters of CCL in comparison with X-Ray diffraction data	115
6.2	Optimized geometrical parameters of CCL in comparison with X-Ray diffraction data	116
6.3	Donor-acceptor interactions results of (2E)-1-(3-Chlorophenyl)-3-(4 Chlorophenyl) prop-2-en-1-one: followed by Fock matrix in Natural bond orbital basis	119
6.4	Definition of internal coordinate of (2E)-1-(3-Chlorophenyl)-3-(4Chlorophenyl) prop-2-en-1-one	122
6.5	Definition of local symmetry coordinates (much like the natural symmetry coordinates) and the corresponding force constant (mdyne /Å) with scale factors used.	124
6.6	Calculated vibrational wavenumbers, observed IR and Raman frequencies, IR and Raman intensities and their assignments with PED% for (2E)-1-(3-Chlorophenyl)-3-(4-Chlorophenyl) prop-2-en-1-one	125
6.7	Comparison of experimental and calculated absorption wavelength (λ , nm), excitation energies (E, ev) and oscillator strength (f) of (2E)-1-(3 Chlorophenyl)-3-(4-	131

	Chlorophenyl) prop-2-en-1-one	
6.8	Bond distances and type between androgen receptor and (2E)-1-(3-Chlorophenyl)-3-(4-chlorophenyl) prop-2-en-1-one (ligand)	135
7.1	Optimized geometrical parameters of PYR in comparison with the XRD data	141
7.2	Donor-acceptor interactions results of PYR followed by Fock matrix in NBO basis	143
7.3	Calculated vibrational wavenumbers, observed IR and Raman frequencies, IR and Raman intensities and their assignments with PED% for PYR	148
7.4	Experimental and calculated absorption wavelengths, energies and oscillator strengths of PYR using the TD-DFT method at CAM B3LYP level	153
8.1	Optimized bond length bond angle of DINO by B3LYP/6-31G** in comparison with the XRD data	161
8.2	Optimized dihedral angle of DINO by B3LYP/6-31 G** in comparison with the XRD data.	162
8.3	Second order perturbation theory analysis of fock matrix in NBO basis.	165
8.4	Calculated vibrational wavenumbers, observed IR and Raman frequencies, IR and Raman intensities and their assignments with PED% for DINO	168
8.5	The bond distances and type between receptor and (2E)-1-(2,4-Dichlorophenyl)-3-(4,5-dimethoxy-2-nitrophenyl)-2-propen-1-one (Ligand)	177

LIST OF FIGURES

FIGURE NO	TITLE	PAGE NO
3.1	Schematic diagram of FT-IR Spectrometer	17
3.2	Schematic representation of Raman lines	21
3.3	Depolarization measurement	25
3.4	Schematic of a Raman spectrophotometer	26
3.5	Schematic diagram of UV-Visible spectrometer	30
4.1	Optimized structure of 1-4-Bromophenyl-3-(2-methoxyphenyl) prop-2-en-1-one calculated at Becke three Lee-Yang-Parr /6-31G** level of theory	56
4.2	Simulated (C, A) and experimental (D, B) Fourier Transform Infrared and Raman spectra of 1-4-Bromophenyl-3-(2-methoxyphenyl) prop-2-en1-one at Becke three Lee-Yang-Parr /6-31G** basis set.	64
4.3	Molecular Hirshfeld surfaces (dnorm, curvedness, di, de and shape index) of 1-4-Bromophenyl -3-(2-methoxyphenyl) prop-2-en-1-one	77
4.4	2-Dimensional fingerprint plot of 1-4-Bromophenyl -3-(2 methoxyphenyl) prop-2-en-1-one	78
4.5	Interactive part of ligand and androgen receptor	80
4.6	Docked protocol reproduces the co-crystallized conformation with the Hydrogen bond (green), π -alkyl (pink), π - π (magenta), and Hydrogen bond receptor surface shown	80
4.7	Schematic for the docked conformation of active site of the title compound at receptor	81
5.1	Molecular structure of DIM one molecule calculated at B3LYP/6-31G**level of theory	87
5.2	Molecular structure of DIM two molecule calculated at B3LYP/6-31G**level of theory	87

5.3	Molecular graph of DIM dimer of different bond path at critical points at (3,-1)	92
5.4	Simulated (A, C) and experimental (B, D) Fourier Transform Infrared and Raman spectra of DIM calculated at B3LYP/6-31G**level of theory	97
5.5	2-Dimensional fingerprint plot of DIM	109
6.1	Optimized structure of (2E)-1-(3-Chlorophenyl)-3-(4-chlorophenyl) prop 2-en-1-one by Becke three Lee-Yang-Parr /6-31G** in comparison with X-Ray diffraction data.	114
6.2	Simulated (C, A) and experimental (D, B) Fourier Transform Infrared and Raman spectra of (2E)-1-(3-Chlorophenyl)-3-(4-chlorophenyl) prop-2-en-1-one at Becke three Lee-Yang-Parr /6-31G** basis set.	120
6.3	Tauc plot of (2E)-1-(3-Chlorophenyl)-3-(4-chlorophenyl) prop-2- en-1-one.	130
6.4	Molecular Hirshfeld surfaces (dnorm, back, de, di, shape index and curvedness) of (2E)-1-(3-Chlorophenyl)-3-(4-chlorophenyl) prop-2-en-1-one. The different colour indicates the distance shorter or equal or longer than Van der walls radii.	133
6.5	2-Dimensional fingerprint plot of (2E)-1-(3-Chlorophenyl)-3-(4-chlorophenyl) prop-2- en- 1one. 2-Dimensional plot highlight atom pair close contact.	133
6.6	Comprehensive perception of receptor and (2E)-1-(3-Chlorophenyl)-3-(4-chlorophenyl) prop-2-en-1-one (ligand) interaction after docking. The ligand binds to receptor via hydrogen and hydrophobic bonds	134
6.7	Docked protocol reproduces the conformation of receptor and ligand the Hydrogen bond (green), π -alkyl (pink), and π - π (magenta) are shown in dots, and the receptor was shown as Hydrogen bond acceptor donor	135

surface. The ligand binds to receptor via hydrogen and hydrophobic bonds.

7.1	Optimized structure of PYR	140
7.2	Simulated (C, A) and experimental (D, B) Fourier Transform Infrared and Raman spectra of (2E)-3-(3-Bromo-4methoxyphenyl)-1-(pyridin-2-yl) prop-2-en1-one at Becke three Lee-Yang-Parr /6-31G** basis set.	145
7.3	Taue plot of PYR	155
7.4	Homo-Lumo Plot of PYR	156
8.1	Optimized structure of DINO	160
8.2	A-Simulated FT-IR Spectrum, B-Experimental FT-IR spectrum	172
8.3	A-Simulated Raman Spectrum, B-Experimental Raman spectrum	172
8.4	Hirshfeld surface showing dmorm, di,de,shape index and curvedness	174
8.5	2-D Finger print plot showing various interactions.	174
8.6 a	Ligand is docked in active site of androgen receptor. Secondary structure of androgen receptor represented by ribbon model, and ligand is represented by ball and stick model	175
8.6 b	Secondary structure of androgen receptor represented by surface model, and ligand is represented by ball and stick (blue colour) model.	176
8.6 c	Interactions of ligand with androgen receptor amino acids omitting androgen receptor, ligand surrounding amino acids are in three letters code represented in dark blue.	176