

LIST OF TABLES

Table No	Title	Page No
3.1	Optimized geometrical parameters of anilazine(compound1) and N-(4,6-dichloro-1,3,5-triazine-2-yl) aniline (compound2)at B3LYP/6-311G(d,p) level	44
3.2	Second order perturbation theory analysis	47
3.3	Detailed assignment of fundamental vibrations of anilazine (compound1) by normal coordinate analysis based on SQM force field calculations	48
3.4	Natural charges of anilazine (compound1) and N-(4,6-dichloro-1,3,5-triazine-2-yl) aniline (compound2)	55
3.5	Potential energy surface scan related data	57
3.6	Calculated and experimental electronic absorption spectrum of anilazine with different solvents (acetone, toluene and gas) at PCM-TDDFT basis set.	60
3.7	Calculated energy values of anilazine and N-(4,6-dichloro-1,3,5-triazine-2-yl) aniline by B3LYP/6-311G(d,p) basis set	63
3.8	The experimental and calculated ¹ H and ¹³ C isotropic chemical shifts with respect to Tetra methyl Silane(TMS) of anilazine	64
3.9	Summary of hydrogen bonding of protein target with anilazine (compound1) and N-(4,6-dichloro-1,3,5-triazine-2-yl) aniline (compound2)	69
3.10	Molecular docking results of anilazine (compound1) and N-(4,6-dichloro-1,3,5-triazine-2-yl) aniline(compound2) with different protein targets	70
3.11	Antifungal activity of 1ml DMSO extracts of anilazine (compound1)	72
4.1	Optimized geometrical parameters of benalaxyl and its related compounds at B3LYP/6-311G(d,p) level	77
4.2	Natural charges of benalaxyl and its related compounds	87
4.3	Possible hydrogen bonding of benalaxyl and it's related compounds	90
4.4	Potential energy surface scan analysis of benalaxyl	91
4.5	The experimental and calculated ¹ H and ¹³ C isotropic chemical shifts with respect to Tetra methyl Silane(TMS) of benalaxyl	95

Table No	Title	Page No
4.6	Molecular docking results of benalaxyl, metalaxyl and furalaxyl with different protein targets	99
4.7	Summary of hydrogen bonding of protein target with benalaxyl, metalaxyl and furalaxyl	100
4.8	Inhibition of the fungicidal growth in benalaxyl performed in biological replicate	101
5.1	Optimized geometrical parameters of benomyl and it's related compounds at B3LYP/6-311G(d,p) level	108
5.2	Natural charges of benomyl and its related compounds	115
5.3	Possible hydrogen bonding of benomyl and it's related compounds	117
5.4	Potential energy surface scan analysis of benomyl	118
5.5	The experimental and calculated ¹ H and ¹³ C isotropic chemical shifts with respect to Tetra methyl Silane(TMS) of benomyl	125
5.6	Molecular docking results of benomyl and its related compounds with different protein targets	128
5.7	Summary of hydrogen bonding of protein target with benomyl and its related compounds	129
5.8	Antifungal activity of 1ml DMSO extracts of benomyl	131
6.1	Structural parameters (bond length in Å and angles in degree) of H-bond in Clotrimazole-H ₂ O complexes calculated at the B3LYP/6-311G(d,p) level	144
6.2	Detailed assignment of fundamental vibrations of clotrimazole by normal coordinate analysis based on SQM force field calculations	146
6.3	The X-H stretching vibrational frequencies (strength) of H-bonds in both clotrimazole-H ₂ O complexes and monomers	159
6.4	Natural charges of clotrimazole and its related compounds	160
6.5	Second order perturbation theory analysis of clotrimazole and it's related compounds	162
6.6	The second order perturbation energies E ⁽²⁾ (kJ/mol) of H-bonds in clotrimazole-H ₂ O complexes obtained by NBO analysis	164
6.7	The experimental and calculated ¹ H and ¹³ C isotropic chemical shifts with respect to Tetra methyl Silane(TMS) of clotrimazole	170

Table No	Title	Page No
6.8	Binding affinity, Hydrogen bond interaction and number of hydrogen bond residues of compound1, related compounds and reference molecule (Fluconazole) with the target protein	177
6.9	Antifungal activity of 1ml DMSO extracts of clotrimazole	179
7.1	Optimized geometrical parameters of flutolanil and mepronil at B3LYP/ 6-311G(d,p) level	188
7.2	The observed and calculated bond length and stretching frequency of modes involved in intermolecular hydrogen bonding	198
7.3	Natural charges of flutolanil and mepronil (monomer and dimer model)	203
7.4	Possible hydrogen bonding of flutolanil and mepronil (monomer and dimer model)	205
7.5	The Docking score, Hydrogen bond interaction and the interacting residues of different poses of the compounds with the target protein 3IS3	221
7.6	Antifungal activity of 1ml DMSO extracts of flutolanil and mepronil	222

LIST OF FIGURES

Figure No	Title	Page No
1.1	Electromagnetic spectrum	8
1.2	Schematic diagram of dispersive infrared spectrophotometer	11
1.3	Schematic diagram of Fourier Transform Infrared spectrophotometer	13
1.4	Rayleigh, stoke's and anti stoke's scattering energy levels	16
1.5	The block diagram of FT-Raman spectrophotometer	20
1.6	Schematic diagram of UV-Visible double beam spectrophotometer	21
1.7	Schematic diagram of Nuclear magnetic resonance spectrophotometer	25
3.1	The optimized molecular structure of anilazine and N-(4,6-dichloro-1,3,5-triazine-2-yl)aniline	44
3.2	Combined FT-IR spectrum of anilazine	53
3.3	Combined FT-Raman spectrum of anilazine	54
3.4	Potential energy surface scan of anilazine	58
3.5	UV-visible spectrum of anilazine (acetone solvent)	59
3.6	HOMO-LUMO energy gap of anilazine and N-(4,6-dichloro-1,3,5-triazine-2-yl)aniline	61
3.7	TDOS spectrum of anilazine	62
3.8	¹ H NMR spectrum of anilazine	65
3.9	¹³ C NMR spectrum of anilazine	66
3.10	Hirshfeld surface mapped with d _c , d _i , d _{norm} , shape index and curvedness of anilazine	67
3.11	Finger print plots of anilazine	68
3.12	The hydrogen bond distances in the docking studies of anilazine with target proteins	70
3.13	Antifungal activity of anilazine at different concentrations for two fungal pathogens (<i>Aspergillus niger</i> and <i>Candida albicans</i>)	72
4.1	Optimized molecular structure of benalaxyl, metalaxyl and furalaxyl	77
4.2	Combined FT-IR spectrum of benalaxyl	86

Figure No	Title	Page No
4.3	Combined FT-Raman spectrum of benalaxyl	86
4.4	Potential energy surface scan of benalaxyl	92
4.5	UV-visible spectrum of benalaxyl (water solvent)	93
4.6	HOMO-LUMO plot of benalaxyl	94
4.7	¹ H NMR spectrum of benalaxyl	97
4.8	¹³ C NMR spectrum of benalaxyl	98
4.9	The hydrogen bond distances in the docking studies of benalaxyl with target protein, PDB ID: 3COZ	99
4.10	Antifungal activity of benalaxyl performed at biological replicate for two fungal pathogens (<i>Aspergillus niger</i> and <i>Candida albicans</i>)	102
5.1	Optimized molecular structure of (a) benomyl (compound1), (b) {Ethyl N-[1-(piperidin-1-ylmethyl)benzimidazol-2-yl]carbamate} (compound2), (c) {EthylN-[1-(pyrrolidin-1-ylmethyl)benzimidazol-2yl]carbamate} (compound3)	107
5.2	Combined FT-IR spectrum of benomyl	114
5.3	Combined FT-Raman spectrum of benomyl	114
5.4	Potential energy surface scan of benomyl	119
5.5	The experimental UV-Visible spectrum of benomyl (methanol solvent)	120
5.6	HOMO-LUMO plot of benomyl and it's related compounds	121
5.7	TDOS spectrum of benomyl	122
5.8	Molecular electrostatic potential of benomyl	123
5.9	¹ H NMR spectrum of benomyl	127
5.10	¹³ C NMR spectrum of benomyl	127
5.11	The highest binding energy for docking studies of benomyl with target protein (PDB ID: 2BYK)	130
5.12	Antifungal activity of benomyl at different concentrations for two fungal pathogens (<i>Aspergillus niger</i> and <i>Candida albicans</i>)	131
5.13	RMSD of benomyl with different target proteins	133
5.14	RMSF of benomyl with different target proteins	134

Figure No	Title	Page No
6.1	Optimized molecular structure of (i) Clotrimazole (compound1), (ii) 1-trityl-1H-imidazole (compound2), (iii) 1-(diphenyl-(2-(trifluoromethyl)phenyl)methyl)imidazole (compound3), (iv) 1-[(2-methylphenyl)-diphenyl methyl]imidazole (compound4) and (v) clotrimazole-H ₂ O complex	141
6.2	Combined FT-IR spectrum of clotrimazole	157
6.3	Combined FT-Raman spectrum of clotrimazole	158
6.4	UV-visible spectrum of clotrimazole (acetone and chloroform solvents)	166
6.5	HOMO-LUMO plot of clotrimazole and its related compounds	167
6.6	TDOS spectrum of clotrimazole	168
6.7	Molecular electrostatic potential of clotrimazole	169
6.8	¹ H NMR spectrum of clotrimazole	172
6.9	¹³ C NMR spectrum of clotrimazole	173
6.10	d _{norm} , d _e , d _i , shape index and curvedness of clotrimazole	174
6.11	Fingerprint plots of clotrimazole	175
6.12	The hydrogen bond interaction of (a) clotrimazole (compound1), (b) compound2, (c) compound3, (d) compound4 (e) Fluconazole (reference molecule) with the protein Lanosterol 14 α -demethylase	178
6.13	Antifungal activity of clotrimazole at different concentrations for two fungal pathogens (<i>Aspergillus niger</i> and <i>Candida albicans</i>)	179
6.14	Time dependencies of the RMSDs	180
6.15	RMSF of ligands with protein	181
6.16	Radius of gyration of ligands with protein	181
7.1	Optimized molecular structure of (a) Flutolanil, (b) Mepronil (monomer form)	186
7.2	Optimized molecular structure of (a) Flutolanil, (b) Mepronil (dimer form)	187
7.3	Combined FT-IR spectrum of (a) Flutolanil (b) Mepronil	200
7.4	Combined FT-Raman spectrum of (a) Flutolanil (b) Mepronil	201
7.5	Potential energy surface scan of ϕ_1 (N ₁₇ -C ₁₅ -C ₁₄ -C ₅)	210
7.6	UV-visible spectrum of (a) Flutolanil, (b) Mepronil (solvent-acetone)	212

Figure No	Title	Page No
7.7	HOMO-LUMO energy gap of (a) Flutolanil, (b) Mepronil	213
7.8	MESP plot of (a) Flutolanil, (b) Mepronil	215
7.9	¹ H spectrum of (a) Flutolanil, (b) Mepronil	217
7.10	¹³ C spectrum of (a) Flutolanil, (b) Mepronil	218
7.11	Docked pose of (a) Mepronil with 3IS3 (b) Flutolanil with 3IS3	221
7.12	Antifungal activity of flutolanil and mepronil at different concentrations for two fungal pathogens (<i>Aspergillus niger</i> and <i>Candida albicans</i>)	222
7.13	Root Mean Square Deviation representation for the protein ligand (i) Mepronil complex (ii) Flutolanil complex	224
7.14	RMSF representation for (i) 17beta-Hydroxysteroid dehydrogenase – Mepronil complex, (ii) 17beta-Hydroxysteroid dehydrogenase – flutolanil complex	225
7.15	Radius of Gyration for (i) 17beta-Hydroxysteroid dehydrogenase – Mepronil complex, (ii) 17beta-Hydroxysteroid dehydrogenase – flutolanil complex	226