

# REFERENCES

## REFERENCES

- Anderson, E.C., Williamson E.G, Thompson E.A (2000)** Monte Carlo evaluation of the likelihood for  $N_e$  from temporally spaced samples. *Genetics*, 156: 2109-2118.
- Arden, W.R& Kapuscinski, A.R (2003)** Demographic and genetic estimates of effective population size ( $N_e$ ) reveals genetic compensation in steelhead trout. *Molecular Ecology*, 12: 35-49.
- Avise, J.C. (1994)** *Molecular Markers Natural History and Evolution*.Chapman and Hall, Inc., New York
- Bagley, M.J. & Gall, G.A.E. (1998)** Mitochondrial and nuclear DNA sequence variability among populations of rainbow trout (*Oncorhynchus mykiss*). *Molecular Ecology*, 7:945-961.
- Bailey, K.M., Stabeno P. J., Powers, D.A (1997)** The role of larval retention and transport features in mortality and potential gene flow of walleye pollock. *Journal of Fish Biology* 51, (Supplement ).
- Balloux, F. & Lugon Moulin, N. (2002)** The estimation of population differentiation with microsatellite markers. *Mol.Ecol*, 11: 155-159.
- Barber, P.H., Palumbi. S.R., Erdmann, MV. Moosa, M.K (2002)** Sharp genetic breaks among populations of *Haptosquilla pulchella* (Stomatopoda) indicate limits to larval transport: patterns, causes, and consequences. *Molecular Ecology*, 11, 659–674.
- Baums, IB., Paris. CB., Cherubin, L.M. (2006)** A bio-oceanographic filter to larval dispersal in a reef-building coral. *Limnology and Oceanography*, 51, 1969–1981.

**Belshaw, R., M. Fitton., E. Herniou., C.Gimeno., and D.J. Quicke. (1998)** A phylogenetic reconstruction of the *Ichneumonoidea* (Hymenoptera) based on the D2 variable region of 28S ribosomal RNA. *Systematic Entomology*, 23: 109-123.

**Benzie, J.A.H., Kenway, M., Troff, L. (1997)** Estimates for the heritability of size in juvenile *P.monodon* from half-sib matings. *Aquaculture*, **152**: 49-53.

**Berry, B. F.(1974)** A revision of the *Panulirus homarus* group of spiny lobsters (Decapoda, Palinuridae). *Crustaceana* 27(1): 31-42.

**Bielawski, J.P& Pumo, D.E. (1997)** Randomly amplified polymorphic DNA (RAPD) analysis of Atlantic Coast striped bass. *Heredity*, **78**: 32–40

**Birky, C.W., Fuerst. P., Maruyama, T. (1989)** Organelle gene diversity under migration, mutation, and drift - equilibrium expectations, approach to equilibrium, effects of heteroplasmic cells, and comparison to nuclear gene. *Genetics*, 121: 613-627.

**Booth JD, Phillips BF (1994)** Early life history of spiny lobster. *Crustacea* 66:271–294

**Borowsky, L., McClelland. M, Cheng. R. &Welsh J. (1995)** Arbitrarily primed DNA fingerprinting for phylogenetic reconstruction in vertebrates: the *Xiphophorous* model. *Molecular Biology and Evolution*, **12**: 1022-1032.

**Bouchon, D., C. Souty-Grosset., & R. Raimond. (1994)** Mitochondrial DNA variation and markers of species identity in two penaeid shrimp species. *Aquaculture*, 127: 131-144.

**Boudry, P., Collet, B. Cornette., F. Hervouet., V. Bonhomme, F. (2002)** High variance in reproductive success of the Pacific oyster (*Crassostrea gigas*, Thunberg) revealed by microsatellite-based parentage analysis of multifactorial crosses. *Aquaculture*, **204**: 283-296.

**Brown, W. M., M. Jr. George., and A. C. Wilson. (1979)** Rapid evolution of animal mitochondrial DNA. *Proceedings of the National Academy of Sciences of the United States of America*, 76: 1967-1971.

**Brown, B. & Epifanio, J. (2003)** Nuclear DNA. E. M. Hallerman (Ed.), *Population genetics: principles and applications for fisheries scientists*. American Fisheries Society, Bethesda, Maryland: 101-123.

**Brower, A. V. Z., R. Desalle., and A. Vogler. (1996)** Gene trees, species trees, and systematics: a cladistic perspective. *Annual Review of Ecology and Systematics*, 27: 423-450.

**Buonaccorsi, V. P., McDowell, J. R., Graves, J. E. (2001)** Reconciling patterns of inter-ocean molecular variance from four classes of molecular markers in blue marlin (*Makaira nigricans*). *Molecular Ecology* 10: 1179–1196.

**Burggraf, S. G., K.O. Stetter, and C.R. Woese. (1992)** A phylogenetic analysis of *Aquifex pyrophilus*. *Systematic and Applied Microbiology* 15: 352-356.

**Caccone, A., Allegrucci, G., Fortunato, C and Sbordoni V. (1997)** Genetic differentiation within the European Sea Bass (*D. labryxo* as revealed by RAPD PCR assays. *J.Herd.* 88: 316-324.

**Chang, C. H., S. C. Chuang., Y. R. Chen, and J. H. Chen. (2005)** NADH Dehydrogenase Subunit 1 gene of the earthworm *Amyntas gracilis* (Kinberg,1867) (Oligochaeta: Megascolecidae), with the discussion on inferring the megascolecid phylogeny using DNA sequences. *Taiwania*, 50(2): 71-79.

**Chavarria, G., and J. M. Carpenter. (1995)** ‘‘Total evidence’’ and the evolution of highly social bees. *Cladistics*, 10: 229-258.

**Chhapgar, B. F., Deshmukh, S. K. (1971)** Lobster fishery of Maharashtra. *Journal of the Indian Fisheries Association* 1(1): 74–86.

**Chen, R. T., C. F. Tsai, and W. N. Tzeng. (2009)** Freshwater prawns (*Macrobrachium Bate*, 1868) of Taiwan with special references to their biogeographic origins. *Journal of Crustacean Biology*, 29: 232-244.

**Ciftci, Y. and Okumus, I. (2002)** Fish Population Genetics and Application of molecular markers to Fisheries and aquaculture- 1: basic Principles of Fish Population Genetics. *Turkish J. Fisheries and Aquatic Sciences*, 2: 145-155.

**Conant, G. C & P. O. Lewis. (2001)** Effects of nucleotide composition bias on the success of the parsimony criterion in phylogenetic inference. *Molecular Biology and Evolution*, 18: 1024-1033.

**Corely-Smith, G.E., Lim, C.J. & Brandhorst, B.P. (1996)** Production of androgenic zebrafish (*Danio rerio*). *Genetics*, 142: 265-276.

**Crandall, K. A., J. W. Fetzner, Jr., S. H. Lawler., M. Kinnersley, and C. M. Austin. (1999)** Phylogenetic relationships among the Australian and New Zealand genera of freshwater crayfishes. *Australian Journal of Zoology*, 47: 199-214.

**Crandall, K. A., O. R. P. Binindamonds., G. M. Mace, and R. K. Wayne. (2000)** Considering evolutionary processes in conservation biology. *Trends in Ecology & Evolution*, 15:290-295.

**Crochet, P.A (2000)** Genetic structure of avian populations - allozymes revisited. *Molecular Ecology* , 9: 1463-1469.

**Crossland, S., Coates, D., Grahame, J., & Mil, P.J. (1993)** Use of random amplified polymorphic DNAs (RAPDs) in separating two sibling species of *Littorina*. *Marine ecology progress series*, 6: 301-305.

**Crozier, W. W., (1994)** Maintenance of genetic variation in hatchery stocks of Atlantic salmon, (*Salmo salar L*): experience from the River Bush, Northern Ireland. *Aquacult. Fish. Manage*, **25**: 383–392.

**Crow, J.F & Denniston, C (1988)** Inbreeding and variance effective population numbers. *Evolution*, 42: 482-495.

**Cunningham, C.O & Mo, T.A. (1997)** Random amplified polymorphic DNA (RAPD) analysis of three Norwegian *Gyrodactylus* salar populations (*Monogenea: Gyrodactylidae*). *J. Parasitol*, **83**: 311-314

**Daan, N (1991)** Bias in virtual population analysis when the unit stock assessed consists of sub-stocks. ICES C.M.1991/D:17.

**Daniels, S. R., M. Hamer, & C. Rogers. (2004)** Molecular evidence suggests an ancient radiation for the fairy shrimp genus *Streptocephalus* (Branchiopoda: Anostraca). *Biological Journal of the Linnean Society*, 82: 313-3.

**De Jong Y., Van Der Wurff AG, Stam WT, Olsen JL (1998)** Studies on Dasyaceae: 3. Towards a phylogeny of the Dasyacean (*Ceramillaes, Rhodophyta*), based on comparative rbcL gene sequences and morphology. *European Journal of Phycology*, 33: 187-201

**Dinesh, K.R., Lim, T.M., Chua, K.L., Chan, W.K. and Phang, V.P. (1993)** RAPD analysis: an efficient method of DNA fingerprinting in fishes. *Zool. Sci*, 10: 849-854.

**Diniz, F.M., Maclean, N., Ogawa, M., Cintra, I.H.A. & Bentzen, P. (2005)** The hypervariable domain of the mitochondrial control region in Atlantic spiny lobsters and its potential as a marker for investigating phylogeographic structuring. *Mar. Biotechnol*, 7: 462-473.

**Dirienzo, A., Peterson, A.C., Garza, J.C., Valdes, A.M., Slatkin, M & Freimer N.B (1994)** Mutational processes of simple-sequence repeat loci in human-populations. *Proceedings of the National Academy of Sciences of the United States of America*, 91: 3166-3170.

**Donini, P., Stephenson, P., Bryan, G.J & Koebner, R.M.D (1998)** The potential of microsatellites for high throughput genetic diversity assessment in wheat and barley. *Genet Res Crop Evol*, 45: 415-421

**Edwards, A., Civitello, A., Hammond, H.A & Caskey, C.T. (1991)** DNA typing and genetic mapping with trimeric and tetrameric tandem repeats. *Am. J. Hum. Genet*, 49. 746-756.

**Einum, S & Fleming, I.A. (1997)** Genetic divergence and interactions in the wild among native, farmed and hybrid Atlantic salmon. *Journal of Fish Biology*, 50: 634-651.

**Ellsworth, D.L., Rittenhouse, K.D & Honeycutt, R.L (1993)** Artifactual variation in randomly amplified polymorphic DNA banding patterns. *Biotechniques*, 14: 214-217.

**Estoup, A. & Angers, B. (1998)** Microsatellites and minisatellites for molecular ecology: theoretical and empirical considerations. In: Carvalho, G. (ed.). *Advances in Molecular Ecology*, IOS Press: 55-86.

**FAO, (2001)** Report on the FAO/DANIDA/CFRAMP/WECAFC Regional Workshops on the assessment of the Caribbean Spiny lobster (*Panulirus argus*). FAOFish. Report No. 619 (ISSN 0429-9337). p. 381.

**Felsenstein, J. (1978)** Cases in which parsimony or compatibility methods will be positively misleading. *Systematic Zoology* 27: 401-410.

**Ferguson, A., Taggart, J.B., Prodohl, P.A., McMeel, O., Thompson, C., Stone, C., McGinnity, P. and Hynes, R.A. (1995)** The application of molecular markers to the study and conservation of fish populations, with special reference to *Salmo*. *Journal of Fish Biology* 47: 103-126.

**Fetzner Jr, J. W., & K. A. Crandall. (2001)** Genetic Variation, pp. 291- 326. In, D. M. Holdich (ed.), *Biology of Freshwater Crayfish. Blackwell Science, Oxford*.27.

**Foo, C.L., Dinesh, K.R., Lim, T.M., Chan, W.K. & Phang, V.P.E. (1995)** Inheritance of RAPD markers in the guppy fish, *Poecilia reticulata*. *Zoological Science*,**12**: 535-541.

**Frankel, O.H. & Soule, M.E. (1981)** Conservation and Evolution. Cambridge University Press, Cambridge.

**Fraser, D.J, & L. Bernatchez (2001)** Adaptive evolutionary conservation: towards a unified concept for defining conservation units. *Molecular Ecology* 10: 2741-2752.

**Fuchs H., Gross R, Stein H and Rottmann O (1998)** Application of molecular genetic markers for the differentiation of bream (*Abramis brama* L.) populations from the rivers Main and Danube. *J Appl Ichthyol*,14:49-55.

**Gaggiotti, O.E., Lange, O., Rassmann, K,& Gliddon, C (1999).** A comparison of two indirect methods for estimating average levels of gene flow using microsatellite data. *Molecular Ecology*, 8: 1513-1520.

**Gaggiotti, OE& Vetter, R.D (1999).** Effect of life history strategy, environmental variability, and overexploitation on the genetic diversity of pelagic fish populations. *Canadian Journal of Fisheries and Aquatic Sciences*. 56: 1376-1388.



**Galindo, H.A., Olson, D.B, & Palumbi, S.R (2006)** Genetics: a coupled oceanographic-genetic model predicts population structure of Caribbean corals. *Current Biology*. **16**, 1622–1626.

**Galtier, N & M. Gouy (1998)** Inferring pattern and process: maximum likelihood implementation of a non-homogeneous model of DNA sequence evolution for phylogenetic analysis. *Molecular Biology and Evolution*. 15: 871-879.

**Galtier, N & M. Gouy. (1995).** Inferring phylogenies from DNA sequences of unequal base compositions. Proceedings of the National Academy of Sciences of the United States of America 92: 11317-11321.

**Garcia, D.K & J.A.H. Benzie (1995)** RAPD markers of potential use in penaeid prawn (*Penaeus monodon*) breeding programs. *Aquaculture*, 130: 137-144.

**Garcia, D.K., Dhar, A.K & Alcivar-Warren A. (1996).** Molecular analysis of a RAPD marker (B20) reveals two microsatellites and differential mRNA expression in *Penaeus vannamei*. *Mol Marine Biol Biotechnol*, **5**:71.

**George, W (1997)** Tectonic plate movements and the evolution of Jasus and Panulirus spiny lobsters (Palinuridae). *Marine freshwater research*. 48: 1121-1130.

**George, R.W & Main, A.R (1967)** "The evolution of spiny lobsters (Palinuridae): a study of evolution in the marine environment". [\*Evolution\*](#) **21** (4): 803–820.

**Grant, W. S & R. S. Waples (2000)** Spatial and temporal scales of genetic variability in marine and anadromous species: *Implications for fisheries oceanography*. Pages 61-93.

**Goldman, N & Z. Yang. (1994)** A codon-based model of nucleotide substitution for protein-coding DNA sequences. *Molecular Biology and Evolution*, 11: 725-736.

**Goldstein, D.B& Schlotterer, C. (1999).** *Microsatellites: Evolution and Application*. Oxford univ. press, NewYork.

**Goldstein, D.B., Linares, A.R., L.L. Cavalli-sforza, and M.W. Feldman (1995)** An evaluation of Genetic distances for use with Microsatellite Loci. *Genetics*, 139(1) 463-471.

**Gonzalez, I. L., J. E. Sylvester., T. F. Smith., D. Stambolian, & R. D. Schmickel (1990)** Ribosomal RNA gene sequences and hominoid phylogeny. *Molecular Biology and Evolution*, 7: 203-219.

**Gopal, K., Tolley, K.A., Groeneveld, J.C., Matthe, C.A., (2006)** Mitochondrial DNA variation in spiny lobster (*Palinurus dalagoae*) suggests genetically structured populations in the southwestern Indian Ocean. *Mar.Ecol.Prog.Ser*, 319,191-198.

**Grace P. McCormack., Richard Powell & Brenden F. Keegan (2000)** Comparative Analysis Of two populations Of the Brittle Star (*Ampiuira filiformis*) (*Echinodermata: Ophiurodea*) with Different Life History Stratagies using RAPD markers. *Mar.Biotechnol*, 2: 100-106.

**Grandjean, F., N. Guoin, M. Frelon, & C. Souty-Grosset (1998)** Genetic and morphological systematic studies on the crayfish (*Austropotamobius pallipes*) *Journal of Crustacean Biology*, 18: 549-555.

**Graves, J.E (1998)** Molecular insights into the population structures of cosmopolitan marine fishes. *The Journal of Heredity*, 89:427-437.

**Hall, T.A. (1999)** BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95-98.

**Hall T.A (1990).** BIOEDIT: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucl. Acids Symp. Ser.* 41, 95-98.

**Hallerman, E.M. (2003)** Population genetics: principles and applications for fisheries scientists. American Fish Hansen, M.M. 2003.

**Hansen, M.M. (2003)** Application of molecular markers ineries Society, Bethesda, Maryland: 458 pp. aquaculture. *World Journal of Microbiology and Biotechnology*, 13: 405-413.

**Harding, G. C., E. L. Kenchington, C. J. Bird, C. S. Pezzack, & D. C. Landry (1997)** Genetic relationships among subpopulations of the American lobster (*Homarus americanus*) as revealed by random amplified polymorphic DNA. *Canadian Journal of Fisheries and Aquatic.*

**Hasegawa, M., H. Kisino, & T. Yano (1985)** Dating of the human-ape splitting by a molecular clock of mitochondrial DNA. *Journal of Molecular Evolution*, 21: 160-174.

**Hasegawa, E& E. Kasuya (2006)** Phylogenetic analysis of the insect order Odonata using 28S and 16S rDNA sequences: a comparison between data sets with different evolutionary rates. *Entomological Science* 9: 55-66.

**Hasegawa, M & T. Hashimoto (1993)** Ribosomal RNA trees misleading? *Nature* 361:23.

**Hauser, L., Adcock, G.J., Smith, P.J., Bernal-Ramirez, J.H & Carvalho, G.R (2002)** Loss of microsatellite diversity and low effective population size in an overexploited population of New Zealand snapper (*Pagrus auratus*). *Proceedings of the National Academy of Sciences of the United States of America*, 99: 11742-11747.

**Heath, D.D., Iwana, G.K, Delvin, R.H (1993).** PCR primed with VNTR core sequences yield species specific patterns and hypervariable probes. *Nucl Acids Res*, 21:5782–5785.

**Hedgecock, D (1994)** Does variance in reproductive success limit effective population sizes of marine organisms? pp 122-134 in *Genetics and Evolution of Aquatic Organisms*, edited by AR Beaumont. Chapman, Hall, London.

**Herbinger, C.M. Doyle, R.W., Pitman., E.R., Paquet, D., Mesa, K.A. Morris., D.B. Wright, J.M& Cook, D. (1995)** DNA fingerprint based analysis of paternal and maternal effects on offspring growth and survival in communically reared rainbow trout. *Aquaculture*, 137: 245-256.

**Holthuis, L.B (1991)** *Marine Lobsters of the World*. An Annotated and Illustrated Catalogue of Species of Interest to Fisheries Known to Date, FAO Fisheries Synopsis 125. FAO, Rome.

**Hutchings, J.A (2000)** Collapse and recovery of marine fishes. *Nature*, 406: 882-885.

**Hutchings, J.A & Gerber, L (2002)** Sex-biased dispersal in a salmonid fish. *Proc. R. Soc. London. B*. 269: 2487-2493.

**Hutchinson, W.F., Van Oosterhout, C., Rogers, S.I& Carvalho, G.R (2003)** Temporal analysis of archived samples indicates marked genetic changes in declining North Sea cod (*Gadus morhua*). *Proceedings of the Royal Society of London Series B-Biological Sciences*, 270: 2125-2132.

**Ihssen, P.E., Booke, H.E., Casselman, J.M., McGlade, J.M., Payne, N.R. & Utter, F.M. (1981)** Stock identification: materials and methods. *Can. J. Fish.Aquat. Sci.*, 38: 1838-1855.

**Jaccard, P. (1901).** Étude comparative de la distribution florale dans une portion des Alpes et des Jura. *Bull. Soc. Vaudoise Sci. Na.* 37: 547-579.

**Jarman, S. N., S. Nicol, N., G. Elliott, & A. McMinn (2000)** 28S rDNA evolution in the Eumalacostraca and the phylogenetic position of krill. *Molecular Phylogenetics and Evolution*, 17: 26-36.

**Jarne, P & Lagoda, P.J.L (1996)** Microsatellites, from molecules to populations and back. *Trends in Ecology and Evolution*, 11: 424-429.

**Jesus, J., D. J. Harris, & A. Brehm (2007)** Relationships of *Afroablepharus* Greer, 1974 skinks from the Gulf of Guinea islands based on mitochondrial and nuclear DNA: Patterns of colonization and comments on taxonomy. *Molecular Phylogenetics and Evolution*, 45: 904-914.

**Jin, L & M. Nei (1990)** Limitations of the evolutionary parsimony of phylogenetic analysis. *Molecular Biology and Evolution*, 7: 82-102.

**Johnson, S.L., Midson, C.N., Ballinger, E.W. & Postlethwaite J.H (1994)** Identification of RAPD primers that reveal extensive polymorphisms between laboratory strains of zebrafish. *Genomics*, 19. 152-156.

**Jorge Fraga (1), Jinnay Rodriguez(2), Omar Fuentes(2), Aymé Fernandez-Calienes (1) & Mayda Castex(2) (2005).** Optimization of random amplified polymorphic DNA techniques for use in genetic studies of *Cuban triatominae*. *Rev. Inst. Med. trop. S. Paulo*, 47(5): 295-300.

**Kagwade, P. V., Manickaraja, M., Deshmukh, V. D., Rajamani, M., Radhakrishnan, E. V., Suresh, V., Kathirvel, M & Rao, G. S (1991).** Magnitude of lobster resources of India. *Journal of Marine Biological Association of India* 33(1&2): 150–158.

**Kaizians, S., Morzot, D.C., McEntire, B.B., Nairn, R.S & Borowsky, R.L (1996).** Genetic mapping in Xiphophorous hybrid fish: assignment of 43 AP-PCR/RAPD and isozyme markers to multipoint linkage groups. *Genome Res.* 6: 280-289.

**Kallio-Nyberg, I & Koljonen, ML (1997).** The genetic consequence of hatchery rearing on life-history traits of Atlantic salmon (*Salmo salar* L.): A comparative analysis of sea-ranched salmon with wild and reared parents. *Aquaculture*, **153**: 207-224

**Kikuchi, M., Ikeda, M & Fujio, Y (1997)** Genetic variability and differentiation within and among populations of Japanese char: a comparison between isozyme and RAPD polymorphism. *Fish Genetics and Breeding Science*, **25**: 27-36.

**Kimura, M. (1980)** A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Biology*. **16**: 111-120.

**Klinbunga, S. A., Boonyapakdee & B. Pratoomchat (2000)** Genetic diversity and species diagnostic markers of mud crabs (genus *Scylla*) in eastern Thailand determined by RAPD analysis. *Marine Biotechnology*, **2**: 180-187.

**Koh T.L., Khoo, G., Fan, L.Q & Phang, V.P.E (1999)** Genetic diversity among wild forms and cultivated varieties of Discus (*Symphysodon spp.*) as revealed by Random Amplified Polymorphic DNA (RAPD) fingerprinting. *Aquaculture*, **173**, 485-497.

**Konstantinou, H & Shen, D.C (1995)** The social and reproductive behavior of the eyed flounder, *Bothus ocellatus*, with notes on the spawning of *Bothus lunatus* and *Bothus ellipticus*. *Environmental Biology of Fishes*, **44**: 311-324.

**Kumar, S.K., K. Tamura, & M. Nei (2004)** MEGA3: Intergrated software for molecular evolutionary genetics analysis and sequence alignment. *Briefings in Bioinformatics*, **5**: 150-163.

**Lecointre, G., & Deleporte, P. (2005).** Total evidence requires exclusion of phylogenetically misleading data. *Zoologica Scripta*, **34**(1):101–117.

**Levinson, G & Gutman, G.A (1987)** Slipped-Strand Mispairing - A major mechanism for DNA-sequence evolution. *Molecular Biology and Evolution*, 4: 203-221.

**Li, W.H., M. Gouy, P.M Sharp, C. O'hUigin, & Y.W Yang (1990)** Molecular phylogeny of Rodentia, Lagomorpha, Primates, Artidactyla and Carnivore and molecular clocks. *Proceedings of the National Academy of Sciences of the United States of America*, 87: 6703-6707.

**Liu, Z.J. P Li, BJ Argue, RA Dunham, (1999)** Random amplified polymorphic DNA markers: usefulness for gene mapping and analysis of genetic variation of catfish *Aquaculture*, 174: pp. 59-68.

**Liu, Z. G., C. Z. Lei, J. Luo, C. Ding, G. H. Chen, H. Chang, K. H. Wang, X. X. Liu, X. Y. Zhang, X. J. Xiao & S. L. Wu (2004)** Genetic variability of mtDNA sequences in Chinese native chicken breeds. *Asian-Aust. J. Anim. Sci*, 17:903-909.

**Liu, M. Y., Y. X. Cai, & C. S. Tzeng (2007)** Molecular systematics of the freshwater prawn genus *Macrobrachium* Bate, 1868 (Crustacea: Decapoda: Palaemonidae) inferred from mtDNA sequences, with emphasis on East Asian species. *Zoological Studies*, 46: 272-289.

**Lockhart, P. J., M. A. Steel, M. D. Hendy, & D. Penny (1994)** Recovering evolutionary trees under a more realistic model of sequence evolution. *Molecular Biology and Evolution*, 11: 605-612.

**Loomis, W.F & D. W. Smith (1990)** Molecular phylogeny of *Dictyostelium discoideum* by protein sequence comparison. *Proceedings of the National Academy of Sciences of the United States of America*, 87: 9093-9097.

**Luikart, G., Cornuet, J.M & Allendorf, F.W (1999)** Temporal changes in allele frequencies provide estimates of population bottleneck size. *Conservation Biology*, 13: 523-530.

**Luikart, G & England, P.R (1999)** Statistical analysis of Microsatellite DNA data. *Trends in Ecology and Evolution*, 14: 253-256.

**Lynch, M. 1994.** The neutral theory of phenotypic evolution, pp. 86-108. In L. Real (ed.) *Ecological Genetics*.

**Machordom, A & E. Macpherson (2004)** Rapid radiation and cryptic speciation in squat lobsters of the genus *Munida* (Crustacea, Decapoda) and related genera in the South West Pacific: molecular and morphological evidence. *Molecular Phylogenetics and Evolution*, 33: 259-279.

**Malau-Aduli, A. E., O. A. Nishimura-Abe, T. Niibayashi, Y. Yasuda, T. Kojima, S. Abe, K. Oshima, K. Hasegawa & M. Komatsu (2004)** Mitochondrial DNA Polymorphism, Maternal Lineage and Correlations with Postnatal Growth of Japanese Black Beef Cattle to Yearling Age. *Asian-Aust. J. Anim. Sci*, 17(11):1484-1490.

**Misof, B., A. M. Rickert, T. R. Buckely, G. Freck & K. P. Sauer (2001)** Phylogenetic signal and its decay in mitochondrial SSU and LSU rRNA gene fragments of Anisoptera. *Molecular Biology and Evolution*, 18: 27-37.

**Miyamoto, M. M., F. Kraus & O. A. Ryder (1990)** Phylogeny and evolution of antlered deer determined from mitochondrial DNA sequences. *Proceedings of the National Academy of Sciences of the United States of America* 87: 6127-6131.

**McVey, J.P. (Ed.). (1993)** CRC, Handbook of mariculture, vol. J. Crustacean Aquaculture, CRC press, Boca raton, F.L, USA.

**McConnell, S.K., O'Reilly, P., Hamilton, L., Wright, J.N & Bentzen, P. (1995)** Polymorphic microsatellite loci from Atlantic salmon (*Salmo salar*) - genetic differentiation of North American and European populations. *Can. J. Fish. Aquat. Sci.*, 52: 1863–1872.



**Mohamed, K. H., George, M. J. (1968):** Results of the tagging experiments on the Indian spiny lobster, *Panulirus homarus* (Linnaeus)—movement and growth. *Indian Journal of Fisheries*, 15: 15–26.

**Morrison, C. L., R. Rios, and J. E. Duffy (2004)** Phylogenetic evidence for an ancient rapid radiation of Caribbean sponge dwelling snapping shrimps (*Synalpheus*). *Molecular Phylogenetics and Evolution*, 30: 563- 581.

**Muraji, M & S. Tachikawa (2000)** Phylogenetic analysis of water striders (*Hemiptera: Gerroidea*) based on partial sequences of mitochondrial and nuclear ribosomal RNA genes. *Entomological Science*, 3(4): 615-626.

**Murphy, N.P & C.M Austin (2005)** Phylogenetic relationships of the globally distributed freshwater prawn genus *Macrobrachium* (Crustacea: Decapoda: Palaemonidae): biogeography, taxonomy and the convergent evolution of abbreviated larval development. *Zoologica Scripta* 34(2): 187-197.

**Murphy, N.P & C.M Austin (2004)** Multiple origins of the endemic Australia *Macrobrachium* (Decapoda: Palaemonidae) based on 16S rRNA mitochondrial sequences. *Australian Journal of Zoology*, 52: 549-559

**Murphy, N.P & C.M Austin (2003)** Molecular taxonomy and phylogenetics of some species of Australian palaemonid shrimps. *Journal of Crustacean Biology*, 23: 169-177.

**Murphy, N.P & C.M Austin (2002)** A preliminary study of 16S rRNA sequence variation in Australian *Macrobrachium* shrimps (Palaemonidae: Decapoda) reveals inconsistencies in their current classification. *Invertebrate Systematics*, 16: 697-701.

**Muse, S. V (1996)** Estimating synonymous and nonsynonymous substitution rates. *Molecular Biology and Evolution* 13: 105-114.

**Muse, S. V (1995)** Evolutionary analyses of DNA sequences subject to constraints on secondary structure. *Genetics* 139: 1429-1439.

**Nadig S.G., Lee KL and Adams SM (1998)** Evaluating alterations of genetic diversity in sunfish populations exposed to contaminants using RAPD assay. *Aquat Toxicol*, 43:163-178.

**Navidi, W.C., G.A. Churchill, & A. Von Haeseler (1991)** Methods for inferring phylogenies from nucleic acid sequence data by using maximum likelihood and linear invariants. *Molecular Biology and Evolution* 8: 128-143.

**Nei, M (1972)** Genetic distance between populations. *Am. Nat*, **106**: 283-292.

**Nei, M., Maruyama, T & Chakraborty, R (1975)** Bottleneck effect and genetic-variability in populations. *Evolution*, 29: 1-10.

**Nei, M., Tajima, F & Tateno, Y (1983)** Accuracy of estimated Phylogenetic tree from molecular data, II: gene frequency data. *J.Mol. Evol*, **19**: 153–170.

**Nei, M (1987)** *Molecular Evolutionary Genetics*. Columbia University Press, New York.

**Nelson, K., and D. Hedgecock. (1977)** Electrophoretic evidence of multiple paternity in the lobster *Homarus americanus* (Milne-Edwards). *American Naturalist*, 111:361-365.

**Nielsen, E.E, Hansen, M.M & Loeschcke, V (1999)** Analysis of DNA from old scale samples: technical aspects, applications and perspectives for conservation. *Hereditas*, 130: 265-276.

**O'Connell, M & Wright, J.M (1997)** Microsatellite DNA in fishes. *Reviews in Fish Biology and Fisheries*, 7: 331-363.

**Ovendon, J. R., J.D. Booth, & A.J. Smolenski (1997)** Mitochondrial DNA phylogeny of red and green rock lobsters. *Marine and Freshwater Research*, 48: 1131-1136.

**Page, R.D.M & Holmes, E.C (1998)** *Molecular Evolution: A Phylogenetic Approach*. Blackwell Science, Oxford, U.K.

**Palumbi, S.R & J.Benzie (1991)** Large mitochondrial DNA differences between morphologically similar penaeid shrimp. *Molecular Marine Biology and Biotechnology* 1: 27-34.

**Palumbi, S.R (2004)** Marine reserves and ocean neighborhoods: The spatial scale of marine populations and their management. *Annual Review of Environment and Resources*, 29, 31–68.

**Pardini, A.T., Jones, C.S., Noble, L.R., Kreiser, B., Malcolm, H., Bruce, B.D., Stevens J.D., Cliff, G., Scholl, M.C., Francis, M., Duffy, C.A.J & Martin, A.P (2001)** Sex-biased dispersal of great white sharks - In some respects, these sharks behave more like whales and dolphins than other fish. *Nature*, **412**: 139-140.

**Partis, L. & R.J. Wells (1996)** Identification of fish species using random amplified polymorphic DNA (RAPD). *Mol. Cell. Probes*, 10:435-441.

**Pawson, M.G & Jennings, S (1996)** A critique of methods for stock identification in marine capture fisheries. *Fisheries Research*, 25: 203-217.

**Pelc, R.A., Warner, R.R & Gaines, S.D (2009)** Geographical patterns of genetic structure in marine species with contrasting life histories. *Journal of Biogeography*, 36, 1881–1890.

**Perez-Enriquez, R., Takemura, M & Taniguchi, N (1999)** Genetic Diversity of red sea bream (*Pagrus major*) in western Japan in relation to stock enhancement. *Fish. Sci*, **67**: 374-379.

**Perez-Enriquez, R & Taniguchi, N (2001)** Genetic structure of red sea bream (*Pagrus major*) population off Japan and Southwest Pacific, Using microsatellites DNA markers. *Fish. Sci.*, **65**: 23-30.

**Pollock, D. E (1993)** Speciation in spiny lobsters – clues to climatically-induced changes in ocean circulation patterns. *Bulletin of marine science*, 53(3): 937-944.

**Ponniah, K & Crandall, A (2005)** Phylogeny and biogeography of the freshwater crayfish *Euastacus* (*Decapoda: Parastacidae*) based on nuclear and mitochondrial DNA. *Molecular Phylogenetics and Evolution*, 37: 249-263.

**Posada, D & K.A. Crandall (1998)** Modeltest: testing the model of DNA substitution. *Bioinformatics*, 14: 817-818.

**Postlethwait, J.H., Jhonson, S.L., Midson, C.N., Talbot, W.S., Gates, M., Ballinger, E.W., Africa, D., Andrews, R., Carl, T., Eisen, J.S., Horne, S., Kimmel, C.B., Hutchison, M., Johnson, M & Rodriguez, A (1994)** A genetic linkage map for the zebrafish. *Science*, **264**: 699-703.

**Primavera, J.H (1985)** A review of maturation and reproduction in closed-*thelycum penaeids*. In: Taki, Y., Primavera J.H. Lobrera. I.A. (Eds), Proceedings of the first International Conference on the Culture of *Penaeid* Prawns, 4-7 December 1984, at Hoito City, Philippines, pp. 47-64.

**Pusey, A (1987)** Sex-biased dispersal and inbreeding avoidance in birds and animals. *Trends in Ecology and Evolution*, 2: 295–299.

**Qi Li, Choulji Park, Takashi Endo & Akhiro Kijima (2004)** Loss of genetic variatioat microsatellite loci in hatchery strains of the Pacific abalone (*Haliotis discus hannai*). *Aquaculture*, **235**: 207-222.

**Rafalski, (2002)** Applications of single nucleotide polymorphisms in crop genetics. *Curr Opin Plant Biol*, 5:94–100.

**Radhakrishnan, E.V & Manisseri, M.K (2003)** Lobsters. *In: Mohan Joseph, M.; Jayaprakash, A. A. ed. Status of exploited marine fishery resources of India. Central Marine Fisheries Research Institute, Kochi–682014, India. PP 195–202.*

**Radhakrishnan, E.V., V.D Deshmukh., Mary K. Manisseri; M. Rajamani., Joe K. Kizhakudan & R. Thangaraja ( 2005)** Status of the major lobster fisheries in India *New Zealand Journal of Marine and Freshwater Research, Vol. 39: 723–732*

**Reilly, P & Wright, J.M (1995)** The evolving technology of DNA fingerprinting and its application to fisheries and aquaculture. *J.Fish. Biol.* **47**: 29-55.

**Rijnsdorp, A.D & Jaworski, A (1990)** Size-selective mortality in plaice and cod eggs - a new method in the study of egg mortality. *Journal du Conseil International pour Exploration de la Mer*, 47: 256-263.

**Rijnsdorp, A.D (1991)** Changes in fecundity of female north-sea plaice (*Pleuronectes platessa* L) between 3 periods since 1900. *ICES Journal of Marine Science*, 48: 253-280.

**Rijnsdorp, A.D (1994)** Population regulating processes during the adult phase in flatfish. *Netherlands Journal of Sea Research*, 32: 207-223.

**Rocha, L.A., Lindeman, K.C, Rocha, C.R. & Lessios, H.A. (2008)** Historical biogeography and speciation in the reef fish genus *Haemulon* (Teleostei: Haemulidae). *Molecular Phylogenetics and Evolution*, 48: 918–928.

**Ryman, N., F.Utter & K. Hindar (1995a)** Introgression, supportive breeding, and genetic conservation. In *Population Management for Survival and Recovery* (Ballou, J.D. et al., eds), pp. 341-365, Columbia University Press.

**Ryman, N., Utter, F, and Laikre,L, (1995b)** Protection of intraspecific biodiversity of exploited fishes. *Rev. Fish Biol. Fish.* 5: 417-446.

**Ryman, N (1991)** Conservation genetics considerations in fishery management. *J. Fish Biol*, **39**: 211-214.

**Rozas, J., J. C. Sanchez-DelBarrio, X. Messeguer & R. Rozas (2003)** DnaSP, DNA polymorphism analyses by the coalescent and other methods. *Bioinformatics*, **19**: 2496-2497.

**Sarver, S.K., J.D. Silberman & P.J. Walsh (1998)** Mitochondrial DNA sequence evidence supporting the recognition of two subspecies or species of the Florida Spiny Lobster. *Journal of Crustacean Biology*, **18**: 177-186.

**Saitou, N & M. Nei (1987)** The Neighbor-Joining Method: a new method for reconstructing Phylogenetic trees. *Molecular Biol. Evol*, **4**: 406-425.

**Sbordoni, V., De Matthaeis, E., Cobolli-Sbordoni, M. Larosa. G. Mattoccia, M (1987)** Genetic changes in seven generations of hatchery stocks of the Kuruma prawn, (*Penaeus japonicas*) (Crustacean, Decapoda). Proc. World symp. On selection, Hybridization and Genetic Engineering in Aquaculture, Bordeaux, **1**:27-30.

**Schoniger, M & A. Von Haeseler (1995)** Performance of the maximum likelihood, neighbor joining, and maximum parsimony methods when sequence sites are not independent. *Systematic Biology*, **44**: 533-547. *Science*, **54**: 1762-1771.

**Schug, M.D., Hutter, C.M., Wetterstrand, K.A., Gaudette, M.S., Mackay, T.F.C & Aquadro, C.F (1998)** The mutation rates of di-, tri- and tetranucleotide repeats in *Drosophila melanogaster*. *Mol. Biol. Evol*, **15(12)**: 1751--1760.

**Shanks, A.L., B.A. Grantham & M.H. Carr (2003)** Propagule dispersal distance and the size and spacing of marine reserves. *Ecological Applications* **13**:S159–S169.

**Shull, H. C., M. Perez-Losada, D. Blair, K. Sewell, E. A. Sinclair, S. Lawler, M.**

**Shy, J. Y & H. P. Yu (1998)** Freshwater Shrimps of Taiwan. National Museum of Marine Biology and Aquarium, Pingdong, Taiwan. 103 pp. Sidow, A., and T. P. Steel. 1992. Estimating the fraction of invariable codons with a capture-recapture method. *Journal of Molecular Evolution*, 35: 253-260.

**Sidow, A & T. P. Steel (1992)** Estimating the fraction of invariable codons with a capture-recapture method. *Journal of Molecular Evolution*, 35:253-260.

**Silberman, J.D., S.K. Sarver & P.J Walsh (1994)** Mitochondrial DNA variation and population structure in the spiny lobster (*Panulirus argus*). *Mar. Biol.*, Vol. 120, pp. 601-608.

**Skibinski, D.O.F (1994)** DNA technology and genetics of aquatic invertebrates. In, *Genetics and Evolution of Aquatic Organisms*. Edited by A.R. Beaumont. Chapman and Hall, London. pp. 177-199.

**Slatkin, M (1995)** A measure of population subdivision based on microsatellite allele frequencies. *Genetics*, 139: 457-462.

**Smith, P.J., Francis, R.I.C.C & McVeagh, M (1991)** Loss of genetic diversity due to fishing pressure. *Fisheries Research*, 10: 309-316.

**Stoner, A.W., Bedja, A.J., Manderson, J.P., Phelan, B.A., Stehlik, L.L & Pessutti, J.P (1999)** Behaviour of winter flounder, (*Pseudopleuronectes americanus*) during the reproductive season: laboratory and field observation on spawning, feeding, and locomotion. *Fisheries Bulletin*, 97: 999-1016.

**Sullivan, J., K. E. Holsinger & C. Simon (1995)** Among-site rate variation and phylogenetic analysis of the 12S rRNA in Sigmodontine rodents. *Molecular Biology and Evolution* 12: 988-1001.

**Sultmann, H., Mayer, W.E., Figuerora, F., Tichy, H & Klein, J (1995)** Phylogenetic analysis of *cichlid fishes* using nuclear DNA markers. *Mol.Biol.Evol*, **12**: 1033-1047.

**Sunden, S.P.F. And Davis S.K (1991)** Evaluation of genetic variation in a domestic populations of *Penaeus vannamei* (Boone): a comparison with three natural populations. *Aquaculture*, **97**:131-142.

**Swofford, D.L., G.L. Olsen, P. J. Waddell & D. M. Hillis (2002)** PAUP\*: Phylogenetic analysis using parsimony (and other methods) 4.0.b10. Sinauer Associates Sunderland, Massachusetts.

**Swofford, D.L., G.L. Olsen, P.J. Waddell & D.M. Hillis (1996)** Phylogenetic inference, pp. 407-514. In, D. M. Hillis, C. Moritz, and B. K. Mable (eds.), *Molecular Systematics*. Sinauer, Sunderland, MA.

**Tamura, K & M. Nei (1993)** Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution* 10: 512-526.

**Tam, Y.K & Kornfield, I (1996)** Characterization of microsatellite markers in *Homarus* (Crustacea, Decapoda). *Molecular Marine Biology and Biotechnology*, 5: 230–238.

**Tassanakajon, A., Pongsomboon, S., Rimphanitchayakil, V., Jarayabhaud, P& Boonsaeng, V ( 1997)** Random amplified polymorphic DNA (RAPD) markers for determination of genetic variation in wild population of the black tiger prawn (*Penaeus monodon*) in Thailand. *Mol. Mar. Biol. Biotechnol*, 6:110–115.

**Tateno, Y., N. Takezaki & M. Nei (1994)** Relative efficiencies of the maximum likelihood, neighbor joining, and maximum parsimony methods when substitution rate varies with site. *Molecular Biology and Evolution*, 11: 261-277.



**Taylor, M.S., & Hellberg, M.E. (2003)** Genetic evidence for local retention of pelagic larvae in a Caribbean reef fish. *Science*, 299:107–109.

**Thompson, and J. D., Higgins, D.G., & Gibson, T.J (1994)** CLUSTALW: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, positions-specific gap penalties and weight matrix choice. *Nucleic Acids Res* 22: 4673-4680.

**Tolley, K.A., Groeneveld, J.C., Gopal, K & Matthee, C.A (2005)** Mitochondrial DNA panmixia in spiny lobster (*Palinurus gilchristi*) suggests a population expansion. *Mar Ecol Prog Ser*, 297:225–231

**Turner, T.F., Wares, J.P & Gold J.R (2002)** Genetic effective size is three orders of magnitude smaller than adult census size in an abundant, estuarine-dependent marine fish (*Sciaenops ocellatus*). *Genetics*, 162, 1329-1339.

**Utter, F.M (1991)** Biochemical genetics and fishery management: an historical perspective. *J Fish Biology*, 39: 1-20.

**Van der Veer, H.W., Pihl, L & Bergman, M.J.N (1990)** Recruitment mechanisms in North Sea plaice (*Pleuronectes platessa*) *Marine Ecology Progress Series*, 64: 1-12.

**Van Oppen, M.J.H., Willis, B.L & Miller, D.J (1999)** A typically low rate of cytochrome b evolution in the scleractinian coral genus *Acropora*. *Proc Roy Soc Lond B* 266: 179-183.

**Vignal, A., Milan, D, San Cristobal, M & Eggen, A (2002)** A review on SNP and other types of molecular markers and their use in animal genetics. *Gen Sel Evol*, 34:275-305.

**Wakeley, J (1993)** Substitution-rate variation among sites and the estimation of transition bias. *Molecular Biology and Evolution*, 11: 426-442.

**Wakeley, J (1996)** The excess of transitions among nucleotide substitutions: new methods of estimating transition bias underscore its significance. *Trends in Ecology & Evolution*, 11(4): 158-162.

**Waldman, J.P (1999)** The importance of comparative studies in stock analysis. *Fish Res* 43:237-246.

**Wang, J.L (2001)** A pseudo-likelihood method for estimating effective population size from temporally spaced samples. *Genetical Research*, 78: 243-257.

**Waples, R.S (1989)** A generalized-approach for estimating effective population-size from temporal changes in allele frequency. *Genetics*, 121: 379-391.

**Webster, M.T., Smith, N.G.C & Ellegren, H (2002)** Microsatellite evolution inferred from human-chimpanzee genomic sequence alignments. *Proceedings of the National Academy of Sciences of the United States of America*, 99: 8748-8753.

**Welsh, J & McClelland, M (1990)** Fingerprinting genomes using PCR with arbitrary primers. *Nucleic Acids Res*, 18: 7213–7218.

**Weir, P. S., and C. C. Cockerham. (1984).** Estimating F-statistics for the analysis of population structure. *Evolution*, 38:1358–1370.

**Wiens, J.J (1998)** Combining data sets with different phylogenetic histories. *Systematic Biology*, 47: 568-581.

**Williams, J.G., Kubelik, A.R., Livak, K.J., Rafalski, J.A & Tingey, S.V (1990)** DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acid Research*.

**Williamson, J.H (2001)** Brood stock management for imperilled and other fishes. In G.A. Wedemeyer, (Ed.). *Fish hatchery management*, second edition. American Fisheries Society, Bethesda, Maryland: 397-482.

**Williamson, E.G & Slatkin, M (1999)** Using maximum likelihood to estimate population size from temporal changes in allele frequencies. *Genetics*, 152: 755-761.

**World Register of Marine Species (WoRMS)**

<http://www.marinespecies.org/aphia.php?p=taxdetails&id=210355>.

**Wolfus, M.G., Denise K. Garcia, Acacia Alcivar-Warren (1997).** Application of the microsatellite technique for analyzing genetic diversity in shrimp breeding programs. *Aquaculture*. 152: 35-47.

**Wright, S (1969)** *Evolution and the genetics of populations*. Vol.2. The theory of gene frequencies. University of Chicago Press, Chicago.

**Wright, J.M & Bentzen, P (1995)** Microsatellites: genetic markers for the future. In: Carvalho, G.R., Pitcher, T.J. (Eds), *Molecular Genetics in Fisheries*. Chapman &hall, London, pp:117-121.

**Xia, X.Z. Xie. (2001)** DAMBE: Data analysis in molecular biology and evolution. *Journal of Heredity*, 92: 371-373.

**Xu, Z., J.H. Primavera, L.D. de la Pena, P. Pettit, J. Belak and A. Alcivar-Warren, (2001).** Genetic diversity of wild and cultured Black Tiger Shrimp (*Penaeus monodon*) in the Philippines using microsatellites. *Aquaculture*. 199: 13-40.

**Yang, Z (1993)** Maximum-likelihood estimation of phylogeny from DNA sequences when substitution rates differ over sites. *Mol Biol Evol* 10:1396-1401.

**Yang, Z (1996)** Among-site rate variation and its impact on phylogenetic analyses. *Trends in Ecology and Evolution* 11(9): 367-372.

**Yeh, F.C., Yang, R.C & T. Boyle (1999)** POPGENE version 1.31, Microsoft Windows-based freeware for population genetics analysis. University of Alberta and Centre for International Forestry Research, Research.

**Yuan, Y. X., Gao, C. N & Zhang, D.X (1992)** Egg hatching and metamorphosis to protozoa of (*Penaeus chinensis*) (osbeck) by removal of heavy metals from rearing systems with polymeric absorbent. *Aquaculture*, 107: 303-311.