

# References

## References

- Abbas AK, Murphy KM, Sher A. Functional diversity of helper T lymphocytes. *Nature*. 1996; 383:787–93.
- Abe T, Iinuma Y, Ando M, Yokoyama T, Yamamoto T, Nakashima K, et al. *NRAMP1* polymorphisms, susceptibility and clinical features of tuberculosis. *J Infect*. 2003; 46:215–20.
- Abecasis GR and Cookson WO. GOLD - Graphical Overview of Linkage Disequilibrium. *Bioinformatics* 2000; 16:182-3.
- Abecasis GR, Ghosh D, Nichols TE. Linkage disequilibrium: ancient history drives the new genetics. *Hum Hered* 2005; 59:118-124.
- Abel B, Thieblemont N, Quesniaux VJ, Brown N, Mpagi J, Miyake K, et al. Toll-like receptor 4 expression is required to control chronic *Mycobacterium tuberculosis* infection in mice. *J Immunol*. 2002; 169:3155-62.
- Abel L, Dessein AJ. The impact of host genetics on susceptibility to human infectious diseases. *Curr Opin Immunol* 1997; 9:509-16.
- Abhimanyu, Mangangcha IR, Jha P, Arora K, Mukerji M, Banavaliker JN, et al. Differential serum cytokine levels are associated with cytokine gene polymorphisms in north Indians with active pulmonary tuberculosis. *Infect Genet Evol*. 2011; 11:1015-22.
- Abu-Raddad, Laith J, Sabatelli, Lorenzo, Achterberg, Jerusha T, et al. Epidemiological benefits of more-effective tuberculosis vaccines, drugs, and diagnostics. *Proc Natl Acad Sci*. 2009; 106: 13980–5.
- Aderem A, Ulevitch RJ. Toll-like receptors in the induction of the innate immune response. *Nature*. 2000; 406:782-7.
- Lalvani A, Behr MA, Sridhar S. Innate Immunity to TB: a druggable balancing act. *Cell* 2012; 148:389-391.
- Akahoshi M, Nakashima H, Miyake K, Inoue Y, Shimizu S, Tanaka Y, et al. Influence of Interleukin-12 receptor  $\alpha 1$  polymorphism on tuberculosis. *Hum Genet*. 2003; 112:237- 43.
- Akgun M, Saglam L, Kaynar H, Yildirim AK, Mirici A, Gorguner M, et al. Serum IL-18 levels in tuberculosis: comparison with pneumonia, lung cancer and healthy controls. *Respirology*. 2005; 10:295-299.
- Aktas E, Ciftci F, Bilgic S, Sezer O, Bozkanat E, Deniz O, et al. Peripheral immune response in pulmonary tuberculosis. *Scand J Immunol*. 2009; 70:300-8.
- Al-Muhsen S, Casanova JL. The genetic heterogeneity of mendelian susceptibility to mycobacterial diseases. *J Allergy Clin Immunol*. 2008; 122:1043-1051.
- Alagarasu K, Selvaraj P, Swaminathan S, Narendran G, Narayanan PR. 5' regulatory and 3' untranslated region polymorphisms of vitamin D receptor gene in south Indian HIV and HIV-TB patients. *J Clin Immunol*. 2009; 29:196–204.
- Alcais A, Abel L. Application of genetic epidemiology to dissecting host susceptibility/resistance to infection illustrated with the study of common mycobacterial infections. In: Bellamy R ed. Cambridge: Cambridge University Press 2004 :7-33.
- Allie N, Keeton R, Court N, Abel B, Fick L, Vasseur V, et al. Limited role for lymphotoxin  $\alpha$  in the host immune response to *Mycobacterium tuberculosis*. *J Immunol* 2010; 185:4292–301.
- Allison MJ, Zappasodi P, Lurie MP. Host-parasite relationships in natively resistant and susceptible rabbits on quantitative inhalation of tubercle bacilli. Their significance for the nature of genetic resistance. *Am. Rev. Respir. Dis*. 1962; 85:553-569.
- Ameglio F, Casarini M, Capoluongo E, Mattia P, Puglisi G, Giosuè S. Post-treatment changes of six cytokines in active pulmonary tuberculosis: differences between patients with stable or increased fibrosis. *Int J Tuberc Lung Dis*. 2005; 98-104.
- Amirzargar AA, Rezaei N, Jabbari H, Danesh AA, Khosravi F, Hajabdolbaghi M, et al. Cytokine single nucleotide polymorphisms in Iranian patients with pulmonary tuberculosis. *Eur Cytokine Netw*. 2006; 17:84-9.
- Andrade DR junior, Santos SA, Castro I, Andrade DR. Correlation between serum tumor necrosis factor  $\alpha$  levels and clinical severity of tuberculosis. *Braz J Infect Dis*. 2008; 12:226-233.
- Ansari A, Talat N, Jamil B, Hasan Z, Razzaki T, Dawood G, et al. Cytokine gene polymorphism across tuberculosis clinical spectrum in Pakistani patients. *PLOS ONE*. 2009; 4: e4478.
- April MM, Garelick JM, Nuovo GJ. Reverse transcriptase in situ polymerase chain reaction in atypical mycobacterial adenitis. *Arch Otolaryngol Head Neck Surg*. 1996; 122:1214–18.
- Ates O, Musellim B, Ongen G, Topal-Sarikaya A, et al. Interleukin-10 and tumor necrosis factor- $\alpha$  gene polymorphisms in tuberculosis. *J Clin Immunol*. 2008; 28:232–6.
- Awomoyi AA, Charurat M, Marchant A, Miller EN, Blackwell JM, McAdam KP, et al. Polymorphism in IL1B: IL1B-511 association with tuberculosis and decreased lipopolysaccharide-induced IL-1 $\beta$  in IFN- $\gamma$  primed ex-vivo whole blood assay. *J Endotoxin Res*. 2005; 11:281-6.
- Babb C, Keet HE, van Helden PD, Hoal EG. SP110 polymorphisms are not associated with pulmonary tuberculosis in a

- 
- South African population. *Hum Genet.* 2007; 121:521–2.
- Babb C, van der Merwe L, Beyers N, Pheiffer C, Walzl G, Duncan K, et al. Vitamin D receptor gene polymorphisms and sputum conversion time in pulmonary tuberculosis patients. *Tuberculosis (Edinb)* 2007; 87:295–302.
- Baghdadi JE, Orlova M, Alter A, Ranque B, Chentoufi M, Lazrak F, et al. An autosomal dominant major gene confers predisposition to pulmonary tuberculosis in adults. *J Exp Med.* 2006; 203:1679-1684.
- Balding DJ. A tutorial on statistical methods for population association studies. *Nat Rev Genet* 2006; 7:781-791.
- Balsa A, Del Amo J, Blanco F, Caliz R, Silva L, Sanmarti R, et al. Prediction of functional impairment and remission in rheumatoid arthritis patients by biochemical variables and genetic polymorphisms. *Rheumatology.* 2010; 49:458-66
- Barnes PF, Abrams JS, Lu S, Sieling PA, Rea TH, Modlin RL. Patterns of cytokine production by mycobacterium-reactive human T-cell clones. *Infect Immun.* 1993; 61: 197–203.
- Barreiro LB, Neyrolles O, Babb CL, Tailleux L, Quach H, Ken McElreavey, et al. Promoter variation in the DC-SIGN-encoding gene CD209 is associated with tuberculosis. *PLoS Med.* 2006; 3:e20.
- Barrett JC, Fry B, Maller J, Daly MJ. Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics.* 2005; 21:263-5.
- Barry C E 3rd, Boshoff HI, Dartois V, Dick T, Ehrt S, Flynn J, Schnappinger D, et al. The spectrum of latent tuberculosis: rethinking the biology and intervention strategies. *Nat Rev Microbiol.* 2009; 7:845–55.
- Bates JH, Stead WW. The history of tuberculosis as a global epidemic. *Med Clin North Am* 1993; 77:1205-17.
- Beals M. The chi-square test. [www.tiem.utk.edu/~gross/bioed/bealsmodules/chisquare.html](http://www.tiem.utk.edu/~gross/bioed/bealsmodules/chisquare.html) . 2004.
- Bean AG, Roach DR, Briscoe H, France MP, Korner H, Sedgwick JD, et al. Structural deficiencies in granuloma formation in TNF gene-targeted mice underlie the heightened susceptibility to aerosol *Mycobacterium tuberculosis* infection, which is not compensated for by lymphotoxin. *J Immunol.* 1999; 162:3504–11.
- Becker T, Knapp M. A powerful strategy to account for multiple testing in the context of haplotype analysis. *Am J Hum Genet.* 2004; 75:561-570.
- Bekker LG, Maartens G, Steyn L, Kaplan G. Selective increase in plasma tumor necrosis factor-alpha and concomitant clinical deterioration after initiating therapy in patients with severe tuberculosis. *J Infect Dis.* 1998; 178:580–4.
- Bekpen C, Hunn JP, Rohde C, Parvanova I, Guethlein L, Dunn DM, et al. The interferon-inducible p47 (IRG) GTPases in vertebrates: loss of the cell autonomous resistance mechanism in the human lineage. *Genome Biol.* 2005; 6:R92.
- Bellamy R, Beyers N, McAdam KP, Ruwende C, Gie R, Samaai P, et al. Genetic susceptibility to tuberculosis in Africans: a genome-wide scan. *Proc Natl Acad Sci.* 2000; 97:8005-8009.
- Bellamy R, Ruwende C, Corrah T, McAdam KP, Thursz M, Whittle HC, et al. Tuberculosis and chronic hepatitis B virus infection in Africans and variation in the vitamin D receptor gene. *J Infect Dis.* 1999; 179:721–4.
- Bellamy R, Ruwende C, Corrah T, McAdam KP, Whittle HC, Hill AVS. Variations in the *NRAMP1* Gene and Susceptibility to Tuberculosis in West Africans. *N Engl J Med.* 1998; 338:640-4.
- Bellamy R. Genetics and pulmonary medicine. 3. Genetic susceptibility to tuberculosis in human populations. *Thorax* 1998; 53:588-93.
- Bellamy RJ, Hill AV. Host genetic susceptibility to human tuberculosis. *Novartis Found Symp.* 1998; 213:3-13.
- Bellamy RJ, Ruwende C, Corrah T, McAdam KP, Whittle, Hill AVS, et al. Assessment of the interleukin 1 gene cluster and other candidate gene polymorphisms in host susceptibility to tuberculosis. *Tuber. Lung Dis.* 1998; 79:83–9.
- Bellamy RR, Hill AV. Genetic susceptibility to mycobacteria and other infectious pathogens in humans. *Curr Opin Immunol.* 1998; 10:483-7.
- Ben-Ali M, Barbouche MR, Bousnina S, Chabbou A, and Dellagi K. Toll-like receptor 2 Arg677Trp polymorphism is associated with susceptibility to tuberculosis in Tunisian patients. *Clin Diagn Lab Immunol.* 2004; 11:625-6
- Biswas D, Qureshi O, Lee WY, Croudace J, Mura M, Lammas DM. ATP-induced autophagy is associated with rapid killing of intracellular mycobacteria within human monocytes/macrophages. *BMC Immunol.* 2008; 9:35.
- Blackwell JM, Goswami T, Evans CA, Sibthorpe D, Papo N, White JK, et al. SLC11A1 (formerly NRAMP1) and disease resistance. *Cell Microbiol.* 2001; 3:773–84.
- Bloch DB, Nakajima A, Gulick T, Chiche JD, Orth D, de La Monte SM, et al. Sp110 localizes to the PML Sp100 nuclear body and may function as a nuclear hormone receptor transcriptional coactivator. *Mol Cell Biol.* 2000; 20:6138–46.
- Bloom BR, Murray CJ. Tuberculosis: commentary on a reemerging killer. *Science* 1992; 257:1055-1064.
- Boraska V, Zeggini E, Groves CJ, Rayner NW, Skrabić V, Diakite M, et al. 2009. Family-based analysis of tumor necrosis factor and lymphotoxin- $\alpha$  tag polymorphisms with type 1 diabetes in the population of South Croatia. *Hum Immunol.* 2009; 70, 195-99.
- Bornman L, Campbell SJ, Fielding K, Bah B, Sillah J, Gustafson P, et al. Vitamin D receptor polymorphisms and susceptibility to tuberculosis in West Africa: a case-control and family study. *J Infect Dis.* 2004; 190:1631–41.
- Bossé Y, Lemire M, Poon AH, Daley D, He JQ, Sandford A, et al. Asthma and genes encoding components of the vitamin D pathway. *Respir Res.* 2009; 10: 98.
- Botstein D, Risch N. Discovering genotypes underlying human phenotypes: past successes for mendelian disease, future

- approaches for complex disease. *Nat Genet* 2003; 33:228-237
- Bottomley MJ, Collard MW, Huggenvik JI, Liu Z, Gibson TJ, Sattler M. The SAND domain structure defines a novel DNA-binding fold in transcriptional regulation. *Nat Struct Biol.* 2001; 8:626–33.
- Brightbill HD, Libraty DH, Krutzik SR, Yang RB, Belisle JT, Bleharski JR , et al. Host defense mechanisms triggered by microbial lipoproteins through tolllike receptors. *Science.* 1999; 285:732–6.
- Brodie D, Schluger NW. The diagnosis of tuberculosis. *Clin Chest Med.* 2005; 26: 247–71, vi.
- Buell GN, Talabot F, Gos A, Lorenz J, Lai E, Morris MA , et al. Gene structure and chromosomal localization of the human P2X7 receptor. *Receptor Channel* 1998; 5:347–54.
- Burgner D, Jamieson SE, and Blackwell JM. Genetic susceptibility to infectious diseases: big is beautiful, but will bigger be even better? *Lancet Infect Dis* 2006; 6:653-663.
- Burgner D, Jamieson SE, Blackwell JM. Genetic susceptibility to infectious diseases: big is beautiful, but will bigger be even better? *Lancet Infect Dis.* 2006; 6:653–63.
- Burgner D, Rockett K, Ackerman H, Hull J, Usen S, Pinder M , et al. Haplotypic relationship between SNP and microsatellite markers at the NOS2A locus in two populations. *Genes Immun.* 2003; 4:506-14.
- Burgner D, Rockett K, Kwiatkowski D. Nitric oxide and infectious diseases. *Arch Dis Childhood.* 1999; 81:185–88.
- Camus JC, Pryor MJ, Medigue C, Cole ST. Re-annotation of the genome sequence of *Mycobacterium tuberculosis H37Rv*. *Microbiology.* 2002; 148:2967-2973.
- Caner S, Köksal D, Ozkara S, Berkoğlu M, Aksaray S, Tarhan D. The relation of serum interleukin-2 and C-reactive protein levels with clinical and radiological findings in patients with pulmonary tuberculosis. *Tuberk Toraks.* 2007; 55:238-45.
- Canetti G, Rist N, Grosset J. Primary drug resistance in tuberculosis. *Am Rev Respir Dis.* 1964; 90:792-99.
- Canonne-Hergaux F, Gruenheid S, Govoni G, Gros P. The Nramp1 protein and its role in resistance to infection and macrophage function. *Proc Assoc Am Physicians.* 1999; 111: 283–9.
- Capon F, Allen MH, Ameen M, Burden AD, Tillman D, Barker JN , et al. A synonymous SNP of the corneodesmosin gene leads to increased mRNA stability and demonstrates association with psoriasis across diverse ethnic groups. *Hum Mol Genet.* 2004; 13:2361–8.
- Cardoso CC, Pereira AC, Brito-de-Souza VN, Dias-Baptista IM, Maniero VC, Venturini J, , et al. IFNG +874 T>A single nucleotide polymorphism is associated with leprosy among Brazilians. *Hum Genet.* 2010;128:481-90.
- Casanova JL, Abel L. Genetic dissection of immunity to mycobacteria: the human model. *Annu Rev Immunol.* 2002; 20:581e620.
- Casarini M, Ameglio F, Alemanno L, Zangrilli P, Mattia P, Paone G , et al. Cytokine Levels Correlate with a Radiologic Score in Active Pulmonary Tuberculosis. *Am J Respir Crit Care Med.* 1993; 159:143–8.
- Caws M, Thwaites G, Dunstan S, Hawn TR, Lan NT, Thuong NT, *et al.*, The influence of host and bacterial genotype on the development of disseminated disease with *Mycobacterium tuberculosis*. *PLoS Pathog* 2008; 4:e1000034.
- Cellier M, Govoni G, Vidal S, Kwan T, Groulx N, Liu J , et al. Human natural resistance-associated macrophage protein: cDNA cloning, chromosomal mapping, genomic organization, and tissue specific expression. *J Exp Med.* 1994; 180:1741–52.
- Cervino AC, Lakiss S, Sow O, Bellamy R, Beyers N, Hoal-van Helden E, et al. Fine mapping of a putative tuberculosis susceptibility locus on chromosome 15q11-13 in African families. *Hum Mol Genet* 2002; 11:1599-1603.
- Chackerian AA and Behar SM. Susceptibility to *Mycobacterium tuberculosis*: lessons from inbred strains of mice. *Tuberculosis (Edinb)* 2003; 83:279-285.
- Chakraborty AK. Epidemiology of tuberculosis: Current status in India. *Indian J Med Res.* 2004; 120:248-76.
- Chakravarti A, Population genetics- making sense out of sequence. *Nat Genet* 1999; 21:56-60.
- Chang K, Deng S, Lu W, Wang F, Jia S, Li F , et al. Association between *CD209* -336A/G and-871A/G Polymorphisms and Susceptibility of Tuberculosis:A Meta-Analysis. *PLoS One.* 2012; 7(7):e41519.
- Che N, Li S, Gao T, Zhang Z, Han Y, Zhang X , et al. Identification of a novel IRGM promoter single nucleotide polymorphism associated with tuberculosis. *Clin Chim Acta.* 2010; 411: 1645–1649.
- Chen XR, Feng YL, Ma Y, Zhang ZD, Li CY, Wen FQ, et al. Study on the association of two polymorphisms of the vitamin D receptor (VDR) gene with the susceptibility to pulmonary tuberculosis (PTB) in Chinese Tibetans. *Sichuan Da Xue Xue Bao Yi Xue Ban.* 2006; 37: 847–51.
- Chinoy H , Salway F , John S , Fertig N, Tait BD , Oddis CV , et al . Interferon-Gamma and Interleukin-4 Gene Polymorphisms in UK Caucasian Idiopathic Inflammatory Myopathy Patients. *Ann Rheum Dis.* 2001 ; 66:970–3.
- Chitale S, Ehrt S, Kawamura I, *et al.*, Recombinant *Mycobacterium tuberculosis* protein associated with mammalian cell entry. *Cell Microbiol* 2001; 3:247-254
- Choudhry S, Coyle NE, Tang H, Salari K, Lind D, Clark SL. Population stratification confounds genetic association studies among Latinos. *Hum Genet.* 2006; 118:652–664.
- Clarke R, Xu P, Bennett D, Lewington S, Zondervan K, Parish S, Palmer A, et al. International Study of Infarct Survival

- 
- (ISIS) Collaborators. Lymphotoxin alpha gene and risk of myocardial infarction in 6,928 cases and 2,712 controls in the ISIS case-control study. *PLoS Genet.* 2006; 2:107.
- Clementi M and Di Gianantonio E. Genetic susceptibility to infectious diseases. *Reprod Toxicol* 2006; 21:345-349.
- Cole ST, Brosch R, Parkhill J, Garnier T, Churcher C, Harris D, et al. Deciphering the biology of *Mycobacterium tuberculosis* from the complete genome sequence. *Nature* 1998; 393:537-544.
- Collins FS, Guyer MS, Charkravarti A. Variations on a Theme: Cataloging Human DNA Sequence Variation. *Science.* 1997; 278:1580-1581.
- Comstock GW. Tuberculosis in twins: a re-analysis of the Proffit survey. *Am Rev Respir Dis.* 1978; 117:621-4.
- Cooke GS, Campbell SJ, Fielding K. Interleukin-8 polymorphism is not associated with pulmonary tuberculosis in the gambia. *J Infect Dis.* 2004; 189:1545-6
- Cooke GS, Campbell SJ, Sillah J, Gustafson P, Bah B, Sirugo G, et al. Polymorphism within the Interferon- $\gamma$  Receptor Complex Is Associated with Pulmonary Tuberculosis. *Am J Respir Crit Care Med.* 2006; 174:339-43.
- Cooke GS, Hill AV. Genetics of susceptibility to human infectious disease. *Nat Rev Genet.* 2001; 2:967-77.
- Coutinho-Silva R, Stahl L, Raymond MN, Jungas T, Verbeke P, Burnstock G, et al. Inhibition of chlamydial infectious activity due to P2X7R-dependent phospholipase D activation. *Immunity.* 2003; 19:403-12.
- Crevel R, Ottenhoff TH, Meer JV. Innate immunity to *Mycobacterium tuberculosis*. *Clin Microbiol Rev.* 2002; 15:294-309.
- Curtis J, Kopanitsa L, Stebbings E, Speirs A, Ignatyeva O, Balabanova Y, et al. Association analysis of the LTA4H gene polymorphisms and pulmonary tuberculosis in 9115 subjects. *Tuberculosis.* 2011; 91:22-5
- Daly AK, Day CP. Candidate gene case-control association studies: advantages and potential pitfalls. *Br J Clin Pharmacol* 2001; 52:489-499.
- Daniel TM. The history of tuberculosis. *Respir Med* 2006; 100:1862-1870.
- Danis VA, Millington M, Hyland VJ, Grennan D. Cytokine production by normal human monocytes: intersubject variation and relationship to an IL-1 receptor antagonist (IL-1Ra) gene polymorphism. *Clin Exp Immunol.* 1995; 99:303-10.
- Davila S, Hibberd ML, Hari Dass R, Wong HE, Sahiratmadja E, Bonnard C, et al. Genetic association and expression studies indicate a role of toll-like receptor 8 in pulmonary tuberculosis. *PLoS Genet.* 2008; 4:10.
- Delgado JC, Baena A, Thim S, Goldfeld AE. Ethnic-specific genetic associations with pulmonary tuberculosis. *J Infect Dis.* 2002; 186:1463-8.
- Deretic V, Levine B. Autophagy, immunity, and microbial adaptations. *Cell Host Microbe.* 2009; 5:527-9.
- Deveci F, Akbulut HH, Turgut T, Muz HM. Changes in Serum Cytokine Levels in Active Tuberculosis With Treatment. *Mediators Inflamm.* 2005; 5: 256-62.
- Devlin B, Roeder K. Genomic control for association studies. *Biometrics.* 1999; 55:997-1004.
- Dewan PK, Grinsdale J, Kawamura LM. Low sensitivity of a whole blood interferon-gamma release assay for detection of active tuberculosis. *Clin Infect Dis.* 2007; 44:69-73.
- Divangahi M, Mostowy S, Coulombe Fo, Kozak R, Guillot Lc, Veyrier F, et al. NOD2-Deficient mice have impaired resistance to *Mycobacterium tuberculosis* infection through defective innate and adaptive immunity. *J Immunol* 2008; 181:7157-65.
- Doffinger R, Dupuis S, Picard C, Fieschi C, Feinberg J, Barcenar-Morales G, et al. Inherited disorders of IL-12 and IFN-gamma-mediated immunity: a molecular genetics update. *Mol Immunol.* 2002; 38:903-9.
- Doherty TM, Andersen P. Vaccines for tuberculosis: novel concepts and recent progress. *Clin Microbiol Rev.* 2005; 18:687-702.
- Drennan MB, Nicolle D, Quesniaux VJ, Jacobs M, Allie N, Mpagi J, et al. Toll-like receptor 2-deficient mice succumb to *Mycobacterium tuberculosis* infection. *Am J Pathol.* 2004; 164:49-57.
- Dubos R, Dubos J. The white plague: tuberculosis, man and society. 1952.
- Ducati RG, Ruffino-Netto A, Basso LA, Santos DS. The resumption of consumption-- a review on tuberculosis. *Mem Inst Oswaldo Cruz* 2006; 101:697-714.
- Duerr RH, Taylor KD, Brant SR *et al.*, A genome-wide association study identifies IL23R as an inflammatory bowel disease gene. *Science* 2006; 314:1461-1463.
- Dunstan SJ, Ho VA, Duc CM, Lanh MN, Phuong CXT, Luxemburger C, et al. Typhoid fever and genetic polymorphisms at the natural resistance-associated macrophage protein. *J Infect Dis.* 2001; 183:1156-60.
- Dupont WD, Plummer WD Jr. Power and sample size calculations for studies involving linear regression. *Control Clin Trials.* 1998; 19:589-601.
- Ehlers S, Benini J, Kutsch S, Endres R, Rietschel ET, Pfeffer K, et al. Fatal granuloma necrosis without exacerbated mycobacterial growth in tumor necrosis factor receptor p55 gene-deficient mice intravenously infected with *Mycobacterium avium*. *Infect. Immun.* 1999; 67:3571-9.
- Engel M, Stössel E, Castiglione K, Schwerdtner N, Wagner M, Bölcskei P, et al. Induction of TNF in human alveolar macrophages as a potential evasion mechanism of virulent *Mycobacterium tuberculosis*. *J Immunol.* 2002; 168:1328-
-

37.

- Evangelou E, Trikalinos TA, Salanti G, Ioannidis JP. Family-based versus unrelated case-control designs for genetic associations. *PLoS Genet* 2006; 2:e123.
- Facchetti F, Vermi W, Fiorentini S, Chilosi M, Caruso A, Duse M, et al. Expression of inducible nitric oxide synthase in human granulomas and histiocytic reactions. *Am J Pathol.* 1999; 154:145–52.
- Fernando SL, Saunders BM, Sluyter R, Skarratt KK, Goldberg H, Marks GB, et al. A Polymorphism in the P2X7 Gene Increases Susceptibility to Extrapulmonary Tuberculosis. *Am J Respir Crit Care Med.* 2007; 175:360–6.
- Fernando SL, Saunders BM, Sluyter R, Skarratt KK, Wiley JS, Britton WJ. Gene dosage determines the negative effects of polymorphic alleles of the P2X7 receptor on adenosine triphosphate-mediated killing of mycobacteria by human macrophages. *J Infect Dis.* 2005; 192:149–155.
- Ferrara G, Losi M, D'Amico R, Roversi P, Piro R, Meacci M, et al. Use in routine clinical practice of two commercial blood tests for diagnosis of infection with *Mycobacterium tuberculosis*: a prospective study. *Lancet.* 2006; 367:1328–34.
- Ferraz JC, Melo FBS, Albuquerque MFPM, Montenegro SML, Abath FGC. Immune factors and immunoregulation in tuberculosis. *Braz J Med Biol Res.* 2006; 39:1387-97.
- Ferwerda B, Kibiki GS, Netea MG, Dolmans WM, van der Ven AJ. The toll-like receptor 4 Asp299Gly variant and tuberculosis susceptibility in HIV infected patients in Tanzania. *Aids.* 2007; 21:1375–7.
- Ferwerda B, Kibiki GS, Netea MG, Dolmans WM, van der Ven AJ. The toll-like receptor 4 Asp299Gly variant and tuberculosis susceptibility in HIV infected patients in Tanzania. *Aids* 2007; 21:1375–7.
- Ferwerda G, Girardin SE, Kullberg BJ, Le BL, de Jong DJ, Langenberg DM, et al. NOD2 and toll-like receptors are nonredundant recognition systems of *Mycobacterium tuberculosis*. *PLoS Pathog.* 2005; 1:279–285.
- Fiorenza G, Rateni L, Farroni MA, Bogue C, Dlugovitzky DG. TNF-alpha, TGF-beta and NO relationship in sera from tuberculosis (TB) patients of different severity. *Immunol Lett.* 2005; 98:45-8.
- Fitness J, Floyd S, Warndorff D K, Sichali L, Malema S, Crampin AC, et al. Large-scale candidate gene study of tuberculosis susceptibility in the Karonga district of northern Malawi. *Am J Trop Med Hyg.* 2004; 71:341–9.
- Fitzgerald D, Sterling TR, Hass DW. *Mycobacterium tuberculosis*. In: Mandell GL, Bennett JE, Dolin R. Principles and practice of infectious diseases 7th ed. Vol2. Philadelphia, PA: Churchill Livingstone Press; p3158.
- Flynn JL, Chan J. Immunology of tuberculosis. *Annu Rev Immunol.* 2001; 19:93–129
- Flynn JL, Goldstein MM, Chan J, Triebold KJ, Pfeffer K, Lowenstein CJ, et al. Tumor necrosis factor alpha is required in the protective immune response against *Mycobacterium tuberculosis* in mice. *Immunity.* 1995; 2:561–72.
- Flynn JL. Lessons from experimental *Mycobacterium tuberculosis* infections. *Microbes Infect* 2006; 8:1179- 88.
- Freidin MB, Rudko AA, Kolokolova OV, Freidin MB, Rudko AA, Kolokolova OV, et al. Association between the 1188 A/C polymorphism in the human IL12B gene and Th1-mediated infectious diseases. *Int J Immunogenet.* 2006; 33:231–2.
- Frieden TR, Sherman LF, Maw KL, Fujiwara PI, Crawford JT, Nivin B, et al. A multi-institutional outbreak of highly drug-resistant tuberculosis: epidemiology and clinical outcomes. *JAMA* 1996; 276:1229-35.
- Frieden TR, Sterling TR, Munsiff SS, Watt CJ, and Dye C. Tuberculosis. *Lancet* 2003; 362:887-99.
- Fulton SA, Cross JV, Toossi ZT, Boom WH. Regulation of interleukin-12 by interleukin-10, transforming growth factor-beta, tumor necrosis factor-alpha, and interferon-gamma in human monocytes infected with *Mycobacterium tuberculosis* H37Ra. *J. Infect. Dis.* 1998; 178:1105–14.
- Gabriel SB, Schaffner SF, Nguyen H, Moore JM, Roy J, Blumenstiel B, et al. The structure of haplotype blocks in the human genome. *Science.* 2002; 296:2225-9.
- Gagneux S, Small PM. Global phylogeography of *Mycobacterium tuberculosis* and implications for tuberculosis product development. *Lancet Infect Dis.* 2007; 7:328–37.
- Gallant CJ, Malik S, Jabado N, Cellier M, Simkin L, Finlay BB, et al. Reduced in vitro functional activity of human *NRAMP1 (SLC11A1)* allele that predisposes to increased risk of pediatric tuberculosis disease. *Genes Immun.* 2007; 8:691–8.
- Gambaro G, Anglani F, D'Angelo A. Association studies of genetic polymorphisms and complex disease. *Lancet* 2000; 355:308-11.
- Gao X, ER martin. Using Allele Sharing Distance for Detecting Human Population Stratification. *Hum Hered.* 2009; 68:182–191.
- Gao YJ, Pei XY, Yang H, Liu F, Jiang XF. A case-control study of the association between VDR gene polymorphism and tuberculosis in Ningxia. *Ningxia Med J.* 2008; 30:673–6.
- Gartner T, Baeten M, Otieno S, Revets H, De Baetselier P, Huygen K. Mucosal prime-boost vaccination for tuberculosis based on TLR triggering OprI lipoprotein from *Pseudomonas aeruginosa* fused to mycolyl-transferase Ag85A. *Immunol Lett.* 2007; 111:26-35.
- Gately MK, Renzetti LM, Magram J, Stern AS, Adorini L, Gubler U, et al. The interleukin-12/ interleukin-12-receptor system: role in normal and pathologic immune responses. *Annu Rev Immunol.* 1998; 16:495–521.

- 
- Geijtenbeek TB, Krooshoop DJ, Bleijs DA, van Vliet SJ, van Duijnhoven GC, Grabovsky V, et al. DC-SIGN-ICAM-2 interaction mediates dendritic cell trafficking. *Nat Immunol.* 2000; 1: 353–7.
- Geijtenbeek TB, Torensma R, van Vliet SJ, van Duijnhoven GC, Adema GJ, van Kooyk Y, et al. Identification of DC-SIGN, a novel dendritic cell-specific ICAM-3 receptor that supports primary immune responses. *Cell.* 2000; 100:575–85.
- Geijtenbeek TB, Van Vliet SJ, Koppel EA, Sanchez-Hernandez M, Vandenbroucke- Grauls CM, Appelmelk B, et al. Mycobacteria target DC-SIGN to suppress dendritic cell function. *J Exp Med* 2003; 197:7–17.
- Geijtenbeek TB. DC-SIGN: escape mechanism for pathogens. *Nat Rev Immunol.* 2003; 3: 697–709.
- Genetics Home reference. 2010; [ghr.nlm.nih.gov/gene/SP110](http://ghr.nlm.nih.gov/gene/SP110)
- Gleeson TD, Decker CF. Treatment of tuberculosis. *Dis Mon* 2006; 52:428–34.
- Gomez LM, Anaya JM, Sierra-Filardi E, Cadena J, Corbi A, Martín J, et al. Analysis of *DC-SIGN (CD209)* functional variants in patients with tuberculosis. *Hum Immunol.* 2006; 67:808–11.
- Gomez LM, Anaya JM, Vilchez JR, Cadena J, Hinojosa R, Velez L, et al. A polymorphism in the inducible nitric oxide synthase gene is associated with tuberculosis. *Tuberculosis.* 2007; 87:288–94.
- Gomez LM, Camargo JF, Castiblanco J, Ruiz-Narváez EA, Cadena J, Anaya JM. Analysis of *IL1B*, *TAP1*, *TAP2* and *IKBL* polymorphisms on susceptibility to tuberculosis. *Tissue Antigens.* 2006; 67: 290–6.
- Gong J, Zhang M, Modlin RL, Linsley PS, Iyer D, Lin Y, et al. Interleukin-10 downregulates *Mycobacterium tuberculosis* induced Th1 responses and CTLA-4 expression. *Infect. Immun.* 1996; 64: 913–8.
- Gordt T and Khamis H. Simple solution to a common statistical problem: interpreting multiple tests. *Clin Ther* 2004; 26:780–786.
- Gordon S. Pattern recognition receptors: doubling up for the innate immune response. *Cell.* 2002; 111:927–30.
- Goren MB, D'Arcy HP, Young MR, Armstrong JA. Prevention of phagosome-lysosome fusion in cultured macrophages by sulfatides of *Mycobacterium tuberculosis*. *Proc Natl Acad Sci. USA* 1976; 73:2510–14.
- Govoni G, Cannone-Hergaux F, Pfeifer CG, Marcus S, Mills S, Hackam D, et al. Functional expression of *Nramp1* in vitro after transfection into murine macrophage line RAW264.7. *Infect Immun.* 1999; 67:2225–32.
- Greenberg SS, Xie J, Kolls J, Mason C, Didier P. Rapid induction of mRNA for nitric oxide synthase II in rat alveolar macrophages by intratracheal administration of *Mycobacterium tuberculosis* and *Mycobacterium avium*. *Proc Soc Exp Biol Med.* 1995; 209:46–53.
- Greenwood CM, Fujiwara TM, Boothroyd LJ *et. al.*, Linkage of tuberculosis to chromosome 2q35 loci, including *NRAMP1*, in a large aboriginal Canadian family. *Am J Hum Genet* 2000; 67:405–416.
- Gros P, Skamene E, Forget A. Genetic control of natural resistance to *Mycobacterium bovis* (BCG) in mice. *J Immunol.* 1981; 127:2417–21.
- Gu BJ, Zhang W, Worthington RA, Sluyter R, Dao-Ung P, Petrou S, et al. Glu-496 to Ala polymorphism leads to loss of function of the human P2X7 receptor. *J Biol Chem.* 2001; 276: 11135–42.
- Gudmundsson J, Sulem P, Manolescu A, Amundadottir LT, Gudbjartsson D, Helgason A, et al. Genome-wide association study identifies a second prostate cancer susceptibility. *Nat Genet.* 2007; 39:631–7.
- Gupta UD and Katoch VM. Animal models of tuberculosis. *Tuberculosis (Edinb)* 2005; 85:277–293.
- Gupta V, Jaiswal V, Behera D, Prasad HK. Disparity in circulating peripheral blood dendritic cell subsets and cytokine profile of pulmonary tuberculosis patients compared with healthy family contacts. *Hum Immunol.* 2010; 71:682–91.
- Gutierrez MC, Brisse S, Brosch R, Fabre M, Omais B, Marmiesse M, et al. Ancient origin and gene mosaicism of the progenitor of *Mycobacterium tuberculosis*. *PLoS Pathog* 2005; 1(1): e5.
- Gutierrez MG, Master SS, Singh SB, Taylor GA, Colombo MI, Deretic V, et al. Autophagy is a defense mechanism inhibiting BCG and *Mycobacterium tuberculosis* survival in infected macrophages. *Cell.* 2004; 119:753–66.
- Hackam DJ, Rostein OD, Zhang W, Gruenheid S, Gros P, Grinstein S. Host resistance to intracellular infection: mutation of natural resistance-associated macrophage protein 1 (*Nramp1*) impairs phagosomal acidification. *J Exp Med.* 1998; 188:351–64.
- Haiman CA, Patterson N, Freedman ML, Myers SR, Pike MC, Waliszewska A, et al. Multiple regions within 8q24 independently affect risk for prostate cancer. *Nat Genet.* 2007; 39:638–44.
- Hampe J, Franke A, Rosenstiel P et al. A genome-wide association scan of nonsynonymous SNPs identifies a susceptibility variant for Crohn disease in *ATG16L1*. *Nat Genet* 2007; 39:207–211.
- Handzel ZT, Barak V, Altman Y, Bibi H, Lidgi M, Iancovici-Kidon M, et al. Increased Th1 and Th2 type cytokine production in patients with active tuberculosis. *Isr Med Assoc J.* 2007; 9:479–83.
- Harpending HC, Batzer MA, Gurven M, Jorde LB, Rogers AR, Sherry ST, et al. Genetic traces of ancient demography. *Proc. Natl. Acad. Sci. USA,* 1998; 95:1961–1967.
- Hasan Z, Jamil B, Khan J, Ali R, Khan MA, Nasir N, et al. Relationship between circulating levels of IFN-gamma, IL-10, CXCL9 and CCL2 in pulmonary and extrapulmonary tuberculosis is dependent on disease severity. *Scand J Immunol.* 2009; 69:259–67.
-

- Haukim N, Bidwell JL, Smith AJP, Keen LJ, Gallagher G, Kimberly R, et al. Cytokine gene polymorphism in human disease: on-line databases. *Gen Immun.* 2002; 3:313-30.
- Healy DG. Case-control studies in the genomic era: a clinician's guide. *Lancet Neurol* 2006; 5:701-707.
- Heiman GA, Hodge SE, Prakash G, Zhang J, Greenberg DA. Effect of Population Stratification on Case-Control Association Studies. *Hum Hered.* 2004; 58:30-39.
- Henderson RA, Watkins SC, Flynn JL. Activation of human dendritic cells following infection with *Mycobacterium tuberculosis*. *J Immunol.* 1997; 159: 635-43.
- Herzog H. History of tuberculosis. *Respiration* 1998; 65:5-15.
- Hill AV. The immunogenetics of human infectious diseases. *Annu Rev Immunol.* 1998; 16:593-617.
- Hirsch CS, Toossi Z, Othieno C, Johnson JL, Schwander SK, Robertson S, et al. Depressed T-cell interferon-gamma responses in pulmonary tuberculosis: analysis of underlying mechanisms and modulation with therapy. *J Infect Dis.* 1999; 180: 2069-73.
- Hirschhorn JN and Daly MJ. Genome-wide association studies for common diseases and complex traits. *Nat Rev Genet* 2005; 6:95-108.
- Hirsh AE, Tsolaki AG, DeRiemer K, Feldman MW, Small PM. Stable association between strains of *Mycobacterium tuberculosis* and their human host populations. *Proc Natl Acad Sci.* 2004; 101:4871-6.
- Hoal EG, Lewis LA, Jamieson SE, Tanzer F, Rossouw M, Victor T, et al. SLC11A1 (NRAMP1) but not SLC11A2 (NRAMP2) polymorphisms are associated with susceptibility to tuberculosis in a high incidence community in South Africa. *Int. J. Tuberc Lung Dis.* 2004; 8:1464-71.
- Humphreys BD, Rice J, Kertesz SB, Dubyak GR. Stress activated protein kinase/JNK activation and apoptotic induction by the macrophage P2X7 nucleotide receptor. *J Biol Chem.* 2000; 275:26792-98.
- Hurme M, Santilla S. IL-1receptor antagonist (IL-1Ra) plasma levels are co-ordinately regulated by both IL-1Ra and IL-1  $\alpha$  genes. *Eur J Immunol.* 1998; 28: 2598-602.
- Indian Genome Variation Consortium. Genetic landscape of the people of India: a canvas for disease gene exploration. *J Genet.* 2008 ;87: 3-20.
- Intemann CD, Thye T, Niemann S, Browne EN, Amanua Chinbuah M, Enimil A, et al. Autophagy gene variant IRGM -261T contributes to protection from tuberculosis caused by *Mycobacterium tuberculosis* but not by *M. africanum* strains. *PLoS Pathog.* 2009; 5: e1000577.
- Jacobs M, Togbe D, Fremont C, Samarina A, Allie N, Botha T, et al. Tumor necrosis factor is critical to control tuberculosis infection. *Microbes Infect.* 2007; 5: 623-8.
- Jamieson SE, Miller EN, Black GF, Peacock CS, Cordell HJ, Howson JM. Evidence for a cluster of genes on chromosome 17q11-q21 controlling susceptibility to tuberculosis and leprosy in Brazilians. *Genes. Immun.* 2004; 5:46-57.
- Johnson R, Streicher EM, Louw GE, Warren RM, van Helden PD, and Victor TC. Drug resistance in *Mycobacterium tuberculosis*. *Curr Issues Mol Biol* 2006; 8:97-111.
- Juffermans NP, Verbon A, Deventer SJ, Deutekom H, Speelman P, van der Poll T. Tumor necrosis factor and interleukin-1 inhibitors as markers of disease activity of tuberculosis. *Am J Respir Crit Care Med.* 1998; 157:1328-31.
- Juffermans NP, Verbon A, van Deventer SJ, van Deutekom H, Belisle JT, Ellis ME, et al. Elevated chemokine concentrations in sera of human immunodeficiency virus (HIV)-seropositive and HIV-seronegative patients with tuberculosis: a possible role for mycobacterial lipoarabinomannan. *Infect Immun.* 1999; 67:4295-7.
- Kadereit S, Gewert DR, Galabru J, Hovanessian AG, Meurs EF. Molecular cloning of two new interferon-induced, highly related nuclear phosphoproteins. *J Biol Chem.* 1993; 268 : 24432-41.
- Kart L, Buyukoglan H, Tekin IO, Altin R, Senturk Z, Gulmez I, et al. Correlation of serum tumor necrosis factor-alpha, interleukin-4 and soluble interleukin-2 receptor levels with radiologic and clinical manifestations in active pulmonary tuberculosis. *Mediators Inflamm.* 2003; 12:9-14.
- Katti MK. Assessment of serum IL-1, IL-2 and IFN- $\gamma$  levels in untreated pulmonary tuberculosis patients: role in pathogenesis. *Arch Med Res.* 2011; 42:199-201.
- Kaufmann SH and McMichael AJ. Annuling a dangerous liaison: vaccination strategies against AIDS and tuberculosis. *Nat Med* 2005; 11:S33-S44.
- Kaufmann SH, Baumann S, and Nasser EA. Exploiting immunology and molecular genetics for rational vaccine design against tuberculosis. *Int J Tuberc Lung Dis* 2006; 10:1068-1079.
- Kaufmann SH, Baumann S, Nasser EA. Exploiting immunology and molecular genetics for rational vaccine design against tuberculosis. *Int J Tuberc Lung Dis.* 2006; 10:1068-79.
- Kaufmann SH, Schaible UE. 100th anniversary of Robert Koch's Nobel Prize for the discovery of the tubercle bacillus. *Trends Microbiol* 2005; 13:469-475.
- Kaufmann SH. Protection against tuberculosis: cytokines, T cells, and macrophages. *Ann Rheum Dis.* 2002; 61:54-58.
- Keane J, Balcewicz-Sablinska MK, Remold HG, Chupp GL, Meek BB, Fenton MJ, et al. Infection by *Mycobacterium tuberculosis* promotes human alveolar macrophage apoptosis. *Infect Immun.* 1997; 65:298-304.



- 
- Keane J, Gershon S, Wise RP, Mirabile-Levens E, Kasznica J, Schwieterman WD, et al. Tuberculosis associated with infliximab, a tumor necrosis factor alpha-neutralizing agent. *N Engl J Med*. 2001; 345:1098–104.
- Keeratan D, Stephan KS, Bingdong ZHU, Richard N. Van ZYL-Smit and Ying Z. The immunology of tuberculosis: From bench to bedside. *Respirology* 2010; 15:433-50.
- Kim DK, Park GM, Hwang YI, Kim HJ, Han SK, Shim YS, et al. Microarray analysis of gene expression associated with extrapulmonary dissemination of tuberculosis. *Respirology* 2006; 11:557–65.
- Kim YK, Uh Y, Lee NS, Cho MY, Eom M, Kim HY. Whole-blood interferon-gamma release assay for diagnosis of tuberculous lymphadenitis. *Tohoku J Exp Med*. 2011; 224:189–193.
- Kindler V, Sappino AP, Grau GE, Piguet PF, Vassalli P, et al. The inducing role of tumor necrosis factor in the development of bactericidal granulomas during BCG infection. *Cell* 1989; 56:731–40.
- King KY, Lew JD, Ha NP, Lin JS, Ma X, Graviss EA, et al. Polymorphic allele of human IRGM1 is associated with susceptibility to tuberculosis in African Americans. *PLoS One*. 2011; 6:e16317.
- Klein RJ, Zeiss C, Chew EY, Tsai JY, Sackler RS, Haynes C, et al. Complement factor H polymorphism in age-related macular degeneration. *Science* 2005; 308:385-389.
- Koch R. Weitere Mittheilungen über das Tuberkulin. *Dt. Med. Wochenschr* 1891; 17:1189-1192.
- Koch W, Wolferstetter H, Schatke A, Schömig A, Kastrati A. Interleukin 18 gene variation and risk of acute myocardial infarction. *Cytokine*. 2011; 56:786-91.
- Kramnik I, Dietrich WF, Demant P, Bloom BR. Genetic control of resistance to experimental infection with virulent *Mycobacterium tuberculosis*. *Proc Natl Acad Sci*. 2000; 97:8560–65.
- Kroeger KM, Carville KS, Abraham LJ. The -308 tumor necrosis factor- $\alpha$  promoter polymorphism affects transcription. *Mol Immunol*. 1997; 34:391–399
- Kumar A, Ghosh B. A single nucleotide polymorphism (A  $\rightarrow$  G) in intron 3 of IFN gamma gene is associated with asthma. *Genes Immun*. 2008; 9:294-301.
- Kumar D, Nath L, Kamal MA, Varshney A, Jain A, Sarman Singh, et al. Genome-wide analysis of the host intracellular network that regulates survival of *Mycobacterium tuberculosis*. *Cell*. 2010; 140:731–43.
- Kunnath-Velayudhan S, Gennaro ML. Immunodiagnosis of tuberculosis: a dynamic view of biomarker discovery. *Clin Microbiol Rev*. 2011; 792–805.
- Kuo HP, Wang CH. Nitric oxide modulates interleukin-1 $\beta$  and tumor necrosis factor- $\alpha$  synthesis by alveolar macrophages in pulmonary tuberculosis. *Am J Respir Crit Care Med*. 2000; 161:192–9.
- Kusner DJ, Barton JA. ATP stimulates human macrophages to kill intracellular virulent *Mycobacterium tuberculosis* via calcium-dependent phagosome–lysosome fusion. *J Immunol*. 2001; 167:3308–15.
- Kusuhara K, Yamamoto K, Okada K, Mizuno Y, Hara T. Association of IL12RB1 polymorphisms with susceptibility to and severity of tuberculosis in Japanese: a gene-based association analysis of 21 candidate genes. *Int J Immunogenet*. 2007; 34:35–44.
- Ladel CH, Blum C, Dreher A, Reifenberg K, Kaufmann SH. Protective role of gamma/delta T cells and alpha/beta T cells in tuberculosis. *Eur J Immunol*. 1995; 25: 2877–81.
- Lander ES., Initial sequencing and analysis of the human genome. *Nature* 2001; 409:860-921.
- Lander ES, The New Genomics: Global Views of Biology. *Science* 1996; 274:536-539.
- Law AJ, Lipska BK, Weickert CS, Hyde TM, Straub RE, Hashimoto R, et al. Neuregulin 1 transcripts are differentially expressed in schizophrenia and regulated by 50 SNPs associated with the disease. *Proc Natl Acad Sci USA* 2006; 103:6747–52.
- Law K, Weiden M, Harkin T, Tchou-Wong K, Chi C, Rom WN. Increased release of interleukin-1  $\beta$ , interleukin-6, and tumor necrosis factor- $\alpha$  by bronchoalveolar cells lavaged from involved sites in pulmonary tuberculosis. *Am J Respir Crit Care Med*. 1996; 153:799–804.
- Leandro AC, Rocha MA, Cardoso CS, Bonecini-Almeida MG. Genetic polymorphisms in vitamin D receptor, vitamin D-binding protein, Toll-like receptor 2, nitric oxide synthase 2, and interferon-gamma genes and its association with susceptibility to tuberculosis. *Braz J Med Biol Res*. 2009; 42:312-22.
- Lee HW, Lee HS, Kim DK, Ko DS, Han SK, Shim YS, et al. Lack of an association between interleukin-12 receptor beta 1 polymorphisms and tuberculosis in Koreans. *Respiration*. 2005; 72:365–8.
- Lesley R., Ramakrishnan L. Insights into early mycobacterial pathogenesis from the zebrafish. *Curr Opin Microbiol*. 2008; 11:277–83.
- Levenstien MA, Ott J, Gordon D. Are molecular haplotypes worth the time and expense? A cost-effective method for applying molecular haplotypes. *PLoS Genet* 2006; 2:e127.
- Levesque MC, Hobbs MR, Anstey NM, Vaughn TN, Chancellor JA, Pole A, et al. Nitric oxide synthase type 2 promoter polymorphisms, nitric oxide production, and disease severity in Tanzanian children with malaria. *J Infect Dis*. 1999; 180:1994–2002.
- Lewis SJ, Baker I, Davey Smith G. Meta-analysis of vitamin D receptor polymorphisms and pulmonary tuberculosis risk.
-

- Int J Tuberc Lung Dis. 2005; 9:1174–77.
- Li CM, Campbell SJ, Kumararatne DS, Bellamy R, Ruwende C, McAdam KP, et al. Association of a polymorphism in the P2X7 gene with tuberculosis in a Gambian population. *J Infect Dis*. 2002; 186:1458–62.
- Li HT, Zhang TT, Zhou YQ, Huang QH, Huang J. SLC11A1 (formerly NRAMP1) gene polymorphisms and tuberculosis susceptibility: a metaanalysis. *Int J Tuberc Lung Dis*. 2006; 10:3–12.
- Liaw YS, Tsai-Wu JJ, Wu CH, Hung CC, Lee CN, Yang PC, et al. Variations in the NRAMP1 gene and susceptibility of tuberculosis in Taiwanese. *Int J Tuberc Lung Dis*. 2002; 6:454–60.
- Lipsitch M, Sousa AO. Histo intensity of natural selection for resistance to tuberculosis *Genetics*. 2002; 161:1599-1607.
- Liu H, Hwangbo Y, Holte S, Lee J, Wang C, Kaupp N, et al. Analysis of genetic polymorphisms in CCR5, CCR2, stromal cell-derived factor-1, RANTES, and dendritic cell-specific intercellular adhesion molecule-3-grabbing nonintegrin in seronegative individuals repeatedly exposed to HIV-1. *J Infect Dis*. 2004; 190:1055–58.
- Liu PT, Stenger S, Li H, Wenzel L, Tan BH, Krutzik SR, et al. Toll-like receptor triggering of a vitamin D-mediated human antimicrobial response. *Science* 2006; 311:1770-73.
- Liu W, Cao WC, Zhang CY, Wu XM, Habbema JD, Zhao QM, et al. VDR and NRAMP1 gene polymorphisms in susceptibility to pulmonary tuberculosis among the Chinese Han population: a case-control study. *Int J Tuberc Lung Dis*. 2004; 8:428–34.
- Lohmueller KE, Pearce CL, Pike M, Lander ES, Hirschhorn JN. Meta-analysis of genetic association studies supports a contribution of common variants to susceptibility to common disease. *Nat Genet* 2003; 33:177-182.
- Lombard Z, Dalton DL, Venter PA, Williams RC, Bornman L. Association of HLA-DR, -DQ, and vitamin D receptor alleles and haplotypes with tuberculosis in the Venda of South Africa. *Hum Immunol*. 2006; 67:643–54.
- Lopez-Maderuelo D, Arnalich F, Serantes R, Gonzalez A, Codoceo R, Madero R, et al. Interferon- $\gamma$  and Interleukin-10 gene polymorphism in pulmonary tuberculosis. *Am J Respir Crit Care Med*. 2003; 167:970-75.
- Lowenstein CJ, Snyder SH. Nitric oxide, a novel biologic messenger. *Cell*. 1992; 70:705–7.
- Lu R, Dou X, Gao X, Zhang J, Ni J, Guo L. A functional polymorphism of lymphotoxin alpha (LTA) gene rs909253 is associated with gastric cancer risk in an Asian population. *Cancer Epidemiol*. 2012 Jun 28. [Epub ahead of print]
- Lucey DR, Clerici M, Shearer GM. Type 1 and type 2 cytokine dysregulation in human infectious, neoplastic, and inflammatory diseases. *Clin Microbiol Rev*. 1996; 9:532–62.
- Ma J, Amos CI. Theoretical formulation of principal components analysis to detect and correct for population stratification. *PLoS One*. 2010; 17;5(9). pii: e12510.
- Ma X, Reich RA, Wright JA, Tooker HR, Teeter LD, Musser JM, et al. Association between interleukin-8 gene alleles and human susceptibility to tuberculosis disease. *J Infect Dis*. 2003; 188:349–55.
- MacMicking JD, North RJ, LaCourse R, Mudgett JS, Shah SK, Nathan CF. Identification of nitric oxide synthase as a protective locus against tuberculosis. *Proc Natl Acad Sci USA* 1997; 94:5243–8.
- MacMicking JD, Taylor GA, McKinney JD. Immune Control of Tuberculosis by IFN-Inducible LRG-47. *Science*. 2003; 302:654–59.
- Malik S, Abel L, Tooker H, Audrey Poon, Leah Simkin, Manon Girard, et al. Alleles of the NRAMP1 gene are risk factors for pediatric tuberculosis disease. *Proc Natl Acad Sci USA* 2005; 102:12183–8.
- Manitchotpisit B, Kunachak S, Kulapraditharom B, Sura T. Combined use of fine needle aspiration cytology and polymerase chain reaction in the diagnosis of cervical tuberculous lymphadenitis. *J Med Assoc Thai*. 1999; 82:363–8.
- Marie C, Losser MR, Fitting C, Kermarrec N, Payen D, Cavaillon JM. Cytokines and soluble cytokine receptors in pleural effusions from septic and nonseptic patients. *Am J Respir Crit Care Med*. 1997; 156:1515-22.
- Martin MP, Lederman MM, Hutcheson HB, Goedert JJ, Nelson GW, van Kooyk Y, et al. Association of DC-SIGN promoter polymorphism with increased risk for parenteral, but not mucosal, acquisition of human immunodeficiency virus type 1 infection. *J. Virol*. 2004; 78:14053–56.
- Massey DC, Parkes M. Genome-wide association scanning highlights two autophagy genes, ATG16L1 and IRGM, as being significantly associated with Crohn's disease. *Autophagy*. 2007; 3:649–51.
- Mathema B, Kurepina NE, Bifani PJ, and Kreiswirth BN. Molecular epidemiology of tuberculosis: current insights. *Clin Microbiol Rev* 2006; 19:658-685.
- Matsuyama W, Hashiguchi T, Matsumuro K, Iwami F, Hirostu Y, Kawabata M, et al. Increased serum level of vascular endothelial growth factor in pulmonary tuberculosis. *Am J Respir Crit Care Med*. 2000; 162:1120-2.
- Mazurek GH, Weis SE, Moonan PK, Daley CL, Bernardo J, Lardizabal AA, et al. Prospective comparison of the tuberculin skin test and 2 whole-blood interferon-gamma release assays in persons with suspected tuberculosis. *Clin Infect Dis*. 2007; 45:837–45.
- McCarroll SA, Huett A, Kuballa P, Chilewski SD, Landry A, Goyette P, et al. Deletion polymorphism upstream of IRGM associated with altered IRGM expression and Crohn's disease. *Nat Genet*. 2008; 40:1107–12.
- McDermid JM, Prentice AM. Iron and infection: effects of host iron status and the iron-regulatory genes haptoglobin

- and *NRAMP1 (SLC11A1)* on hostpathogen interactions in tuberculosis and HIV. *Clin Sci (Lond)* 2006; 110:503–24.
- McShane H. Susceptibility to tuberculosis - the importance of the pathogen as well as the host. *Clinical and Experimental Immunology* 2003; 133:201.
- Means TK, Jones BW, Schromm AB, Shurtleff BA, Smith JA, Keane J , et al. Differential effects of a Toll-like receptor antagonist on *Mycobacterium tuberculosis* induced macrophage responses. *J Immunol* 2001; 166:4074–82.
- Mege JL , Meghari S, Honstette A , Capo C, Raoult D. The two faces of interleukin 10 in human infectious diseases. *Lancet Infect Dis.* 2006; 557–69.
- Migliori, GB, De Iaco, G, Besozzi, G, Centis R, Cirillo DM. *Eurosurveillance.* 2007, 12, 3194 .
- Miller EN, Jamieson SE, Joberty C, Fakiola M, Hudson D, Peacock CS, et al. Genome-wide scans for leprosy and tuberculosis susceptibility genes in Brazilians. *Genes Immun* 2004; 5:63-67.
- Mohan AK, Coté TR, Block JA, Manadan AM, Siegel JN, Braun MM , et al. Tuberculosis following the use of etanercept, a tumor necrosis factor inhibitor. *Clin Infect Dis.* 2004; 39:295–9.
- Mohapatra PR, Janmeja AK. Tuberculous Lymphadenitis. *J Assoc Physicians India.* 2009; 57:585 –90.
- Mokrousov I, Sapozhnikova N, Narvskaya O. *Mycobacterium tuberculosis* co-existence with humans: making an imprint on the macrophage P2X(7) receptor gene? *J Med Microbiol.* 2008; 57:581–84.
- Möller M, de Wit E, Hoal EG. Past, present and future directions in human genetic susceptibility to tuberculosis. *FEMS Immunol Med Microbiol.* 2010; 58:3-26.
- Möller M, Hoal EG. Current findings, challenges and novel approaches in human genetic susceptibility to tuberculosis. *Tuberculosis.* 2010; 90:71-83.
- Möller M, Nebel A, Helden PDV, Schreiber S, Hoal EG. Analysis of eight genes modulating interferon gamma and human genetic susceptibility to tuberculosis: a case-control association study. *BMC Infectious Diseases.* 2010; 10:154.
- Moncada S. Nitric oxide in the vasculature: physiology and pathophysiology. *Ann N Y Acad Sci.* 1997; 811:60–7.
- Morahan G, Kaur G, SinghM, , Rappthap CC, Kumar N, Katoch K , et al. Association of variants in the IL12B gene with leprosy and tuberculosis. *Tissue Antigens.* 2007; 69:234–6.
- Morosini M, Meloni F, Bianco AM, Paschetto E, Uccelli M, Pozzi E , et al. The assessment of IFN-gamma and its regulatory cytokines in the plasma and bronchoalveolar lavage fluid of patients with active pulmonary tuberculosis, *Int J Tuberc Lung Dis.* 2003; 7:994-1000.
- Morrison NA, Qi J C, Tokita A, Kelly PJ, Crofts L, Nguyen TV , et al. Prediction of bone density from vitamin D receptor alleles. *Nature.* 1994; 367:284–87.
- Mostowy S, Behr MA. Comparative genomics in the fight against tuberculosis: diagnostics, epidemiology, and BCG vaccination. *Am J Pharmacogenomics* 2002; 2:189-196.
- Motsinger AA, Haas DW, Hulgan T, Ritchie MD. Human genomic association studies: a primer for the infectious diseases specialist. *J Infect Dis.* 2007; 195:1737–44.
- Motsinger-Reif AA, Antas PRZ, Okil NO, Levy S, Holland SM, Sterling T R. Polymorphisms in IL-1 $\beta$ , vitamin D receptor Fok1, and Toll-like receptor 2 are associated with extrapulmonary tuberculosis. *BMC Medical Genetics.* 2010; 11:37.
- Motulsky AG. Metabolic polymorphisms and the role of infectious diseases in human evolution. *Hum Biol* 1960; 32:28-62.
- Motulsky H. *Intuitive Biostatistics.* Oxford University Press, Inc, 1995.
- Mueller JC. Linkage disequilibrium for different scales and applications. *Brief Bioinform* 2004; 5:355-364.
- Nathan CF, Murray HW, Wiebe ME, Rubin BY. Identification of interferon- $\gamma$  as the lymphokine that activates human macrophage oxidative metabolism and anti-microbial activity. *J Exp Med.* 1983; 58:670–9.
- Nathan CF, Murray HW, Wiebe ME, Rubin BY. Identification of interferon- $\gamma$  as the lymphokine that activates human macrophage oxidative metabolism and anti-microbial activity. *J Exp Med.* 1983; 58:670–89.
- Nations JA, Lazarus AA, and Walsh TE. Drug-resistant tuberculosis. *Dis Mon* 2006; 52:435-440.
- Nava-Aguilera E, Andersson N, Harris E, Mitchell S, Hamel C, Shea B , et al. Risk factors associated with recent transmission of tuberculosis: systematic review and meta-analysis. *Int J Tuberc Lung Dis.* 2009; 13:17–26.
- Nedwin GE, Naylor SL, Sakaguchi AY, Smith D, Jarrett-Nedwin J, Pennica D, , et al. Human lymphotoxin and tumor necrosis factor genes: structure, homology and chromosomal localization. *Nucleic Acids Res.* 1985; 13:6361-73.
- Newport MJ, Allen A, Awomoyi AA, Dunstan SJ, McKinney E, Marchant A , et al. The toll-like receptor 4 Asp299Gly variant: no influence on LPS responsiveness or susceptibility to pulmonary tuberculosis in the Gambia. *Tuberculosis.* 2004; 84:347–52.
- Newport MJ, Nejentsev S. Genetics of susceptibility to tuberculosis in humans. *Monaldi Arch Chest Dis.* 2004 ; 61:102–11.
- Newton CR, Graham A, Heptinstall LE, Powell SJ, Summers C, Kalsheker N , et al. Analysis of any point mutation in DNA. The amplification refractory mutation system (ARMS). *Nucleic Acids Res* 1989; 17:2503-16.
- Niño-Moreno P, Portales-Pérez D, Hernández-Castro B, Portales-Cervantes L, Flores-Meraz V, Baranda L , et al. P2X7

- and NRAMP1/SLC11 A1 gene polymorphisms in Mexican mestizo patients with pulmonary tuberculosis. *Clin Exp Immunol.* 2007; 148:469–477.
- Nicholson S, Bonecini-Almeida Mda G, Lapa e Silva JR, C Nathan, Q W Xie, R Mumford , et al. Inducible nitric oxide synthase in pulmonary alveolar macrophages from patients with tuberculosis. *J Exp Med.* 1996; 183:2293–302.
- Nicke A, Baumert HG, Rettinger J, Eichele A, Lambrecht G, Mutschler E, Schmalzing G. P2X1 and P2X3 receptors form stable trimers: a novel structural motif of ligand-gated ion channels. *EMBO J* 1998; 17:3016–28.
- Nicodemus KK, Liu W, Chase GA, Tsai YY, and Fallin MD. Comparison of type I error for multiple test corrections in large single-nucleotide polymorphism studies using principal components versus haplotype blocking algorithms. *BMC Genet* 2005; 6:S1-S78.
- Nnoaham KE, Clarke A. Low serum vitamin D levels and tuberculosis: a systematic review and meta-analysis. *Int J Epidemiol.* 2008; 37:113–9.
- North RJ, Jung YJ. Immunity to tuberculosis. *Annu Rev Immunol.* 2004; 22:599–623.
- Nyholt DR. A simple correction for multiple testing for single-nucleotide polymorphisms in linkage disequilibrium with each other. *Am J Hum Genet* 2004; 74:765-769.
- Ogarkov O, Mokrousov I, Sinkov V, Zhdanova S, Antipina S, Savilov E , et al. 'Lethal' combination of Mycobacterium tuberculosis Beijing genotype and human CD209–336G allele in Russian male population. *Infect Genet Evo.* 2012; 12:732–6.
- Ogus AC, Yoldas B, Ozdemir T, Uguz A, Olcen S, Keser I , et al. The Arg753Gln polymorphism of the human Toll-like receptor 2 gene in tuberculosis disease. *Eur Respir J.* 2004; 23:219-23.
- Okayama N, Fujimura K, Nakamura J, Suehiro Y, Hamanaka Y, Hinoda Y. Evaluation of a new efficient procedure for single-nucleotide polymorphism genotyping: tetra-primer amplification refractory mutation system-polymerase chain reaction. *Clin Chem Lab Med.* 2004; 42:13-26.
- Olesen R, Wejse C, Velez D R, Bisseye C, Sodemann M, Aaby P , et al. DC-SIGN (CD209), pentraxin 3 and vitamin D receptor gene variants associate with pulmonary tuberculosis risk in West Africans. *Genes Immun.* 2007; 8:456–67.
- Oral HB, Budak F, Uzaslan EK, Baştürk B, Bekar A, Akalin H , et al. Interleukin-10 (IL-10) gene polymorphism as a potential host susceptibility factor in tuberculosis. *Cytokine* 2006; 35:143–7.
- Orme IM, Cooper AM. Cytokine/chemokine cascades in immunity to tuberculosis. *Immunol Today.* 1999; 20:307–312.
- Ozaki K, Tanaka T. Genome-wide association study to identify single nucleotide polymorphisms conferring risk of myocardial infarction. *Methods Mol Med* 2006; 128:173-180.
- Ozaki K, Inoue K, Sato H et al. Functional variation in LGALS2 confers risk of myocardial infarction and regulates lymphotoxin-alpha secretion in vitro. *Nature* 2004; 429:72-75.
- Ozaki K, Ohnishi Y, Iida A, Sekine A, Yamada R, Tsunoda T, et al. Functional SNPs in the lymphotoxin-alpha gene that are associated with susceptibility to myocardial infarction. *Nat Genet* 2002; 32:650-654.
- Pace JL, Russell SW, Torres BA, Johnson HM, Gray PW. Recombinant mouse  $\gamma$ -interferon induces the priming step in macrophage activation for tumor cell killing. *J Immunol.* 1983; 130:2011–3.
- Parkes M, Barrett JC, Prescott NJ, Tremelling M, Anderson CA, Fisher SA , et al. Sequence variants in the autophagy gene IRGM and multiple other replicating loci contribute to Crohn's disease susceptibility. *Nat Genet.* 2007; 39:830–2.
- Pathak SK, Basu S, Basu KK, Banerjee A, Pathak S, Bhattacharyya A, et al. Direct extracellular interaction between the early secreted antigen ESAT-6 of Mycobacterium tuberculosis and TLR2 inhibits TLR signaling in macrophages. *Nat Immunol.* 2007; 8:610-8.
- Patil N, Berno AJ, Hinds DA et al. Blocks of limited haplotype diversity revealed by high-resolution scanning of human chromosome 21. *Science* 2001; 294:1719-23.
- Pease AS. Some remarks on the diagnosis and treatment of tuberculosis in antiquity. *Isis* 1940; 3:380-93.
- Perrey C, Pravica V, Sinnott PJ, Hutchison IV. Genotyping for polymorphisms in interferon-gamma, interleukin-10, transforming growth factor-beta 1 and tumour necrosis factor-alpha genes: a technical report. *Transpl. Immunol.* 1998; 6:193–7.
- Phillips MS, Lawrence R, Sachidanandam R *et al.*, Chromosome-wide distribution of haplotype blocks and the role of recombination hot spots. *Nat Genet* 2003; 33:382-387.
- Pieters, J, Garfield J. Hijacking the host: Survival of pathogenic mycobacterial inside macrophages. *Trends Microbiol.* 2002; 10:142-6.
- Png E, Alisjahbana B, Sahiratmadja E, A genome wide association study of pulmonary tuberculosis susceptibility in Indonesians. *BMC Med Genet.* 2012; 13:13:5.
- Polesky A, Grove W, Bhatia G. Peripheral tuberculous lymphadenitis: epidemiology, diagnosis, treatment and outcome. *Medicine. (Baltimore)* 2005; 84: 350–62.
- Poort SR, Rosendaal FR, Reitsma PH, Bertina RM. A common genetic variation in the 3'-untranslated region of the prothrombin gene is associated with elevated plasma prothrombin levels and an increase in venous thrombosis. *Blood.* 1996; 88:3698–703.

- 
- Poveda F, Camacho J, Arnalich F, Codoceo R, Arco A, Hernández PM. Circulating cytokine concentrations in tuberculosis and other chronic bacterial infections. *Infection*. 1999; 27:272-4.
- Powrie F, Coffman RL. Inhibition of cell-mediated immunity by IL-4 and IL-10. *Res Immunol*. 1993; 144:639-43.
- Price AL, Patterson J, Plenge RM, Weinblatt ME, Shadick NA, Reich D. Principal components analysis corrects for stratification in genome-wide association studies. *Nat Genet*. 2006; 38:904-9.
- Pritchard JK, Seielstad MT, Perez-Lezaun A, Feldman MW. Population growth of human Y chromosomes: a study of Y chromosome microsatellites. *Mol Biol Evol* 1999; 16:1791-1798.
- Rajalingam R, Mehra NK, Pande J, Jain RC, Singla R. Correlation of serum interleukin-2 receptor alpha levels with clinical manifestations in pulmonary tuberculosis. *Tuber Lung Dis*. 1996; 77:374-9.
- Rangaka MX, Wilkinson KA, Glynn JR. Predictive value of interferon- $\gamma$  release assays for incident active tuberculosis: a systematic review and meta-analysis. *Lancet Infect Dis* 2012; 12:45-55.
- Raviglione M, Snider D, Kochi A. Global epidemiology of tuberculosis: morbidity and mortality of a worldwide epidemic. *JAMA* 1995; 273:220-6.
- Reich DE and Lander ES. On the allelic spectrum of human disease. *Trends Genet* 2001; 17:502-510.
- Reichel H, Koefler HP, Norman AW. The role of the vitamin D endocrine system in health and disease. *N Engl J Med*. 1989; 320:980-91.
- Reiling N, Hölscher C, Fehrenbach A, Kröger S, Kirschning CJ, Goyert S, et al. Cutting edge: toll-like receptor (TLR)2- and TLR4-mediated pathogen recognition in resistance to airborne infection with *Mycobacterium tuberculosis*. *J Immuno*. 2002; 169:3480-4.
- Remus N, El Baghdadi J, Fieschi C, Feinberg J, Quintin T, Chentoufi M, et al. Association of IL12RB1 polymorphisms with pulmonary tuberculosis in adults in Morocco. *J Infect Dis*. 2004; 190:580-87.
- Rich EA, Torres M, Sada E, Finegan CK, Hamilton BD, Toossi Z. *Mycobacterium tuberculosis* (MTB)-stimulated production of nitric oxide by human alveolar macrophages and relationship of nitric oxide production to growth inhibition of MTB. *Tuber Lung Dis*. 1997; 78:247-55.
- Risch N, Merikangas K. The future of genetic studies of complex human diseases. *Science* 1996; 273:1516-1517.
- Roach DR, Briscoe H, Saunders B, France MP, Riminton S, Britto WJ. Secreted Lymphotoxin-a Is Essential for the Control of an Intracellular Bacterial Infection. *J Exp Med*. 2001; 193:239-246.
- Roach TI, Barton CH, Chatterjee D, Blackwell JM. Macrophage activation: lipoarabinomannan from avirulent and virulent strains of *Mycobacterium tuberculosis* differentially induces the early genes c-fos, KC, JE, and tumor necrosis factor-alpha. *J Immunol*. 1993; 150:1886-96.
- Roberts M, Mock BA, and Blackwell JM. Mapping of genes controlling *Leishmania major* infection in CXS recombinant inbred mice. *Eur J Immunogenet* 1993; 20:349-62.
- Rockett KA, Brookes R, Udalova I, Vidal V, Hill AV, Kwiatkowski D. 1,25-Dihydroxyvitamin D3 induces nitric oxide synthase and suppresses growth of *Mycobacterium tuberculosis* in a human macrophage-like cell line. *Infect Immun*. 1998; 66:5314-21.
- Rodrigues R, Resende TC, Johnson JL. Sputum cytokine levels in patients with pulmonary tuberculosis as early markers of mycobacterial clearance. *Clin Diagn Lab Immunol*. 2002; 9:818-23.
- Rom WN and Garay SM. *Tuberculosis*. Philadelphia: Lippincott Williams & Wilkins, 2004.
- Rook GA, Steele J, Ainsworth M and Leveton C. A direct effect of glucocorticoid hormones on the ability of human and murine macrophages to control the growth of *Mycobacterium tuberculosis*. *Eur J Respir Dis* 1987; 71:286-291.
- Rosenberg PS, Che A, and Chen BE. Multiple hypothesis testing strategies for genetic case-control association studies. *Stat Med* 2006; 25:3134-3149.
- Rosenwasser LJ, Klemm DJ, Dresback JK, Inamura H, Mascali JJ, Klinnert M, et al. Promoter polymorphisms in the chromosome 5 gene cluster in asthma and atopy. *Clin Exp Allergy*. 1995; 25:74-8; discussion95-6.
- Roth DE, Soto G, Arenas F, Bautista CT, Ortiz J, Rodriguez R, et al. Association between vitamin D receptor gene polymorphisms and response to treatment of pulmonary tuberculosis. *J Infect Dis*. 2004; 190:920-7.
- Roth S, Carr M, Springer T. C-C chemokines, but not the C-X-C chemokines, interleukin-8 and interferon-gamma inducible protein-10, stimulate transendothelial chemotaxis of T lymphocytes. *Eur J Immunol*. 1995; 25:3482-8.
- Ryu S, Park YK, Bai GH, Kim SJ, Park SN, Kang S. 3'UTR polymorphisms in the NRAMP1 gene are associated with susceptibility to tuberculosis in Koreans. *Int J Tuberc Lung Dis*. 2000; 4:577-80.
- Sakuntabhai A, Turbpaiboon C, Casademont I, Chuansumrit A, Lowhnoo T, Kajaste-Rudnitski A, et al. A variant in the CD209 promoter is associated with severity of dengue disease. *Nat Genet*. 2005; 37:507-13.
- Sambasivan V, Murthy KJ, Reddy R, Vijayalakshmi V, Hasan Q. P2X7 gene polymorphisms and risk assessment for pulmonary tuberculosis in Asian Indians. *Dis Markers*. 2010; 28:43-48.
- Schaid DJ. Transmission disequilibrium, family controls, and great expectations. *Am J Hum Genet* 1998; 63:935-41.
- Schultz RM, Kleinschmidt W. Functional identity between murine interferon and macrophage activating factor. *Nature*. 1983; 305:239-40.
-

- Schwandner R, Dziarski R, Wesche H, Rothe M, Kirschning CJ. Peptidoglycan- and lipoteichoic acid-induced cell activation is mediated by toll-like receptor 2. *J Biol Chem.* 1999; 274:17406–9.
- Scola L, Crivello A, Marino V, Gioia V, Serauto A, Candore G, et al. IL-10 and TNF-alpha polymorphisms in a sample of Sicilian patients affected by tuberculosis: implication for ageing and life span expectancy. *Mech Ageing Dev.* 2003; 124:569–72.
- Sebastiani P, Abad MM, Alparagu G, Ramoni MF. Robust transmission/disequilibrium test for incomplete family genotypes. *Genetics* 2004; 168:2329-337.
- Selvaraj P, Alagarasu K, Harishankar M, Vidyarani M, Narayanan PR. Regulatory region polymorphisms of vitamin D receptor gene in pulmonary tuberculosis patients and normal healthy subjects of South India. *Int J Immunogenet.* 2008; 35:251–4.
- Selvaraj P, Alagarasu K, Harishankar M, Vidyarani M, Nisha RD, Narayanan PR. Cytokine gene polymorphisms and cytokine levels in pulmonary tuberculosis. *Cytokine.* 2008; 43:26–33.
- Selvaraj P, Alagarasu K, Swaminathan S, Harishankar M, Narendran G. *CD209* gene polymorphisms in South Indian HIV and HIV-TB patients. *Infect Genet Evol.* 2009; 9:256–62.
- Selvaraj P, Chandra G, Jawahar MS, Rani MV, Rajeshwari DN, Narayanan PR. Regulatory role of vitamin D receptor gene variants of BsmI, ApaI, TaqI, and FokI polymorphisms on macrophage phagocytosis and lymphoproliferative response to *Mycobacterium tuberculosis* antigen in pulmonary tuberculosis. *J Clin Immunol.* 2004; 24:523–32.
- Selvaraj P, Chandra G, Kurian SM, Reetha AM, Charles N, Narayanan PR. *NRAMP1* gene polymorphism in pulmonary and spinal tuberculosis. *Curr Sci.* 2002; 82:4.
- Selvaraj P, Kurian SM, Chandra G, Reetha AM, Charles N, Narayanan PR. Vitamin D receptor gene variants of BsmI, ApaI, TaqI, and FokI polymorphisms in spinal tuberculosis. *Clin Genet.* 2004; 65:73-6.
- Selvaraj P, Kurian SM, Uma H, Reetha AM, Narayanan PR. Influence of non-MHC genes on lymphocyte response to *Mycobacterium tuberculosis* antigens and tuberculin reactive status in pulmonary tuberculosis. *Indian J Med Res.* 2000; 112:86–92.
- Selvaraj P, Prabhu Anand S, Harishankar M, Alagarasu K. Plasma 1,25 dihydroxy vitamin D(3) level and expression of vitamin D receptor and cathelicidin in pulmonary tuberculosis *J Clin Immunol.* 2009; 29:470–478.
- Selvaraj P, Prabhu anand S, Jawahar MS. Promoter polymorphism of IL-8 gene and IL-8 production in pulmonary tuberculosis. *Curr Sci.* 2006;90:952–4.
- Selvaraj P, Vidyarani M, Alagarasu K, Prabhu Anand S, Narayanan P R. Regulatory role of promoter and 3' UTR variants of vitamin D receptor gene on cytokine response in pulmonary tuberculosis. *J Clin Immunol.* 2008; 28:306–13
- Serbina NV, Flynn JL. Early emergence of CD8(+) T cells primed for production of type 1 cytokines in the lungs of *Mycobacterium tuberculosis*-infected mice. *Infect Immun* 1999; 67: 3980–88.
- Sharma PR, Singh S, Jena M, Mishra G, Prakash R, Das PK, et al. Coding and non-coding polymorphisms in VDR gene and susceptibility to pulmonary tuberculosis in tribes, castes and Muslims of Central India. *Infect Genet Evol.* 2011; 11:1456-61.
- Sharma S, Rathored J, Ghosh B, Sharma SK. Genetic polymorphisms in *TNF* genes and tuberculosis in North Indians. *BMC Infectious Diseases.* 2010; 10:165.
- Sharma SK, Mohan A. Extrapulmonary tuberculosis. *Indian J Med Res.* 2004; 120:316-53.
- Shaw MA, Collins A, Peacock CS, Miller EN, Black GF, Sibthorpe D, et al. Evidence that genetic susceptibility to *Mycobacterium tuberculosis* in a Brazilian population is under oligogenic control: linkage study of the candidate genes *NRAMP1* and *TNFA*. *Tuber Lung Dis.* 1997; 78:35–45.
- Shaw TC, Thomas LH, Friedland JS. Regulation of IL-10 secretion after phagocytosis of *Mycobacterium tuberculosis* by human monocytic cells. *Cytokine.* 2000; 12: 483–6.
- Shemon AN, Sluyter R, Fernando SL, Clarke AL, Dao-Ung LP, Skarratt KK, et al. A Thr357 to Ser polymorphism in homozygous and compound heterozygous subjects causes absent or reduced P2X7 function and impairs ATP-induced mycobacterial killing by macrophages. *J Biol Chem.* 2006; 281:2079–86.
- Shin HD, Park BL, Kim LH, Cheong HS, Lee IH, Park SK. Common interleukin 10 polymorphism associated with decreased risk of tuberculosis. *Exp Mol Med.* 2005; 37:128-32.
- Singh SB, Davis AS, Taylor GA, Deretic V. Human IRGM induces autophagy to eliminate intracellular mycobacteria. *Science.* 2006; 313:1438–41.
- Smolonska J, Wijmenga C, Postma DS, Boezen HM. Meta analyses on suspected chronic obstructive pulmonary disease genes: a summary of 20 years' research. *Am J Respir Crit Care Med.* 2009; 180:618-31.
- Soborg C, Andersen AB, Range N, Malenganisho W, Friis H, Magnussen P, et al. Influence of candidate susceptibility genes on tuberculosis in a high endemic region. *Mol Immunol* 2007; 44:2213–20.
- Soilleux EJ, Barten R, Trowsdale J. DC-SIGN; a related gene, DCSIGNR; and CD23 form a cluster on 19p13. *J Immunol.* 2000; 165:2937–42.
- Song KH, Jeon JH, Park WB, Kim SH, Park KU, Kim NJ, et al. Usefulness of the whole-blood interferon-gamma

- 
- release assay for diagnosis of extrapulmonary tuberculosis. *Diagn Microbiol Infect Dis.* 2009; 63:182–7.
- Songane M, Kleinnijenhuis J, Alisjahbana B, Sahiratmadja E, Parwati I, Oosting M, et al. Polymorphisms in autophagy genes and susceptibility to tuberculosis. *PLoS ONE* 2012 7(8): e41618.
- Spielman RS, Ewens WJ. A sibship test for linkage in the presence of association: the sib transmission/disequilibrium test. *Am J Hum Genet* 1998; 62:450–458.
- Spielman RS, McGinnis RE, and Ewens WJ. Transmission test for linkage disequilibrium: the insulin gene region and insulin-dependent diabetes mellitus (IDDM). *Am J Hum Genet* 1993; 52:506–516.
- Stach JL, Gros P, Forget A and Skamene E. Phenotypic expression of genetically controlled natural resistance to *Mycobacterium bovis* (BCG). *J Immunol* 1984; 132:888–92.
- Stead WW, Senner JW, Reddick WT, Lofgren JP. Racial differences in susceptibility to infection by *Mycobacterium tuberculosis*. *N Engl J Med.* 1990; 322:422–7.
- Sternsdorf T, Jensen K, Reich B, Will H. The nuclear dot protein sp100, characterization of domains necessary for dimerization, subcellular localization, and modification by small ubiquitin-like modifiers. *J Biol Chem.* 1999; 274:12555–66.
- Sun F, Flanders WD, Yang Q, and Khoury MJ. Transmission disequilibrium test (TDT) when only one parent is available: the 1-TDT. *Am J Epidemiol* 1999; 150:97–104.
- Surdo PL, Bottomley MJ, Sattler M, Scheffzek K. Crystal structure and nuclear magnetic resonance analyses of the SAND domain from glucocorticoid modulatory element binding protein-1 reveals deoxyribonucleic acid and zinc binding regions. *Mol Endocrinol.* 2003; 17:1283–95.
- Syvänen AC. Accessing genetic variation: genotyping single nucleotide polymorphisms. *Nat Rev Genet* 2001; 2:930–942.
- Szeszko SJ, Healy B, Stevens H, Balabanova Y, Drobniowski F, Todd JA, et al. Resequencing and association analysis of the SP110 gene in adult pulmonary tuberculosis. *Hum Genet* 2007; 121:155–60.
- Tailleux L, Maeda N, Nigou J, Gicquel B, Neyrolles O. How is the phagocyte lectin keyboard played? Master class lesson by *Mycobacterium tuberculosis*. *Trends Microbiol.* 2003; 11:259–263.
- Tang S, Xiao H, Fan Y, Wu F, Zhang Z, Li H, et al. Changes of proinflammatory cytokines and their receptors in serum from patients with pulmonary tuberculosis, *Zhonghua Jie He He Hu Xi Za Zhi.* 2002; 25:325–9.
- The International HapMap Consortium. A haplotype map of the human genome. *Nature* 2005; 437:1299–1320.
- Thuong NT, Hawn TR, Thwaites GE, Chau TT, Lan NT, Quy HT, et al. A polymorphism in human TLR2 is associated with increased susceptibility to tuberculous meningitis. *Genes Immun.* 2007; 8:422–8.
- Thye T, Browne EN, Chinbuah MA, Gyapong J, Osei I, Owusu Dabo E, et al. No associations of human pulmonary tuberculosis with Sp110 variants. *J Med Genet.* 2006; 43.
- Thye T, Owusu-Dabo E, Vannberg FO *et al.*, Common variants at 11p13 are associated with susceptibility to tuberculosis. *Nat Genet* 2012; 44:257–259.
- Tindall EA, Hayes VM, Petersen DC. Inflammatory Genetic Markers of Prostate Cancer Risk. *Cancers.* 2010; 2:1198–220.
- Tobin DM, Ramakrishnan L. Comparative pathogenesis of *Mycobacterium marinum* and *Mycobacterium tuberculosis*. *Cell Microbiol.* 2008; 10:1027e39.
- Tobin DM, Roca FJ, Oh SF, McFarland R, Vickery TW, Ray JP, et al. Host genotype specific therapies can optimize the inflammatory response to mycobacterial infections. *Cell.* 2012; 148:434–46.
- Tobin DM, Vary Jr JC, Ray JP, Walsh GS, Dunstan SJ, Bang ND, et al. The *Ita4h* locus modulates susceptibility to mycobacterial infection in zebrafish and humans. *Cell* 2010; 140:717e30.
- Torrico F, Heremans H, Rivera MT, Marck VE, Billiau A, Carlier Y. Endogenous IFN- $\gamma$  is required for resistance to acute *Trypanosoma cruzi* infection in mice. *J Immunol.* 1991; 146:3626–32.
- Tosh K, Campbell SJ, Fielding K, Sillah J, Bah B, Gustafson P, et al. Variants in the SP110 gene are associated with genetic susceptibility to tuberculosis in West Africa. *Proc Natl Acad Sci. USA* 2006; 103:10364–68.
- Trinchieri G, Pflanz S, Kastelein RA. The IL-12 family of heterodimeric cytokines: new players in the regulation of T cell responses. *Immunity.* 2003; 19:641–4.
- Trinchieri G. Interleukin-12 and the regulation of innate resistance and adaptive immunity. *Nat Rev Immunol.* 2003; 3:133–46.
- Tsao CT, Li L, Hsieh M, Liao S, Chang KS. Soluble TNF-alpha receptor and IL-1 receptor antagonist elevation in BAL in active pulmonary TB. *Eur Respir J.* 1999; 14:490–505.
- Tsao TC, Hong J, Huang C, Yang P, Liao SK, Chang KS. Increased TNF-alpha, IL-1 beta and IL-6 levels in the bronchoalveolar lavage fluid with the upregulation of their mRNA in macrophages lavaged from patients with active pulmonary tuberculosis. *Tuber Lung Dis.* 1999; 79:279–85.
- Tsilidis KK, Helzlsouer KJ, Smith MW, Grinberg V, Hoffman-Bolton J, Clipp SL. Association of common polymorphisms in IL10, and in other genes related to inflammatory response and obesity with colorectal cancer. *Cancer Causes Control.* 2009; 20: 1739–51.
-

- Tso HW, Lau YL, Tam CM, Wong HS, Chiang AK, et al. Associations between IL12B polymorphisms and tuberculosis in the Hong Kong Chinese population. *J Infect Dis*. 2004; 190:913–19.
- Tsuji S, Matsumoto M, Takeuchi O, Akira S, Azuma I, Hayashi A, et al. Maturation of human dendritic cells by cell wall skeleton of *Mycobacterium bovis* bacillus Calmette-Guerin: involvement of toll-like receptors. *Infect Immun*. 2000; 68:6883–90.
- Turgut T, Akbulut H, Deveci F, Kacar C, Muz MH. Serum interleukin-2 and neopterin levels as useful markers for treatment of active pulmonary tuberculosis. *Tohoku J Exp Med*. 2006; 209:321–8.
- Udwadia ZF, Amala RA, Ajbani KK, Rodrigues C. Totally drug-resistant tuberculosis in India. *Clin. Infect*. 2012; 54(4):579–581.
- Underhill DM, Ozinsky A, Smith KD, Aderem A. Toll-like receptor-2 mediates mycobacteria-induced proinflammatory signaling in macrophages. *Proc Natl Acad Sci. USA* 1999; 96:14459–63.
- Valdivielso J M, Fernandez E. Vitamin D receptor polymorphisms and diseases. *Clin Chim Acta*. 2006; 371:1–12.
- Vallinoto ACR, Graça ES, Araújo MS, Azevedo VN, Cayres-Vallinoto I, Machado LF. IFN  $\gamma$  +874T/A polymorphism and cytokine plasma levels are associated with susceptibility to *Mycobacterium tuberculosis* infection and clinical manifestation of tuberculosis. *Hum Immunol*. 2010; 71: 692–6.
- Van Crevel R, Ottenhoff TH, van der Meer JW. Innate immunity to *Mycobacterium tuberculosis*. *ClinMicrobiol Rev*. 2002; 15:294–309.
- van Crevel R, Parwati I, Sahiratmadja E, Marzuki S, Ottenhoff TH, Netea MG, et al., Infection with *Mycobacterium tuberculosis* Beijing genotype strains is associated with polymorphisms in SLC11A1/NRAMP1 in Indonesian patients with tuberculosis. *J Infect Dis* 2009;200:1671–1674.
- Van Kooyk Y, Appelmelk B, Geijtenbeek TB. A fatal attraction: *Mycobacterium tuberculosis* and HIV-1 target DC-SIGN to escape immune surveillance. *Trends Mol Med* 2003; 9:153–159.
- Vannberg FO, Chapman SJ, Khor CC, Tosh K, Floyd S, Sillah DJ, et al. CD209 genetic polymorphism and tuberculosis disease. *PLoS One* 2008; 3:e1388.
- Velayati AA, Masjedi MR, Farnia P, Tabarsi P, Gahnavi J, Ziazarifi AH, et al. Emergence of new forms of totally drug-resistant tuberculosis bacilli: super extensively drug-resistant tuberculosis or totally drug-resistant strains in Iran. *Chest* 2006; 136: 420–425.
- Velez DR, Hulme WF, Myers JL, Stryjewski ME, Abbate E, Estevan R, et al. Association of SLC11A1 with tuberculosis and interactions with NOS2A and TLR2 in African-Americans and Caucasians. *Int J Tuberc Lung Dis* 13:1068–1076.
- Velez DR, Hulme WF, Myers JL, Weinberg JB, Levesque MC, Stryjewski ME, et al. NOS2A, TLR4, and IFNGR1 interactions influence pulmonary tuberculosis susceptibility in African-Americans. *Hum Genet*. 2009; 126:643–53.
- Venter JC, Adams MD, Myers EW, Li PW, Mural RJ, Sutton GG, Smith HO, et al. The Sequence of the Human Genome. *Science* 2001; 1304–1351.
- Verbon A, Juffermans N, Deventer SV, Speelman P, Deutekom H, Poll V. Serum concentrations of cytokines in patients with active tuberculosis (TB) and after treatment. *Clin Exp Immunol*. 1999; 115:110–3.
- Vergne I, Chua J, Singh SB, Deretic V. Cell biology of *Mycobacterium tuberculosis* phagosome. *Annu Rev Cell Dev Biol*. 2004; 20: 367–94.
- Vidyanani M, Selvaraj P, Prabhu Anand S, Jawahar MS, Adhilakshmi AR, Narayanan PR. Interferon gamma (IFN $\gamma$ ) & interleukin-4 (IL-4) gene variants & cytokine levels in pulmonary tuberculosis. *Indian J Med Res*. 2006; 124:403–10.
- Vidyanani M, Selvaraj P, Raghavan S, Narayanan P R. Regulatory role of 1, 25-dihydroxyvitamin D3 and vitamin D receptor gene variants on intracellular granzyme A expression in pulmonary tuberculosis. *Exp Mol Pathol*. 2009; 86:69–73.
- Walker S, Chandra P, Maniava M, Axe E, Ktistakis NT. Making autophagosomes: localized synthesis of phosphatidylinositol 3-phosphate holds the clue. *Autophagy*. 2008; 4: 1093–96.
- Wall JD and Pritchard JK. Haplotype blocks and linkage disequilibrium in the human genome. *Nat Rev Genet* 2003; 4:587–597.
- Wallis RS, Pai M, Menzies D, Doherty TM, Walzl G, Perkins MD, Zumla A. 2010. Biomarkers and diagnostics for tuberculosis: progress, needs, and translation into practice. *Lancet* 2010; 375:1920 – 37.
- Walzl G, Ronacher K, Hanekom W, Scriba TJ, Zumla A. Immunological biomarkers of tuberculosis. *Nat Rev Immunol*. 2011; 11:343–54.
- Wanchu A, Bhatnagar A, Talreja J, Sapra S, Suryanarayana BS, Suresh P. Immunophenotypic and intracellular cytokine profile of Indian patients with tuberculosis with and without human immunodeficiency virus co-infection. *Indian J Chest Dis Allied Sci*. 2009; 51:207–11.
- Wang B, Henao-Tamayo M, Harton M, Ordway D, Shanley C, Basaraba RJ, et al. A Toll-like receptor-2-directed fusion protein vaccine against tuberculosis. *Clin Vaccine Immunol*. 2007; 14:902–6.



- Wang CH, Liu CY, Lin HC, Yu CT, Chung KF, Kuo HP. Increased exhaled nitric oxide in active pulmonary tuberculosis due to inducible NO synthase upregulation in alveolar macrophages. *Eur Respir J*. 1998; 11:809–15.
- Wang D, Johnson A D, Papp AC, Kroetz, DL, Sadee W. Multidrug resistance polypeptide 1 (MDR1, ABCB1) variant 3435C>T affects mRNA stability. *Pharmacogenet Genomics*. 2005; 15:693–704.
- Wang L, Cai Y, Cheng Q, Hu Y, Xiao H, Imbalance of Th1/Th2 cytokines in patients with pulmonary tuberculosis. *Zhonghua Jie He He Hu Xi Za Zhi*. 2002; 25:535-537.
- Wang N, Akey JM, Zhang K, Chakraborty R, Jin L. Distribution of recombination crossovers and the origin of haplotype blocks: the interplay of population history, recombination, and mutation. *Am J Hum Genet* 2002; 71:1227-1234.
- Wasylyk C, Schlumberger SE, Criqui-Filipe P, Wasylyk B. Sp100 interacts with ETS-1 and stimulates its transcriptional activity. *Mol Cell Biol*. 2002; 22:2687–702.
- Watashi K, Hijikata M, Tagawa A, Doi T, Marusawa H, Shimotohno K. Modulation of retinoid signaling by a cytoplasmic viral protein via sequestration of Sp110b, a potent transcriptional corepressor of retinoic acid receptor, from the nucleus. *Mol Cell Biol*. 2003; 23:7498–509.
- Werts C, Tapping RI, Mathison JC, Chuang TH, Kravchenko V, Saint Girons I , et al. Leptospiral lipopolysaccharide activates cells through a TLR2-dependent mechanism. *Nat. Immunol*. 2001; 2:346–52.
- White JH. Vitamin D signaling, infectious diseases, and regulation of innate immunity. *Infect Immun* 2008; 76:3837–43.
- Wichukchinda N, Kitamura Y, Rojanawiwat A, Nakayama EE, Song H, Pathipvanich P , et al. The polymorphisms in DC-SIGNR affect susceptibility to HIV type 1 infection. *AIDS Res Hum Retroviruses*. 2007; 23:686–92.
- Wigginton JE, Cutler DJ, Abecasis GR. A note on exact tests of Hardy- Weinberg equilibrium. *Am J Hum Genet*. 2005; 76:887-93.
- Wilbur A K, Kubatko L S, Hurtado A M, Hill K R, Stone A C. Vitamin D receptor gene polymorphisms and susceptibility *M. tuberculosis* in native Paraguayans. *Tuberculosis. (Edinb)* 2007; 87:329–37.
- Wilkinson RJ, Llewelyn M, Toossi Z, Patel P, Pasvol G, Lalvani A , et al. Influence of vitamin D deficiency and vitamin D receptor polymorphisms on tuberculosis among Gujarati Asians in West London: a case-control study. *Lancet*. 2000; 355: 618–21.
- Wilkinson RJ, Patel P, Llewelyn M, Hirsch CS, Pasvol G, Snounou G , et al. Influence of polymorphism in the genes for the interleukin (IL)-1 receptor antagonist and IL-1beta on tuberculosis. *J Exp Med*. 1999; 189:1863–74.
- Wilson AG, di Giovine FS, Blakemore AI, Duff GW. Single base polymorphism in the human TNF $\alpha$  gene detectable by NcoI restriction of PCR product. *Hum Mol Genet*. 1993; 1:353
- World Health Organization. (2011). Global tuberculosis control: WHO report 2011 (Geneva: World Health Organization).
- World Health Organization. Frequently asked questions - XDR-TB. <http://www.who.int/tb/challenges/xdr/faqs/en/index.html>
- Wu F, Qu Y, Tang Y, Cao D, Sun P, Xia Z , et al. Lack of association between cytokine gene polymorphisms and silicosis and pulmonary tuberculosis in Chinese iron miners. *J Occup Health*. 2008; 50:445–54.
- Wurfel MM, Gordon AC, Holden TD, Radella F, Strout J, Kajikawa O, et al. Toll-like Receptor 1 Polymorphisms Affect Innate Immune Responses and Outcomes in Sepsis. *Am J Respir Crit Care Med*. 2008; 178:710– 20.
- Xiao J, Sun L, Jiao WW, Li Z, Zhao S, Li H , et al. Lack of association between polymorphisms in the P2X7 gene and tuberculosis in a Chinese Han population. *FEMS Immunol Med Mic*. 2009; 55: 107–11.
- Xie Z, Klionsky DJ. Autophagosome formation: core machinery and adaptations. *Nat Cell Biol*. 2007; 9:1102–9.
- Xu J, Turner A, Little J, Bleecker ER, Meyers DA. Positive results in association studies are associated with departure from Hardy-Weinberg equilibrium: hint for genotyping error? *Hum Genet*. 2002; 111:573-4.
- Yamada G, Shijubo N, Shigehara K, Okamura H, Kurimoto M, Abe S. Increased levels of circulating interleukin-18 in patients with advanced tuberculosis. *Am J Respir Crit Care Med*. 2000; 161:76.
- Yang BF, Han CL. Meta-analysis of relationship of vitamin D receptor polymorphism and tuberculosis. *China Trop Med*. 2006; 6:1347–9.
- Ye S, Dhillon S, Ke X, Collins AR, Day IN. An efficient procedure for genotyping single nucleotide polymorphisms. *Nucleic Acids Res*. 2001; 29:E88.
- Yeager M, Orr N, Hayes RB, Jacobs KB, Kraft P, Wacholder S , et al. Genome-wide association study of prostate cancer identifies a second risk locus at 8q24. *Nat Genet*. 2007;39:645–9.
- Yilmaz V, Yentur SP, Saruhan-Direskeneli G. IL-12 and IL-10 polymorphisms and their effects on cytokine production. *Cytokine* . 2005; 30:188–194.
- Yim JJ, Ding L, Schäffer AA, Park GY, Shim YS, Holland SM , et al. A microsatellite polymorphism in intron 2 of human Toll-like receptor 2 gene: functional implications and racial differences. *FEMS Immunol Med Microbiol*. 2004; 40:163–9.
- Yim JJ, Lee HW, Lee HS, Kim YW, Han SK, Shim YS , et al. The association between microsatellite polymorphisms in intron II of the human Toll-like receptor 2 gene and tuberculosis among Koreans. *Genes Immun*. 2006; 7:150–5.
- Yim JJ, Selvaraj P. Genetic susceptibility in tuberculosis. *Respirology*. 2010; 15:241-56.

- YongPing X, GuoMei L, GuangXing C. Detection serum IL-2 level in patients with active tuberculosis before and after therapy. *Modern Preventive Medicine*. 2009; 36:4382-3.
- Yoshimura A, Lien E, Ingalls RR, Tuomanen E, Dziarski R, Golenbock D , et al. Cutting edge: recognition of Gram-positive bacterial cell wall components by the innate immune system occurs via Toll-like receptor 2. *J Immunol*. 1999; 163:1–5.
- Young D, Hessel T, Dougan G. Chronic bacterial infections: Living with unwanted guests. *Nat Immunol*. 2002; 3:1026–32.
- Yuk JM, Shin DM, Lee HM, Yang CS, Jin HS, Kim KK , et al. Vitamin D3 induces autophagy in human monocytes/macrophages via cathelicidin. *Cell Host Microbe*. 2009; 6:231–43.
- Zhang HQ, Guo CF, Chen J, et al. Influence of vitamin D receptor gene polymorphism on susceptibility of spinal tuberculosis in Chinese Han nationality. *Chin J Spine Spinal Cord*. 2006; 16:920–3.
- Zhang J, Zhang X, Fu J, Bi Z, Arheart KL, Barreiro LB , et al. Protective role of DC-SIGN (CD209) neck-region alleles with <5 repeat units in HIV-1 transmission. *J Infect Dis*. 2008; 198:68–71.
- Zhang W, Shao L, Weng X, Hu Z, Jin A, Chen S , et al. Variants of the natural resistance associated macrophage protein 1 gene (NRAMP1) are associated with severe forms of pulmonary tuberculosis. *Clin Infect Dis*. 2005; 40:1232–6.
- Zhang Y, Broser M, Cohen H, Bodkin M, Law K, Reibman J, et al. Enhanced interleukin-8 release and gene expression in macrophages after exposure to *Mycobacterium tuberculosis* and its components. *J. Clin. Invest*. 1995; 95: 586–92.