

LIST OF TABLES

Table No.	Description	Page No.
3.1	Composition of Murashige and Skoog's medium	18
3.2	Composition of MS medium supplemented with different concentrations of growth regulators used for <i>in vitro</i> establishment of papaya variety Pusa Nanha	20
3.3	Composition of various media supplemented with different growth hormones used for <i>in vitro</i> multiplication of papaya variety Pusa Nanha	22
3.4	Composition of various media supplemented with additives used for <i>in vitro</i> multiplication of papaya variety Pusa Nanha	23
3.5	Composition of MS half strength medium supplemented with various types of auxins with varying concentrations used for <i>in vitro</i> rooting	24
3.6	Different RAPD primers along with their sequence used for testing of genetic fidelity	27
3.7	Different ISSR primers along with their sequence used for testing of genetic fidelity	27
3.8	Reagents used for isolation of DNA from papaya leaves	28
3.9	Reagents used for electrophoresis of genomic DNA	28
3.10	PCR reagents for testing of genetic stability (RAPD)	29
3.11	PCR reagents for testing of genetic stability (ISSR)	30
3.12	PCR amplification cycle for RAPD primer	30
3.13	PCR amplification cycle for ISSR primer	30
3.14	Papaya known varieties and unknown genotypes collected from various regions of India	31
3.15	ISSR primers used for studying genetic diversity in papaya collected from different regions of India	32
3.16	RAPD primers used for studying genetic diversity in papaya collected from different regions of India	33
3.17	Varieties of papaya used for sex determination	34
3.18	SCAR and ISSR marker sequences used for sex determination	35
3.19	PCR amplification cycle for ISSR primer	35
3.20	PCR amplification cycle for SCAR primer	35
4.1	Effect of different durations of sterilizing agents on <i>in vitro</i> establishment of explants of <i>C. papaya</i> (survival percentage on 20 th day of inoculation)	39

4.2	Effect of different growth regulators on <i>in vitro</i> establishment of nodal segments	41
4.3	Effect of different growth regulators on <i>in vitro</i> multiplication of nodal segments (M1-M5)	47
4.4	Effect of different growth regulators on <i>in vitro</i> multiplication of nodal segments (M6-M10)	48
4.5	Effect of different growth regulators on <i>in vitro</i> multiplication using additives	56
4.6	Effect of different growth regulators on <i>in vitro</i> rooting	60
4.7	Effect of different potting mixture on percent survival of <i>in vitro</i> raised plantlets of <i>Carica papaya</i>	61
4.8	DNA amplification profile generated in Pusa Nanha of papaya using ISSR primers	62
4.9	DNA amplification profile generated in Pusa Nanha of papaya using RAPD primers	63
4.10	SCAR and ISSR markers sequences used for sex determination	64
4.11	Results of SCAR & ISSR for sex determination in five different varieties of papaya	66
4.12	Varieties of <i>C.papaya</i> collected	67
4.13	Different papaya varieties with Purity and Concentration values of their genomic DNA	68
4.14	DNA amplification profile and polymorphism generated in 18 different varieties of papaya using RAPD primers	70
4.15	DNA amplification profile and polymorphism generated in 18 different varieties of papaya using ISSR primers	72

LIST OF FIGURES

Fig. No.	Description	Page No.
3.1	Seedlings of Pusa Nanha in the greenhouse of CPB, Hisar	17
4.1	The percentage survival on sterilization treatments of nodal and shoot tip explants	40
4.2	<i>In vitro</i> establishment in <i>Carica papaya</i> using nodal segments after fix interval of time of culture (A) Nodal segments regeneration after 10 days (B) after 15 days (C) after 20 days (D) after 30 days	43
4.3	Effect of different concentration of growth regulators (PE ₀ - PE ₂₅) on number of buds and average days for initiation	44
4.4	Effect of different concentration of growth regulators (PE ₂₆ - PE ₅₀) on number of buds and average days for initiation	44
4.5	Effect of different concentration of growth regulators (PE ₀ - PE ₂₅) on percentage response for initiation	45
4.6	Effect of different concentration of growth regulators (PE ₂₆ - PE ₅₀) on percentage response for initiation	45
4.7	<i>In vitro</i> multiplication in <i>Carica papaya</i> after fixed interval of time (A) Shoot bud initiation after 28 days (B) Shoot multiplied on medium PM 27 (C) Shoot multiplied (highest) on medium PM 38	49
4.8	Effect of BAP(0.5mg/l) and GA ₃ (1-5mg/l) on mean no of shoots obtained on 28 th day of culture	50
4.9	Effect of BAP(0.5mg/l) and NAA(0.1-0.5mg/l) on mean no of shoots obtained on 28 th day of culture	50
4.10	Effect of BAP(0.5mg/l) and IBA(0.1-0.5mg/l) on mean no of shoots obtained on 28 th day of culture	51
4.11	Effect of BAP(0.5mg/l) and IAA(0.1-0.5mg/l) on mean no of shoots obtained on 28 th day of culture	51
4.12	Effect of BAP (0.5mg/l), GA ₃ (4.0mg/l) and NAA(0.1-0.5mg/l) on mean no of shoots obtained on 28 th day of culture	52
4.13	Effect of KIN(1.0mg/l),GA ₃ (1mg/l) and NAA(0.2mg/l) on mean no of shoots obtained on 28 th day of culture	52
4.14	Effect of KIN(1.0mg/l) and GA ₃ (1-5mg/l) on mean no of shoots obtained on 28 th day of culture	53
4.15	Effect of KIN(1.0mg/l) and NAA(0.1-0.5mg/l) on mean no of shoots obtained on 28 th day of culture	53
4.16	Effect of Zeatin (2.0mg/l) and NAA (0.1-0.5mg/l) on mean no of shoots obtained on 28 th day of culture	54
4.17	Effect of TDZ (1.35 mg/l) and NAA (0.1-0.5mg/l) on mean no of shoots obtained on 28 th day of culture	54
4.18	<i>In vitro</i> multiplication with additives in <i>Carica papaya</i> after fixed interval of time (A) Shoot bud initiation after 14 days (B) after 28 days	57

	(C) Shoot multiplied on medium PMA3 (D) Shoot multiplied (highest) on medium PMA2	
4.19	Effect of conc. Of AgNO ₃ on mean number of shoots and percentage of leaf fall on 28 th day of culture	58
4.20	Effect of conc. Of adenine sulphate (Ads) on mean number of shoots and percentage of leaf fall on 28 th day of culture	58
4.21	Effect of conc. Of PEG on mean number of shoots and percentage of leaf fall on 28 th day of culture	59
4.22	<i>In vitro</i> rooting in <i>Carica papaya</i> after three weeks of culture (A) Rooting on R5 media (B) Rooting on R4 media (C) Rooting on R3 media	60
4.23	Hardening of regenerated plants of <i>Carica papaya</i> (A) Freshly transferred regenerated plants in polybags in Green House B) Well developed plant transferred to soil under field conditions	61
4.24	PCR profile of mother plant and 17 <i>in vitro</i> regenerated plants using ISSR AY 21	63
4.25	PCR profile SCAR and ISSR markers used in Sex determination A) ISSR B) SCAR 1 C) SCAR 2 D) SCAR 3	65
4.26	PCR profile of Standard Male (M) and Female (F) with 12 unknown DNA samples of papaya seedlings, showing Sample 6 as female plant DNA	66
4.27	Variations in DNA concentrations in different papaya varieties	69
4.28	Gel profile of purified genomic DNA sample of 18 varieties of papaya	69
4.29	Gel profile of RAPD OPA-1 with three polymorphic bands	71
4.30	Gel profile of RAPD OPA-9 with two monomorphic bands	71
4.31	Gel profile of ISSR AY-27 with five monomorphic bands	73
4.32	Gel profile of ISSR AY-45 with two polymorphic bands	73
4.33	Dendogram based on RAPD and ISSR analysis among 18 papaya varieties	75
4.34	Similarity matrix among 18 papaya varieties based on RAPD and ISSR analysis	76
4.35	2D map constructed among 18 papaya varieties based on RAPD and ISSR analysis	77
4.36	3D map constructed among 18 papaya varieties based on RAPD and ISSR analysis	77