

List of Figures

2.1	Schematic diagram of the 90° beamline at LEIBF with the molecular dissociation setup at its end.	9
2.2	Single field time-of-flight mass spectrometer.	11
2.3	Double field linear time-of-flight mass spectrometer.	13
2.4	Composite mechanical drawing of the IUAC TOFMS with detectors.	17
2.5	The principle of a delay line anode.	19
2.6	Electronic setup of the position-sensitive multi-hit detector system. The different abbreviations mean the following. DA: differential amplifier, CFD: constant fraction discriminator, TA: timing amplifier, PA: pre-amplifier, D: delay, NEC: NIM-ECL converter, TDC: time-to-digital converter, DAC: data acquisition computer	21
2.7	Collection efficiency as a function of fragment charge and fragment energy for an extraction field of 114 V/cm	23
2.8	Flowchart showing conversion of raw binary data into ROOT format	25
2.9	A typical time-of-flight spectra from nitrogen dissociation. The subscript f is for the fragment directed along the applied extraction field and b is for the fragment directed opposite to the applied extraction field.	27
2.10	Typical TOF spectra of a monoatomic gas, Argon.	28
2.11	A typical two dimensional histogram : the coincidence map	29
3.1	Potential energy curves depicting dissociative states of diatomic molecules.	39

3.2	Potential energy curves for N_2^{3+} . The curves represent the results of MCSCF energy calculation as a function of inter-nuclear distance.	42
3.3	Potential energy curves for N_2^{4+} . The curves represent the results of MCSCF energy calculation as a function of inter-nuclear distance.	42
3.4	Potential energy curves for N_2^{5+} . The curves represent the results of MCSCF energy calculation as a function of inter-nuclear distance.	43
3.5	Potential energy curves for N_2^{6+} . The curves represent the results of UHF energy calculation as a function of inter-nuclear distance.	44
3.6	Potential energy curves for CO^{4+} . The curves represent the results of UHF energy calculation as a function of inter-nuclear distance.	46
3.7	Potential energy curves for CO^{5+} . The curves represent the results of UHF energy calculation as a function of inter-nuclear distance.	47
3.8	Potential energy curves for CO^{6+} . The curves represent the results of UHF energy calculation as a function of inter-nuclear distance.	47
4.1	Coincidence map of the N_2 fragmentation	53
4.2	Kinetic energy release spectra of the $N^+ - N^+$ fragmentation channel	56
4.3	Kinetic energy release spectra of the $N^+ - N^{2+}$ fragmentation channel	57
4.4	Kinetic energy release spectra of the $N^{2+} - N^{2+}$ and $N^{1+} - N^{3+}$ fragmentation channel	59
4.5	Kinetic energy release spectra of the $N^{2+} - N^{3+}$ fragmentation channel	60
4.6	Kinetic energy release spectra of the $N^{3+} - N^{3+}$ and $N^{2+} - N^{4+}$ fragmentation channel	61
4.7	Kinetic energy release spectra of the $N^{1+} - N^{1+}$ fragmentation channel obtained for different incident projectile charge state.	62
4.8	Kinetic energy release spectra of the $N^{1+} - N^{2+}$ fragmentation channel obtained for different incident projectile charge state.	62

4.9	Kinetic energy release spectra of the $N^{2+} - N^{2+}$ fragmentation channel obtained for different incident projectile charge state. . . .	63
4.10	Kinetic energy release spectra of the $N^{1+} - N^{3+}$ fragmentation channel obtained for different incident projectile charge state. . . .	63
4.11	Kinetic energy release spectra of the $N^{2+} - N^{3+}$ fragmentation channel obtained for different incident projectile charge state. . . .	64
4.12	Kinetic energy release spectra of the $N^{3+} - N^{3+}$ fragmentation channel obtained for different incident projectile charge state. . . .	64
4.13	Kinetic energy release spectra of the $N^{2+} - N^{4+}$ fragmentation channel obtained for different incident projectile charge state. . . .	65
5.1	Kinetic energy release spectra for CO^{2+} dissociation. Theoretical values are indicated by vertical lines at the top of the graph.	75
5.2	Kinetic energy release spectra for CO^{3+} dissociation. Theoretical values are indicated by vertical lines at the top of the graph.	76
5.3	Kinetic energy release spectra for CO^{4+} dissociation. Theoretical values are indicated by vertical lines at the top of the graph.	77
5.4	Potential energy curves of CO^{4+}	79
5.5	Kinetic energy release spectra for CO^{5+} dissociation. Theoretical values are indicated by vertical lines at the top of the graph.	79
5.6	Kinetic energy release spectra for CO^{6+} dissociation. Theoretical values are indicated by vertical lines at the top of the graph.	81
6.1	A parallel beam of diameter 10mm and energy 30 keV/q is focused to a spot of 4mm diameter and 5 eV/q energy at the target with 100% transmission efficiency.	89
6.2	Equipotential surfaces for deceleration from 30 keV/q to 5 eV/q	90
6.3	A parallel beam of diameter 10mm and energy 30 keV/q is focused to a spot of 2mm diameter and 500 eV/q energy at the target with 100% transmission efficiency.	91
6.4	A beam of 5mm radius and ~ 35 mrad divergence having initial energy of 30 keV/q is focused with 85% of the beam within a radius of 1.5mm at the target.	92
6.5	The variation of energy along the length of the deceleration lens	93

LIST OF FIGURES

viii

6.6	<i>SOLIDWORKS</i> : Cut out view of the mechanical design of the deceleration lens	94
6.7	Schematic of experimental chamber (beam level).	95
6.8	<i>SOLIDWORKS</i> : Chamber lid designed for finer target alignment .	96
6.9	<i>SOLIDWORKS</i> : Assembly of deceleration lens with experimental chamber.	96
6.10	Schematic of the beamline of the PKDELIS ECR ion source.	97
6.11	Actual photograph of the deceleration lens assembly.	98
6.12	Repeller plate analyser	100
6.13	Picture of the beamline with the deceleration lens and experimental chamber	101