

References

1. **Affinity Chromatography: Principles and Methods.** Pharmacia fine chemicals, Uppsala, Sweden, 1979.
2. **Algarra, I., Ohlen, C., Perez, M., Ljunggren, H.G., Klein, G., Garrido, F. and Karre, K.** NK sensitivity and lung clearance of MHC-class I-deficient cells within a heterogeneous fibrosarcoma. *Int. J. Cancer.* 44, 675, 1989.
3. **Anderson, M., Paabo, S., Nilsson, T. and Peterson, P.A.** Impaired intracellular transport of class I MHC antigens as a possible means for adenoviruses to evade immune surveillance. *Cell.* 43, 215, 1985.
4. **Andersson, P., Caliguri, M., Ritz, J. and Scholassman, S.F.** CD3- negative natural killer cells express TCR as part of a novel molecular complex. *Nature,* 341,159, 1989.
5. **Bean, P., Agah, R. and Mazumder, A.** Heterogeneity of cell surface structures involved in cytotoxicity mediated by LAK cells. *J. Biol. Resp. Modif.* 9, 92, 1990.
6. **Bernards, R., Dessain, S.K. and Weinberg, R.A.** N-myc amplification causes down modulation of MHC class I antigen expression in neuroblastoma. *Cell.* 47, 667, 1986.
7. **Bialas, T., Kolitz, J., Levi, E., Polivka, A., Oez, S., Miller, G. and Welte, K.** Distinction of partially purified human natural killer cytotoxic factor from recombinant human tumor necrosis factor and recombinant human lymphotoxin. *Cancer Res.* 48, 891, 1988.
8. **Biron, C.A., Van Den Elsen, P., Tutt, M.M., Medveczky, P., Kumar, V. and Terhorst, C.** Murine natural killer cells stimulated in vivo do not express the T cell receptor α , β , γ , T3 δ , T3 ϵ genes. *J. Immunol.* 139, 1704, 1987.
9. **Biron, C.A., Byron, K.S. and Sullivan, J.L.** Severe herpes virus infection in an adolescent without natural killer cells. *New Engl. J. Med.* 320, 1731, 1989.
10. **Blottiere, H.M., Zennadi, R., Burg, C., Douillard, J.Y., Meflah, K. and Le Pendu, J.** Relationship between sensitivity to natural killer cells and MHC class-I antigen expression in colon carcinoma cell lines. *Int. J. Cancer.* 50, 659, 1992.
11. **Bouwens, L., Remels, L., Baekeland, M., van Bossuyr, H. and Wisse, E.** Large granular lymphocytes of "Pit cells" from rat liver: isolation, ultrastructural characterization and natural killer activity. *Eur. J. Immunol.* 17, 37, 1987.

12. **Brooks, C.G., Burton, R.C., Pollack, S.B. and Henney, C.S.** The presence of NK alloantigens on cloned cytotoxic T lymphocytes. *J. Immunol.* 131, 1391, 1983.
13. **Brunda, M.J., Herberman, R.B. and Holden, H.T.** Antibody induced augmentation of murine natural cell killer activity. *Int. J. Cancer.* 27, 205, 1981.
14. **Carbone, E., Pacioppi, L., La Cava, A., Portella, G., Velotti, F., Zappacosta, S. and Fontana, S.** NK and LAK susceptibility varies inversely with target cell MHC class I antigen expression in a rat epithelial tumor system. *Scand. J. Immunol.* 33, 185, 1991.
15. **Carpen, O., Virtanen, I. and Saksela, E.** Ultrastructure of human natural killer cells: nature of the cytolytic contacts in relation to cellular secretion. *J. Immunol.* 128, 2691, 1982.
16. **Carpen, O., Virtanen, I., Letho, V.P. and Saksela, E.** Polarization of NK cell cytoskeleton upon conjugation with sensitive target cells. *J. Immunol.* 131, 2695, 1983.
17. **Chambers, W.H., Vujanovic, N.L., Delso, A.B., Olszowy, M.W., Herberman, R.B. and Hiserodt, J.C.** Monoclonal antibodies to a triggering structure expressed on rat natural killer cells and adherent lymphokine activated killer cells. *J. Exp. Med.* 169, 1373, 1989.
18. **Chervenak, R. and Wolcott, R.M.** Target cell expression of MHC antigens is not (always) a turn-off signal to natural killer cells. *J. immunol.* 140, 3712, 1988.
19. **Dawson, J.R., Storkus, W.J., Patterson, E.B. and Cresswell, P.** Adenovirus inversely modulates target cell class I MHC antigen expression and sensitivity to natural killing. In *Natural Killer cells and host defenses*, Ades, E.W. and Lopez, C. Eds., S. Karger, Basel, 156, 1989.
20. **De Fries, R.U. and Golub, S.H.** Characteristics and mechanisms of Interferon-gamma induced protection of human tumor cells from lysis by LAK cells. *J. Immunol.* 140, 3686, 1988.
21. **De Fries, R.U. and Golub, S.H.** Interferon-gamma reduces the sensitivity of cultured and fresh human tumor cells to lysis by lymphokine-activated killer cells. *Nat. Immun. Cell Growth Regul.* 7, 65, 1988.
22. **De Mayer, E. and De Mayer Guignard, J.** Interferons as multifunctional gene activators. In *interferons and other regulatory cytokines*. John Wiley and Sons, New York, 91, 1982.

23. **Doherty, P.C., Knowles, B.B. and Wettstein, P.J.** Immunological surveillance of tumors in the context of major histocompatibility complex restriction on T cell function. *Adv. Cancer Res.* 42, 1, 1984.
24. **Due, C., Simonsen, M. and Olsson, L.** The major histocompatibility complex class I heavy chain as a structural subunit of the human cell membrane insulin receptor: implications for the range of biological functions of histocompatibility antigens. *Proc. Natl. Acad. Sci. U.S.A.* 83, 6007, 1986.
25. **Evans, C. H., Heinbaugh, J.A. and Ransom, J.H.** Flow cytometric evaluation of leukoregulin as an intrinsic molecular mediator of natural killer lymphocyte cytotoxicity. *Lymphokine Res.* 2, 277, 1987.
26. **Gately, M.K., Desal, B.B., Wolitzky, A.G., Quinn, P.M., Dawyer, C.M., Podlaski, F.J., Familletti, P.C., Sinigaglia, F., Chizzonite, R., Gubler, U. and Stern, A.S.** Regulation of human lymphocyte proliferation by heterodimeric cytokine, IL-2 (cytotoxic lymphocyte maturation factor). *J. Immunol.* 147, 874, 1991.
27. **Gillis, S., Ferm, M.M., Ou, W. and Smith, K.A.** T cell growth factor: parameters of production and a quantitative microassay for activity. *J. Immunol.* 120, 2027, 1978.
28. **Grimm, E.A., A. Mazumder, K.Z.Zhang, and S.A. Rosenberg.** Lymphokine-activated killer cell phenomenon: lysis of natural-killer cell resistant fresh solid tumor cells by interleukin-2 activated autologous human peripheral blood lymphocytes. *J. Exp. Med.* 155, 1823, 1982.
29. **Gubler, U., Chua, A.O., Schoenhaut, D.S., Dawyer, C.M., Mc Comas, W., Motyka, R., Nabavi, N., Wolitzky, A.G., Quinn, P.M., Familletti, P.C. and Gately, M.K.** Coexpression of two distinct genes is required to generate secreted bioactive cytotoxic lymphocyte maturation factor. *Proc. Natl. Acad. Sci. U.S.A.* 88, 143, 1991.
30. **Gaopas, J., Segal, S., Hammerling, G., Bar-Eli, M. and Reyer-Zisman.** Influence of H-2K transfection on susceptibility of fibrosarcoma tumor cell to natural killer (NK) cells. *Immunol. Lett.* 17, 261, 1988.
31. **Gorelik, E., Gunji, Y. and Herberman, R.B.** H-2 antigen expression and sensitivity of BL6 melanoma cells to natural killer cell cytotoxicity. *J. Immunol.* 140, 2096, 1988.

32. **Hanson, M., Karre, K., Bakacs, T., Kiessling, R. and Klein, G.** Intra- and interspecies reactivity of human and mouse natural killer (NK) cells, *J. Immunol.* 121, 6, 1979.
33. **Hanson, M., Karre, K., Kiessling, R. and Roder, J.** NK cell-sensitive T-cell subpopulation in thymus: inverse correlation to host NK activity. *Nature.* 278, 174, 1979.
34. **Hanson, M., Kiessling, R., Andersson, B. and Welsh, R.M.** Effect of interferon and interferon inducers on the NK sensitivity of normal mouse thymocytes. *J. Immunol.* 125, 2225, 1980.
35. **Hanson, M. and Kiessling, R.** Natural killing of hematopoietic cells. in *NK and Other Natural Effector Cells*, Herberman, R.B., Ed., Academic Press, New York, 1077, 1987.
36. **Harel-Bellan, A., Quillet, A., Marchiof, C., DeMars, R., Tursz, T. and Fradelizi, D.** Natural killer susceptibility of human cells may be regulated by genes in the HLA regions of chromosome. *Proc. Nat. Acad. Sci. U.S.A.* 83, 5688, 1986.
37. **Hemler, M.E.** VLA proteins in the integrin family: structure, function and their role in leukocytes. *Annu. Rev. Immunol.* 8, 365, 1990.
38. **Herberman, R.B. and Holden, H.T.** Natural cell mediated immunity. *Adv. Cancer Res.* 27, 377, 1978.
39. **Herberman R.B., Reynold C.B. & Ortaldo J:** Mechanisms of cytotoxicity by natural killer cells. *Ann. Rev. of Immunol.* 4, 651, 1986.
40. **Herberman, R., Balch, C., Bolhius, R., Golub, S., Hiserodt, J., Lanier, L., Philips, J., Riccardi, C., Ritz, J., Santoni, A., Schimdt, R., Uchidi, A. and Vujanovic, N.** Most lymphokine activated killer (LAK) activity mediated by blood and splenic lymphocytes is attributable to stimulation of natural killer (NK) cells by interleukin-2 (IL-2). *Immunol. Today.* 8, 178, 1987.
41. **Hiserodt, J.C., Britvan, L.J. and Targan, S.R.** Differential effects of various pharmacologic agents on the cytolytic reaction mechanism of the natural killer lymphocyte: further resolution of programming for lysis and KCIL into discrete stages. *J. Immunol.* 129, 2266, 1982.
42. **Hiserodt, J.C., Laybourn, K.A. and Varani, J.** Expression of laminin-like substances on the surface of murine natural killer lymphocytes and its role in NK recognition. *J. Immunol.* 135, 1484, 1985.

43. **Hiserodt, J.C., Laybourn, K.A. and Varani, J.** Laminin inhibits the recognition of tumor target cells by murine natural killer (NK) natural cytotoxic (NC) lymphocytes. *Am. J. Pathol.* 121, 148, 1985.
44. **Hiserodt, J.C., and Schwartz, R.E.** A surface structure involved in target cell recognition by non-MHC-restricted cytotoxic effector cells. *Tissue Antigens* 33, 301, 1989.
45. **Jennings, S.R., Rice, P.L., Kloszewski, E.D., Anderson, R.W., Thompson, D.L. and Tevethia, S.S.** Effect of herpes simplex virus types 1 and 2 on surface expression of class I major histocompatibility complex antigens on infected cells. *J. Virol.* 56, 757, 1985.
46. **Jondal, M., Kallman, C., Alter, M.B. and Ljunggren, K.** Monoclonal antibodies against the NK cell-FcR and T3-complex potentiate normal lymphocyte killing. *Cell. Immunol.* 100, 158, 1986.
47. **Kalland, T.H., Belfrage, P. Bhiladvala, and G. Hedlund.** Analysis of the murine lymphokine-activated killer (LAK) cell phenomenon: Dissection of effectors and progenitors into NK- and T- like cells. *J. Immunol.* 138, 3640, 1987.
48. **Karre, K., Ljunggren, H.G., Piontek, G., Kiessling, R., Klein, G., Taniguchi, K. and Gronberg, A.** Activation of cell mediated immunity by absence or deleted expression of normal cellular gene products i.e. by "no self" rather than "non self". *Immunobiology.* 167, 43, 1984.
49. **Karre, K., Ljunggren, H.G., Piontek, G. and Kiessling, R.** Selective rejection of H-2 deficient lymphoma variants suggests alternative immune defence strategy. *Nature (Lond.).* 319, 675, 1986.
50. **Karlhofer, F.M., Ribaldo, R.K. and Yokoyama, W.M.** MHC class I alloantigen specificity of Ly-49⁺ IL-2 activated natural killer cells. *Nature.* 358, 66, 1992.
51. **Kawano, Y.I., Taniguchi, K., Toshihara, A. and Nomoto, K.** Synergistic defense system by cooperative natural effectors against metastasis of B16 melanoma cells in H-2 associated control: different behavior of H-2⁺ and H-2⁻ cells in metastatic processes. *J. Immunol.* 136, 4729, 1986.
52. **Kiessling, R., Hochman, P.S., Haller, O., Shearer, G.M., Wigzell, H. and Cudkowicz, G.** Evidence for a similar or common mechanism for natural killer cell activity and resistance to haematopoietic grafts. *Eur. J. Immunol.* 7, 655, 1977.

53. **Klein, J.** Biology of the Mouse Histocompatibility-2 complex. Springer Verlag, New York, 1975.
54. **Krangel, M.S., Orr, H.T. and Strominger, J.L.** Assembly and maturation of HLA-A and HLA-B antigens in vivo. *Cell*. 18, 979, 1979.
55. **Kupfer, A., Dennert, G. and Sinfer, S.J.** Polarization of the Golgi apparatus and the microtubule-organizing center within cloned natural killer cells bound to their targets. *Proc. Nat. Acad. Sci. U.S.A.* 80, 7224, 1983.
56. **Lanier, L.L., Kipps, T.J. and Philips, J.H.** Functional properties of a unique subset of cytotoxic CD3⁺ lymphocytes that express Fc receptors for IgG (CD16/Leu 11 antigen). *J. Exp. Med.* 163, 2089, 1985.
57. **Lanier, L.L., Le, A.M., Civin, C.I., Loken, M.R. and Phillips, J.H.** The relationship of CD16 (Leu-11) and Leu-19 (NKH-1) antigen expression on human peripheral blood NK cells and cytotoxic T lymphocytes. *J. Immunol.* 136, 4480, 1986.
58. **Lanier, L.L. and Phillips, J.H.** NK-cells: membrane-structures triggering cell mediated cytotoxicity, in *Leukocyte Typing IV. White Cell Differentiation Antigens*. Knapp, W., Dorken, B., Gilks, W.R., Rieber, E.P., Schmidt, R.E., Stein, R.H. and von dem Borne, A.E.G.Kr., Eds., Oxford University press, Oxford, 1064, 1989.
59. **Lanier, L.L., Yu, G. and Phillips, J.H.** Co-association of CD36 with a receptor (CD16) for IgG Fc on human natural killer cells. *Nature*. 342, 803, 1989.
60. **Lanier, L.L.** NK and T-lymphocytes: membrane structure triggering cell-mediated cytotoxicity. *J. Cell. Biochem.* 14B, 52, 1990.
61. **Laybourn, K.A., Hiserodt, J.C. and Varani, J.** Laminin receptor expression on murine tumor cells: correlation with sensitivity to natural cell-mediated cytotoxicity. *Int. J. Cancer*. 43, 737, 1989.
62. **Ledbetter, J.A., Tonks, N.K., Fischer, E.H. and Clark, E.A.** Cd45 regulates signal transduction and lymphocyte activation by specific association with receptor molecules on T or B cells. *Proc. Natl. Acad. Sci. U.S.A.* 85, 8628, 1988.

63. **Leiden, J.M., Gottesdiener, K.M., Quertermous, T., Coury, T., Bray, R.A., Gottschalk, L., Gebel, H., Seidman, J.G., Strominger, J.L, Landay, A.L. and Kornbluth, J.** T cell receptor arrangement and expression in human natural killer cells: natural killer activity is not dependent on the rearrangement and expression of T-cell receptor alpha, beta and gamma genes. *Immunogenetics* 27, 231, 1988.
64. **Ljunggren, H.G. and Karre, K.** Host resistance directed selectivity against H-2 deficient lymphoma variants. analysis of the mechanism. *J. Exp. Med.* 162, 1745, 1985.
65. **Ljunggren, H.G., Paabo, S., Cochet, M., Kling, G., Kuorilsky, P. and Karre, K.** Molecular analysis of H-2 deficient lymphoma lines. Distinct defects in biosynthesis and association of MHC class I heavy chain and β -microglobulin observed in cells with increased sensitivity to NK lysis. *J. Immunol.* 142, 2911, 1989.
66. **Ljunggren, H.G. and Karre, K.** In search of the "missing self":MHC molecules and NK cell recognition. *Immunol. Today* 11, 237, 1990.
67. **Lopez-Nevot, M.A., Esteban, F., Ferron, A., Guitierrez, J., Oliva, M.R., Romero, C., Havelin, C., Ruiz-Cabello, F. and Garrido, F.** HLA class I gene expression on human primary tumors and autologous metastasis:demonstration of selective losses of HLA antigens on colorectal, gastric and laryngeal carcinomas. *Br. J. Cancer* 59, 221, 1989.
68. **Luger, T.A., Krutmann, J., Kirnbauer, R., Urbanski, A., Schwartz, T., Klappacher, G., Kock, A., Micksche, M., Malejczyk, J., Schauer, E., May, L.T. and Sehgal, P.B.** IFN- β , IL-6, augments the activity of human natural killer cells. *J. Immunol.* 143, 1206, 1989.
69. **Main, E.K., Lampson, L.A., Hart, M.K., Kornbluth, J. and Wilson, D.B.** Human neuroblastoma cell lines are susceptible to lysis by natural killer cells but not cytotoxic T lymphocytes. *J. Immunol.* 135, 242, 1985.
70. **Masuyama, K., Ochiai, H., Ishizawa, S., Tazawa, K., Niwayama, S. and Fujimaki, M.** Relation of H-2 expression on murine RCT (+) sarcoma cells to lung colonization and sensitivity to NK cells. *J. Cancer Res. Clin. Onc.* 114, 487, 1988.
71. **Miyatake, S.I., Haruhiko, K., Oda, Y., Nishioka, T., Aoki, T., Kasakura, S. and Namba, Y.** Decreased susceptibility of lined human gliosarcoma cells to lymphokine-activated killer cell cytolysis by gamma interferon treatment. *Cancer Res.* 50, 596, 1990.

72. **Moore, M., White, W.J. and Potter, M.R.** Modulation of target cell susceptibility to human natural killer cells by IFN. *Int. J. Cancer.* 25, 565, 1980.
73. **Mule, J.J., Smith, C.A. and Rosenberg, S.A.** Interleukin 4 (B cell stimulatory factor I) can mediate the induction of lymphokine-activated killer cell activity directed against fresh tumor cells. *J. Exp. Med.* 166, 792, 1982.
74. **Nagler, A., Lanier, L.L. and Phillips, J.H.** Comparative studies of studies of human FcγIII-positive and negative natural killer cells. *J. Immunol.* 143, 3183, 1989.
75. **Nakane, P.K. and Kawaoi, A.** Peroxidase-labelled antibody: A new method of conjugation. *J. Histochem. Cytochem.* 22, 1084, 1974.
76. **Obexer, G., Rumpold, H. and Kraft, D.** Target-effector-cell interactions in the human natural killer (NK) system: isolation of target structures. *J. Immunol.* 165, 15 1983.
77. **Ohlen, C., Bejarano, M.T., Gronberg, A., Torsteinsdottir, S., Franksson, L., Ljunggren, H.G., Klein, G. and Karre, K.** Studies of sublines selected for loss of HLA expression from an EBV-transformed lymphoblastoid cell line: changes in sensitivity to cytotoxic T cells activated by IFN or IL 2. *J. Immunol.* 142, 3336, 1989.
78. **Ortaldo, J.R., and Herberman, R.B.** Heterogeneity of natural killer cells. *Annu. Rev. Immunol.* 2, 359, 1984.
79. **Ortaldo, J.R., Kantor, R., Segal, D., Bolhuis, R.H. and Bino, T.** Identification of a proposed NK receptor. in *Natural Killer Cells and Host Defense*, Ades, E.D. and Lopez, C., Eds., Karger. Basel, 221, 1989.
80. **Ortaldo, J.R., Anderson, S., Roder, J., Okamura, K., Sugamura, K., Frey, J., Smyth, M. and Young, H.A.** Mechanism of lysis by human natural killer cells. *J. Cell. Biochem.* 14B, 8 1990.
81. **Ostensen, M.E., Thiele, D.L. and Lipsky, P.E.** Tumor necrosis factor- α enhances cytolytic activity of human natural killer cells. *J. Immunol.* 138, 4185, 1987.
82. **Owen-Schaub, L.B., Gutterman, J.U. and Grimm, E.A.** Synergy of tumor necrosis factor α and interleukin 2 in the activation of human cytotoxic lymphocytes: Effect of tumor necrosis factor and interleukin 2 in the generation of human lymphokine-activated killer cell cytotoxicity. *Cancer Res.* 48, 788, 1988.

83. Palmieri, G., Morrone, S., Lollini, P.L., DeGeovanni, C., Nicoletti, G., Nanni, P., Frati, L. and Santoni, A. TNF impairs in vivo and in vitro natural killer (NK) susceptibility of B16 melanoma cells. *Scand. J. Immunol.* 35, 279, 1992.
84. Pawelec, G., Newman, W., Schwulera, U. and Warner, P. Heterogeneity of human natural killer recognition demonstrated by cloned effector cells and differential blocking of cytotoxicity with monoclonal antibodies. *Cell. Immunol.* 92, 31, 1985.
85. Pena, J., Alonso, C., Solona, R., Serrano, R., Carracedo, J. and Ramirez, R. Natural killer susceptibility is independent of HLA class I antigen expression on cell lines obtained from human solid tumors. *Eur. J. Immunol.* 20, 2445, 1990.
86. Pestka, S., Langer, J.A., Zoon, K.C. and Samuel, C.E. Interferons and their actions. *Annu. Rev. Biochem.* 56, 727, 1987.
87. Piontek, G.E., Taniguchi, K., Ljunggren, H.G., Gronberg, A., Kiessling, R., Klein, G. and Karre, K. YAC-I MHC class I variants reveal an association between decreased NK sensitivity and increased H-2 expression after interferon treatment or in vivo passage. *J. Immunol.* 135, 4281, 1985.
88. Pollack, S.B. and Rosse, C. The primary role of murine bone marrow in the production of natural killer cells. A cytokinetic study. *J. Immunol.* 139, 2149, 1987.
89. Quan, P.C., Ishizaka, T. and Bloom, B.R. Studies on the mechanisms of NK lysis. *J. Immunol.* 128, 1786, 1982.
90. Quillet, A., Presse, F., Marchio-Fournigault, C. Harel-Bellan, A., Benbunam, M., Ploegh, H. and Fradelizi, D. Increased resistance to non-MHC-restricted cytotoxicity related to HLA A, B expression. direct demonstration using β_2 -microglobulin-transferred Daudi cells. *J. Immunol.* 141, 17, 1988.
91. Ramirez, R., Solona, R., Carracedo, J., Alonso, M.C. and Pena, J. Mechanism involved in NK resistance induced by IFN- gamma. *Cell. Immunol.* 140, 248, 1992.
92. Reitzer, Z., Fischer, D.G. and Rubinstein, M. The protective effect of interferon against natural killing activity is not via the expression of class I MHC antigens. *Immunol. Lett.* 17, 323, 1988.

93. Reynold, C.W., Sharrow, S.O., Ortaldo, J.R. and Herberman, R.B. Natural killer activity in the rat. II Analysis of surface antigens on LGL by flow cytometry. *J. Immunol.* 127, 2204, 1981.
94. Ritz, J., Schmidt, R.E., Michon, J., Hercend, T. and Schlossman, S.F. Characterization of functional surface structures on human natural killer cells. *Adv. Immunol.* 42, 181, 1988.
95. Roder, J.C., Rosen, A., Fenyo, E.M. and Troy, F.A. Target effector interaction in the natural killer system: isolation of target structure. *Proc. Nat. Acad. Sci. U.S.A.* 76, 1405, 1979.
96. Roder, J.C. and Pross, H.F. The biology of natural killer cell. *J. Clin. Immunol.* 2, 249, 1982.
97. Roder, J.C., Karre, K. and Kiessling, R. Natural killer cells. *Prog. Allergy.* 28, 66, 1991.
98. Rosenberg, S.A., M.T. Lotze, L.M. Muul, A.E. Chang, F.P. Avis, S. Leitman, W.M. Lineham, C.N. Robertson, R.E. Lee, J.T. Rubin, C.A. Seipp, C. Simpson, and D.E. White. A progress report on the treatment of 157 patients with advanced cancer using lymphokine-activated killer cells and interleukin-2 or high dose interleukin-2 alone. *N. Engl. J. Med.* 316, 889, 1987.
99. Rosenstein, M., I.Yron, Y. Kaufmann, and S.A. Rosenberg. Lymphokine-activated cells: Lysis of fresh syngeneic natural killer resistant murine tumor cells by lymphocytes cultured in interleukin-2. *Cancer Res.* 44, 1946, 1984.
100. Routes, J.M. and Cook, J.L. Adenovirus persistence in man. Defective E1A gene product targeting of infected cells for elimination by natural killer cells. *J. Immunol.* 142, 4022, 1989.
101. Ryan, J.C., Turck, J. and Seaman, W.E. Molecular cloning of the NK 1.1 antigen, a member of the NKRP-1 family of natural killer cell activation molecules. *J. Immunol.* 149, 1631, 1992.
102. Salup, R.R., B.J. Mathieson, and R.H. Wiltrot. Precursor phenotype of lymphokine-activated killer cells in the mouse. *J. Immunol.* 138, 3635, 1987.

103. Santoli, A., Gismondi, A., Marrone, S., Procopio, A., Modesti, A., Scarpa, S., D'Orazi, G., Piccoli, M. and Frati, L. Rat natural killer cells synthesize fibronectin: possible involvement in the cytotoxic function. *J. Immunol.* 143, 2415, 1989.
104. Sarin, A. and Saxena, R.K. Quantitative estimation of major histocompatibility antigens on live tumor cells. *Indian. J. Exp. Biol.* 28, 1017, 1990.
105. Sarin, A., Haridas, V. and Saxena, R.K. MHC class I upregulation protects some but not all targets against LAK mediated lysis: Possible implications for the existence of LAK cell subpopulations using different recognition strategies. *Immunol. Lett.* 38, 59, 1993.
106. Sarin, A. and Saxena, R.K. Reversal of Interferon induced LAK resistance in two murine cell lines by exposure to acid pH. *Current Science.* 65, 705, 1994.
107. Saxena, R.K. and Adler, W.H. Modulation of natural cytotoxicity by alloantibodies. I. Alloantiserum enhancement of cytotoxicity of mouse spleen cells towards a human myeloid cell line. *J. Immunol.* 123, 846, 1979.
108. Saxena, R.K., Adler, W.H. and Nordin, A.A. Modulation of natural cytotoxicity by alloantibodies. IV. A comparative study of the activation of mouse spleen cell cytotoxicity by anti H-2 antisera, interferon and mitogens. *Cell. Immunol.* 63, 28, 1981.
109. Saxena, R.K., Saxena, Q.B. and Adler, W.H. Identity of effector cell participating in the reverse antibody dependent cell mediated cytotoxicity. *Immunology* 46, 459, 1982.
110. Saxena, R.K., Saxena, Q.B. and Adler, W.H. Properties and characterization of a rat spleen cell derived factor which induces resistance to NK cell lysis in YAC lymphoma cells. *J. Immunol.* 141, 1782, 1988.
111. Saxena, R.K., Saxena, Q.B. and Adler, W.H. Lectin induced cytotoxic activity in spleen cells from young and old mice. Age related changes in types of effector cells, lymphokine production and response. *Immunology.* 64, 457, 1988.
112. Schwartz, R.E. and Hiserodt, J.C. A surface structure involved in target-cell recognition by non-MHC-restricted cytotoxic effector cells. in *Leukocyte Typing IV. White Cell Differentiation Antigens.* Knapp, W., Dorken, B., Gilks, W.R., Rieber, E.P., Schmidt, R.E., Stein, R.H. and von dem Borne, A.E.G.Kr., Eds., Oxford University press, Oxford, 741, 1979.

113. **Schwartz, R.E. and Hiserodt, J.C.** The expression and functional involvement of laminin-like molecules in non-MHC-restricted cytotoxicity by human NK cells. *J. Immunol.* 141, 3318, 1988.
114. **Schwinzer, R.** Cluster report: CD45/CD45R, in *Leukocyte Typing IV. White Cell Differentiation Antigens.* Knapp, W., Dorken, B., Gilks, W.R., Rieber, E.P., Schmidt, R.E., Stein, R.H. and von dem Borne, A.E.G.Kr., Eds., Oxford University press, Oxford, 628, 1989.
115. **Shellani, G.R.** Studies on a gross virus induced lymphoma in the rat. V. Natural cytotoxic cells are non T cells. *Int. J. Cancer.* 19, 225, 1977.
116. **Shimizu, Y. and DeMars, R.** Demonstration by class I gene transfer that reduced susceptibility of human cells to natural killer cell mediated lysis is inversely correlated with HLA class I antigen expression. *Eur. J. Immunol.* 19, 447, 1989.
117. **Shimizu, Y., Weidman, E., Iwatsuki, S., Herberman, R.B. and Whiteside, T.** Characterization of human anti tumor reactive T cell clones obtained from tumor infiltrating lymphocytes in liver lymphocytes in liver metastasis gastric carcinoma. *cancer Res.* 51, 6153, 1991.
118. **Skibber, J.M., Lotze, M.T., Muul, L.M., Uppenkamp, I.K., Ross, W. and Rosenberg, S.A.** Human lymphokine activated killer cells: Further isolation and characterization of the precursor and effector cells. *Nat. Immun. Cell Growth Regul.* 6, 291, 1987.
119. **Sparrow, R.L., and McKenzie, I.F.C.** A function for human T200 in natural killer cytotoxicity. *Transplantation* 36, 166, 1983.
120. **Stein-Streilein, J., Bennet, M., Mann, D. and Kumar, V.** Natural killer cells in mouse lung : surface phenotype target preference and response to local influenza virus infection. *J. Immunol.* 131, 2699, 1983.
121. **Stern, A.S., Podlaski, F.J., Hulmes, J.D., Pan, Y.C.E., Quinn, P.M., Wolitzky, A.G., Familletti, P.C., Stremio, D.L., Tritt, T., Chizzonite, R. and Gately, M.K.** Purification to homogeneity and partial characterization of cytotoxic lymphocyte maturation factor from human B-lymphoblastoid cells. *Proc. Natl. Acad. Sci. U.S.A.* 87, 6808, 1990.
122. **Storkus, W.J. and Dawson, J.R.** B cell sensitivity to natural killing: correlation with target cell stage of differentiation and state of activation. *J. Immunol.* 136, 1542, 1986.

123. **Storkus, W.J., Howell, D.N., Salter, R.D., Dawson, J.R. and Cresswell, P.** NK susceptibility varies inversely with target cell class I HLA antigen expression. *J. Immunol.* 138, 1657, 1987.
124. **Storkus, W.J., Alexander, J., Payne, J.A., Dawson, J.R. and Cresswell, P.** Reversal of natural killing susceptibility in target cells expressing transfected class I HLA genes. *Proc. Natl. Acad. Sci. U.S.A.* 86, 2361, 1989.
125. **Stotter, H., Custer, M.C., Bolton, E.S., Guedez, L. and Lotze, M.T.** IL-7 induces human lymphokine-activated killer cell activity and is regulated by IL-4. *J. Immunol.* 146, 150, 1991.
126. **Strueli, M., Hall, L.R., Saga, Y., Schlossman, S.F. and Bolhuis, R.L.H.** Differential usage of three exons generates at least five different mRNAs encoding human human leukocyte common antigens. *J. Exp. Med.* 166, 1548, 1987.
127. **Sturmhofel, K. and Hammerling, G.J.** Reconstitution of H-2 class I expression by gene transfection decreases susceptibility to natural killer cells of an EL4 class I loss variant. *Eur. J. Immunol.* 20, 171, 1990.
128. **Suguwara, S., Abo, T. and Kumagai, K.** A simple method to eliminate the antigenicity of surface class I MHC molecules from the membrane of viable cells by acid treatment at pH 3.0. *J. Imm. Methods.* 100, 83, 1987.
129. **Tanaka, K., Yoshioka, T., Bieberich, C. and Jay, G.** Role of the major histocompatibility complex class I antigens in tumor growth and metastasis. *Annu. Rev. Immunol.* 6, 359, 1988.
130. **Tangiuchi, K., Karre, K. and Klein, G.** Lung colonization and metastasis by disseminated B16 melanoma cells: H-2 associated control at the level of the host and the tumor cell. *Int. J. Cancer* 36, 503, 1985.
131. **Targan, B., Britvan, L., Kendal, R., Vimadala, S. and Soll, A.** Isolation of spontaneous and interferon inducible natural killer cells from human colonic mucosa: lysis of lymphoid and autologous epithelial target cells. *Cell. Exp. Immunol.* 54, 14, 1983.
132. **Timonen, T., Ortaldo, J.R., and Herberman, R.B.** Characteristics of human large granular lymphocytes and relationship to natural killer and K cells. *J. Exp. Med.* 153, 583, 1981.

133. **Tutt, M.M., Kuziel, W.A., Hackett, J., Bennett, M., Tucker, P.W. and Kumar, V.** Murine natural killer cells do not express functional transcripts of alpha, beta or gamma chain genes of the T cell receptor. *J. Immunol.* 137, 2998, 1986.
134. **Twomey, J.J. and Kouttab, N.M.** The null cell compartment of the mouse spleen. *Cell. Immunol.* 63, 106, 1978.
135. **Vanky, F., Stubber, G., Willem, J., Sjowall, K., Larsson, B., Book, K., Ivert, T., Peteroffy, A. and Klein, E.** Importance of MHC antigen expression on solid tumors in the in vitro interaction with autologous blood lymphocytes. *Clin. Immunol. Immunother.* 27, 213, 1988.
136. **Van Rood, J.J., DeVries, R.R.P. and Bradely, B.A.** Genetics and biology of the HLA system, in the role of the Major Histocompatibility Complex in Immunobiology, Dorf, M.E., Ed., Garland, New York, 1981.
137. **Versteeg, R., Peltenburg, L.T.C., Plomp, A.C. and Schrier, P.I.** c-myc downregulates class I HLA expression in human melanomas. *EMBO. J.* 7, 1023, 1988.
138. **West, W.H., Tauer, K.W., Yanneill, J.R., Marshall, G.D., Orr, D.W., Thurmann, G.B. and Oldham, R.K.** Constant-infusion recombinant interleukin-2 in adoptive immunotherapy of advanced cancer. *N. Engl. J. Med.* 316, 898, 1987.
139. **Wiebke, E.A., Custer, M.C., Rosenberg, S.A. and Lotze, M.T.** Cytokines alter target cell susceptibility to lysis: Evaluation of non Major Histocompatibility Complex restricted effectors reveal differential effects on natural and lymphokine activated killing. *J. Biol. Resp. Modifiers.* 9, 113, 1990.
140. **Wright, S.C. and Bonavida, B.** Studies on the mechanism of natural killer (NK) cell-mediated cytotoxicity (CMC) I. Release of cytotoxic factors specific for NK-sensitive target cells (NKCF) during co-culture of NK effector cells with NK target cells. *J. Immunol.* 129, 433, 1982.
141. **Wright, S.C. and Bonavida, B.** Studies on the mechanism of natural killer cell mediated cytotoxicity. IV. Interferon induced inhibition of NK target cell susceptibility to lysis is due to a defect in their ability to stimulate release of natural killer cytotoxic factors (NKCF). *J. Immunol.* 130, 2965, 1983.

142. Yamaski, T., Klein, G., Ljunggren, H.G., Hoglund, P., Ohlen, C., Petersson, M.G.E. and Karre, K. Effects of dimethyl sulfoxide treatment on H-2 expression and susceptibility to NK or cytotoxic T lymphocyte-mediated lysis of the YAC-1 lymphoma and its β -microglobulin-deficient variant. J. Nat. Cancer Inst. 80, 263, 1988.
143. Yanagawa, E., Uchida, A. and Micksche, M. Natural cytotoxicity of lymphocytes from lymph nodes draining breast carcinoma and its augmentation by interferon and OK432. Cancer. Immunol. Immunother. 17, 1, 1984.
144. Yates, A.J., Icard-Liepkains, C., Sirinack, L.P., Stephens, R.E., Elder, P.J., Liepkains, V.A. and Whisler, R.L. Natural killer activity against cultured human neural tumor and fetal brain cells. Cell. Immunol. 90, 485, 1985.
145. Zanden, J., Werkmeister, J., McCarthy, W.H. and Hersey, P. Characterization of antigens recognized by natural killer cells in cell-culture supernatants. Br. J. Cancer 43, 5, 1981.
146. Zinkernagel, R.M. and Doherty, P.C. MHC-restricted cytotoxic T cells: studies on the biological role of polymorphic major transplantation antigens determining T-cell restriction-specificity, function and responsiveness. Adv. Immunol., 27, 51, 1979.