

## 6. REFERENCES

---

- Adams MD, Kelley JM, Gocayne JD, Dubnick M, Polymeropoulos MH, Xiao H, Merril CR, Wu A, Olde B, Moreno RF, Kerlavage AR, McCombie WR, Venter JC (1991) Complementary DNA sequencing: expressed sequence tags and human genome project. *Science* 252: 1651-1656
- Ahn S, Anderson JA, Sorrells ME, Tanksley SD (1993) Homoeologous relationships of rice, wheat and maize chromosomes. *Mol Gen Genet* 241: 483-490
- Akbari M, Wenzl P, Caig V, Carling J, Xia L, Yang S, Uszynski G, Mohler V, Lehmensiek A, Kuchel H, Hayden MJ, Howes N, Sharp P, Vaughan P, Rathmell B, Huttner E, Kilian A (2006) Diversity arrays technology (DArT) for high-throughput profiling of the hexaploid wheat genome. *Theor Appl Genet* 113: 1409-1420
- Akhunov ED, Akhunova AR, Dvorak J (2005) BAC libraries of *Triticum urartu*, *Aegilops speltoides* and *Ae tauschii*, the diploid ancestors of polyploid wheat. *Theor Appl Genet* 111: 1617-1622
- Akhunov ED, Goodyear AW, Geng S, Qi L-L, Echaliier B, Gill BS, Miftahudin, Gustafson JP, Lazo G, Chao S, Anderson OD, Linkiewicz AM, Dubcovsky J, La Rota M, Sorrells ME, Zhang D, Nguyen HT, Kalavacharla V, Hossain K, Kianian SF, Peng J, Lapitan NLV, Gonzalez-Hernandez JL, Anderson JA, Choi DW, Close TJ, Dilbirligi M, Gill KS, Walker-Simmons MK, Steber C, McGuire PE, Qualset CO, Dvorak J (2003) The organization and rate of evolution of wheat genomes are correlated with recombination rates along chromosome arms. *Genome Res* 13: 753-763
- Akkaya MS, Bhagwat AA, Cregan PB (1992) Length polymorphism of simple sequence repeat DNA in soybean. *Genetics* 132: 1131-1139
- Alonso-Blanco C, Goicoechea PG, Roca A, Alvarez E, Giraldez R (1994) Genetic mapping of cytological and isozyme markers on chromosome 1R, 3R, 4R and 6R of rye. *Theor Appl Genet* 88: 208-214
- Balyan HS, Gupta PK, Goyal A, Mohan A, Kumar S (2008) Integrated physical map of bread wheat containing 2,120 simple sequence repeat loci. *Adv Chromosome Sci* 3: 87-91
- Bandopadhyay R, Sharma S, Rustgi S, Singh R, Kumar A, Balyan HS, Gupta PK (2004) DNA polymorphism among 18 species of *Triticum-Aegilops* complex, using wheat EST-SSRs. *Plant Sci* 166: 349-356
- Barakat A, Carels N, Bernardi G (1997) The distribution of genes in the genomes of Graminae. *Proc Natl Acad Sci USA* 94: 6857-6861

- Barakat A, Matassi G, Bernardi G (1998) Distribution of genes in the genome of *Arabidopsis thaliana* and its implications for the genome organization of plants. *Proc Natl Acad Sci USA* 95: 10044-10049
- Bariana HS, Hydan MJ, Ahmed NU, Khatkar S, Conner RL, Wellings CR, Haley S, Sharp PJ, Laroche A (2002) Characterization of *Triticum vavilovi* - derived stripe rust resistance using, cytogenetic and molecular analysis and its marker assisted selection. *Theor Appl Genet* 104: 315-320
- Bennett MD, Leitch IJ (2005) Plant genome size research – -6
- Bennett MD, Leitch IJ (2011) Nuclear DNA amounts in angiosperms: targets, trends and tomorrow. *Ann Bot*, doi:10.1093/aob/mcq258
- Bennetzen JL (2000) Sequence analysis of plant nuclear genomes: microcolinearity and its many exceptions. *Plant Cell* 12: 1021-1030
- Berard A, LePaslier MC, Dradevet M, Exbravet-Vinson F, Bonnin I, Cenci A, Haudry A, Brunel D, Ravel C (2009) High-throughput single nucleotide polymorphism genotyping in wheat (*Triticum* spp.). *Plant Biotechnol J* 7: 364-374
- Biagetti M, Vitellozzi F, Ceoloni C (1999) Physical mapping of wheat-*Aegilops longissima* breakpoints in mildew-resistant recombinant lines using FISH with highly repeated and low-copy DNA probes. *Genome* 42: 1013-1019
- Bohn M, Utz HF, Melchinger AE (1999) Genetic similarities among winter wheat cultivars determined on the basis of RFLPs, AFLPs, and SSRs and their use for predicting progeny variance. *Crop Sci* 39: 228-237
- Bolot S, Abrouk M, Masood-Quraishi U, Stein N, Messing J, Feuillet C, Salse J (2009) The 'inner circle' of the cereal genome. *Curr Opin Plant Biol* 12: 1-7
- Bortiri E, Coleman-Derr D, Lazo GR, Anderson OD, Gu YQ (2008) The complete chloroplast genome sequence of *Brachypodium distachyon*: sequence comparison and phylogenetic analysis of eight grass plastomes. *BMC Res Notes* 1: 61
- Bossolini E, Wickler T, Knobel PA, Keller B (2007) Comparison of orthologous loci from small grass genomes *Brachypodium* and rice: implications for wheat genomics and grass genome annotation. *Plant J* 49: 704-717
- Boyko E, Kalendar R, Korzun V, Fellers J, Korol A, Schulman AH, Gill BS (2002) A high-density cytogenetic map of the *Aegilops tauschii* genome incorporating retrotransposons and defense-related genes: insights into cereal chromosome structure and function. *Plant Mol Biol* 48: 767-790
- Brunner S, Srichumpa P, Yahiaoui N, Keller B (2005) Positional cloning and evolution of powdery mildew resistance gene at *Pm3* locus of hexaploid wheat. In: *Plant and Animal Genome XIII Conference, Town & Country Convention Center, San Diego CA*, pp 73

- Bryan GJ, Collins AJ, Stephenson P, Orry A, Smith JB, Gale MD (1997) Isolation and characterization of microsatellites from hexaploid bread wheat. *Theor Appl Genet* 94: 557-563
- Cadalen T, Boeuf C, Bernard S, Bernard M (1997) An intervarietal molecular marker map in *Triticum aestivum* L. em Thell and comparison with a map from a wide cross. *Theor Appl Genet* 94: 367-377
- Cadalen T, Maitre B, Duranton N, Gay G, Gandon B, Chiquet V, Murigneux A, Dufour P, Sourdille P, Bernard M (2002) Potential of the deletion mapping system in wheat for the establishment of genetic map – physical map relationships through the chromosomal assignment of mapped markers. In: *Plant, Animal & Microbe Genomes X Conference, Town & Country Convention Center, San Diego CA*, pp 349
- Cardle L, Ramsay L, Milbourne D, Macaulay M, Marshall D, Waugh R (2000) Computational and experimental characterization of physically clustered simple sequence repeats in plants. *Genetics* 156: 847-854
- Catalan P, Olmstead RG (2000) Phylogenetic reconstruction of the genus *Brachypodium* P. Beauv. (Poaceae) from combined sequences of chloroplast *ndhF* gene and nuclear ITS. *Plant Syst Evol* 220: 1-19
- Catalan P, Ying S, Armstrong L, Draper J, Stace CA (1995) Molecular phylogeny of the grass genus *Brachypodium* P. Beauv. based on RFLP and RAPD analysis. *Bot J Linn Soc* 117: 263-280
- Chalmers KJ, Campbell AW, Kretschmer J, Karakousis A, Henschke PH, Pierens S, Harker N, Pallota M, Cornish GB, Shariflou MR, Rampling LR, McLauchlan A, Daggard G, Sharp P, Holton TA, Sutherland MW, Appels R, Langridge P (2001) Construction of three linkage maps in bread wheat, *Triticum aestivum*. *Aust J Agric Res* 52: 1089-1119
- Chang YL, Tao Q, Scheuring C, Meksem K, Zhang H-B (2001) An integrated map of *Arabidopsis thaliana* for functional analysis of its genome sequence. *Genetics* 159: 1231-1242
- Chantret N, Cenci A, Sabot F, Anderson O, Dubcovsky J (2004) Sequencing of the *Triticum monococcum* *Hardness* locus reveals good microcolinearity with rice. *Mol Genet Genomics* 271: 377-386
- Chao S, Sharp PJ, Worland AJ, Koebner RMD, Gale MD (1989) RFLP-based genetic maps of homoeologous group 7 chromosomes. *Theor Appl Genet* 78: 495-504
- Charlesworth D, Charlesworth B, McVean GAT (2000) Genome sequences and evolutionary biology, a two-way interaction. *Trends Ecol Evol* 16: 235-242
- Chen M, Presting G, Barbazuk WB, Goicoechea JL, Blackmon B, Fang G, Kim H, Frisch D, Yu Y, Sun S, Higingbottom S, Phimphilai J, Phimphilai D, Thumond S, Gaudette B, Li P,

- Liu J, Hatfield J, Main D, Farrar K, Hendrson C, Barnett L, Costa R, Williams B, Walser S, Atkins M, Hall C, Budiman MA, Tomkins JP, Luo M, Bancroft I, Salse J, Regad F, Mohapatra T, Singh NK, Tyagi AK, Soderlund C, Dean RA, Wing RA (2002) An integrated physical and genetic map of the rice genome. *Plant Cell* 14: 537-545
- Chen, JM Gustafson JP (1995) Physical mapping of restriction fragment length polymorphisms (RFLPs) in homoeologous group 7 chromosomes of wheat by *in-situ* hybridization. *Heredity* 75: 225-233
- Cheng ZK, Buell CR, Wing RA, Jiang J (2002) Resolution of fluorescence *in-situ* hybridization mapping on rice mitotic prometaphase chromosomes, meiotic pachytene chromosomes and extended DNA fibers. *Chromosome Res* 10: 379-387
- Cheung WY, Chao S, Gale MD (1991) Long-range physical mapping of the  $\alpha$ -amylase 1 (*Amy-1*) loci on homoeologous group 6 chromosomes of wheat and rye. *Mol Gen Genet* 229: 373-379
- Cho YG, Ishii T, Temnykh S, Chen X, Lipovich L, McCouch SR, Park WD, Ayres N, Cartinhour S (2000) Diversity of microsatellites derived from genomic libraries and GenBank sequences in rice (*Oryza sativa* L.). *Theor Appl Genet* 100: 713-722
- Christiansen MJ, Andersen SB, Ortiz R (2002) Diversity changes in an intensively bread wheat germplasm during the 20<sup>th</sup> century. *Mol Breed* 9: 1-11
- Condit R, Hubbel S (1991) Abundance and DNA sequence of two-base repeat regions in tropical tree genomes. *Genome* 34: 66-71
- Cone K, McMullen M, Bi IV, Davis G, Yim Y-S, Gardiner J, Polacco M, Sanchez-Villeda H, Fang Z, Schroeder SG, Havermann SA, Bowers JE, Paterson AH, Soderlund CA, Engler FW, Wing RA, Coe Jr EH (2002) Genetic, physical and informatics resources for maize: on the road to an integrated map. *Plant Physiol* 130: 1598-1605
- Conley EJ, Nduati V, Gonzalez-Hernandez JL, Mesfin A, Trudeau-Spanjers M, Chao S, Lazo GR, Hummel DD, Anderson OD, Qi LL, Gill BS, Echaliier B, Linkiewicz AM, Dubscovsky J, Akhunov ED, Dvorak J, Peng JH, Lapitan NLV, Pathan MS, Nguyen HT, Ma X-F, Miftahudin, Gustafson JP, Greene RA, Sorrells ME, Hossain KG, Kalavacharla V, Kianian SF, Sidhu D, Dilbirligi M, Gill KS, Choi DW, Fenton RD, Close TJ, McGuire PE, Qualset CO, Anderson JA (2004) A 2600-locus chromosome bin map of wheat homoeologous group 2 reveals interstitial gene-rich islands and colinearity with rice. *Genetics* 168: 625-637
- Cordoba JM, Chavarro C, Schlueter JA, Jackson SA, Blair MW (2010) Integration of physical and genetic maps of common bean through BAC-derived microsatellite markers. *BMC Genomics* 11: 436-446

- Delaney DE, Nasuda S, Endo TR, Gill BS, Hulbert SH (1995a) Cytologically-based physical maps of the group 2 chromosomes of wheat. *Theor Appl Genet* 91: 568-573
- Delaney DE, Nasuda S, Endo TR, Gill BS, Hulbert SH (1995b) Cytologically-based physical maps of the group 3 chromosomes of wheat. *Theor Appl Genet* 91: 780-782
- Devos KM (2010) Grass genome organization and evolution. *Curr Opin Plant Biol* 13: 139-145
- Devos KM, Atkinson MD, Chinoy CN, Gale MD (1992) RFLP-based genetic maps of the homoeologous group-3 chromosomes of wheat and rye. *Theor Appl Genet* 83: 931-939
- Devos KM, Gale MD (1993) Extended genetic maps of the homoeologous group 3 chromosomes of wheat, rye and barley. *Theor Appl Genet* 85: 649-652
- Devos KM, Gale MD (2000) Genome relationship: the grass model in current research. *Plant Cell* 12: 637-646
- Devos KM, Ma J, Potaro AC, Pratt LH, Bennetzen JL (2005) Analysis and mapping of randomly chosen bacterial artificial chromosome clones from hexaploid bread wheat. *Proc Natl Acad Sci USA* 52: 19242-19248
- Devos KM, Moore G, Gale MD (1995) Conservation of marker synteny during evolution. *Euphytica* 85: 367-372
- Dillon SL, Lawrence PK, Henry RJ (2005) The new use of *Sorghum bicolor* derived SSR markers to evaluate genetic diversity in seventeen Australian *Sorghum* species. *Plant Genet Res* 3: 19-28
- Dolezel J, Kubalaková M, Paux E, Bartos J, Feuillet C (2007) Chromosome-based genomics in the cereals. *Chromosome Res* 15: 51-66
- Dolezel J, Lysak MA, Kubalaková M, Simkova H, Macas J (2001) Sorting of plant chromosomes. In: Darzynkiewicz Z, Crissman HA, Robinson JP (eds) *Methods in Cell Biology*, 3<sup>rd</sup> edition, Part B. Academic Press, San Diego, pp 3-31
- Dong F, Song J, Naess SK, Helgeson JP, Gebhardt C, Jiang J (2000) Development and applications of a set of chromosome specific cytogenetic DNA markers in potato. *Theor Appl Genet* 101: 1001-1007
- Dong FG, Miller JT, Jackson SA, Wang GL, Ronald PC (1998) Rice (*Oryza sativa*) centromeric regions consist of complex DNA. *Proc Natl Acad Sci USA* 95: 8135-8140
- Draper J, Mur LAJ, Jenkins G, Ghosh-Biswas GC, Bablak P, Hasterok R, Routledge APM (2001) *Brachypodium distachyon*, a new model system for functional genomics in grasses. *Plant Physiol* 127: 1539-1555
- Dweikat I, Ohm H, Patterson F, Cambron S (1997) Identification of RAPD markers for 11 hessian fly resistance genes in wheat. *Theor Appl Genet* 94: 419-423

- Edwards KJ, Barker JHA, Daly A, Jones C, Carp A (1996) Microsatellite libraries enriched for several microsatellite sequences in plants. *Biotechniques* 20: 758-760
- Elouafi I, Nachit MM (2004) A genetic linkage map of the Durum  $\times$  *Triticum dicoccoides* backcross population based on SSRs and AFLP markers, and QTL analysis for milling traits. *Theor Appl Genet* 108: 401-413
- Endo TR (1988) Chromosome mutations induced by gametocidal chromosome in common wheat. In: Miler TE, Koebner RMD (eds) *Proceedings 7<sup>th</sup> International Wheat Genetics Symposium*, Cambridge UK, pp 259-265
- Endo TR, Gill BS (1996) The deletion stocks of common wheat. *J Hered* 87: 295-307
- Erayman M, Sandhu D, Sidhu D, Dilbirligi M, Baenziger PS, Gill KS (2004) Demarcating the gene-rich regions of the wheat genome. *Nucl Acids Res* 32: 3546-3565
- Eujayl I, Sorrells ME, Baum M, Wolters P, Powell W (2001) Assessment of genotypic variation among cultivated durum wheat based on EST-SSRs and genomic SSRs. *Euphytica* 119: 39-43
- Eujayl I, Sorrells ME, Baum M, Wolters P, Powell W (2002) Isolation of EST-derived microsatellite markers for genotyping the A and B genomes of wheat. *Theor Appl Genet* 104: 339-407
- Ewing RM, Kahla AB, Poirot O, Lopez F, Audic S, Calverie JM (1999) Large-scale statistical analyses of rice ESTs reveal correlated patterns of gene expression. *Genome Res* 9: 950-959
- Faris JD, Fellers JP, Brooks SA, Gill BS (2003) A bacterial artificial chromosome contig spanning the major domestication locus *Q* in wheat and identification of a candidate gene. *Genetics* 164: 311-321
- Faris JD, Haen KM, Gill BS (2000) Saturation mapping of a gene-rich recombination hot spot region in wheat. *Genetics* 154: 823-835
- Faris JD, Zhang Z, Fellers JP, Gill BS (2008) Micro-colinearity between rice, *Brachypodium* and *T. monococcum* at the wheat domestication locus *Q*. *Funct Integr Genomics* 8: 149-164
- Fernandes J, Brendel V, Gai X, Lal S, Chandler VL, Elumalai RP, Galbraith DW, Pierson EA, Walbot V (2002) Comparison of RNA expression profile based on maize expressed sequence tag frequency analysis and micro-array hybridization. *Plant Physiol* 128: 896-910
- Feuillet C, Keller B (1999) High gene density is conserved at syntenic loci of small and large grass genomes. *Proc Natl Acad Sci USA* 96: 8265-8270
- Feuillet C, Messmer M, Schachermayr G, Keller B (1995) Genetic and physical characterization of the *LRI* leaf rust resistance locus in wheat (*Triticum aestivum* L.). *Mol Gen Genet* 248: 553-562

- Feuillet C, Travella S, Stein N, Albar L, Nublat A, Keller B (2003) Map-based isolation of the leaf rust disease resistance gene *Lr10* from the hexaploid wheat (*Triticum aestivum* L.) genome. *Proc Natl Acad Sci USA* 100: 15253-15258
- Flavell AJ, Bolshakov VN, Booth A, Jing R, Russell J, Ellis TH, Isaac P (2003) A microarray-based high throughput molecular marker genotyping method: the tagged microarray marker (TAM) approach. *Nucleic Acids Res* 31: e115
- Flavell RB, Bennett MD, Smith JB, Smith DB (1974) Genome size and the proportion of repeated nucleotide sequence DNA in plants. *Biochem Genet* 12: 257-269.
- Florijn RJ, Bonden LAJ, Vrolijk H, Wiegant J, Vaandrager JW, Baas F, den Dunnen JT, Tanke HJ, van Ommen GJB, Raap AK (1995) High-resolution DNA mapping and colour bar-coding of large genes. *Hum Mol Genet* 5: 831-836
- Foote T, Roberts M, Kurata N, Sasaki T, Moore G (1997) Detailed comparative mapping of cereal chromosome regions corresponding to the *Ph1* locus in wheat. *Genetics* 147: 801-807
- Fransz PF, Alfonso-Blanco C, Liharska TB, Peeters AJM, Zabel P, de Jong JH (1996) High-resolution physical mapping in *Arabidopsis thaliana* and tomato by fluorescence *in situ* hybridization to extended DNA fibers. *Plant J* 9: 421-430
- Fransz PF, Armstrong S, de Jong JH, Parnell LD, Drunen CV, Dean C, Zabel P, Bisseling T, Jones GH (2000) Integrated cytogenetic map of chromosome arm 4S of *Arabidopsis thaliana*: structural organization of heterochromatic knob and centromeric region. *Cell* 100: 367-376
- Freeling M (2001) Grasses as a single genetic system. Reassessment 2001. *Plant Physiol* 125: 1191-1197
- Fu Y-B, Peterson GW, Yu J-K, Gao L, Jia J, Richards KW (2005) Impact of plant breeding on genetic diversity of the Canadian hard red spring wheat germplasm as revealed by EST-derived SSR markers. *Theor Appl Genet* 112: 1239-1247
- Furuta Y, Nishikawa K (1991) Variation in nuclear and individual chromosomal DNA content and its role in evolution of plants. In: Gupta PK, Tsuchiya T (eds) *Chromosome Engineering in Plants: Genetics, Breeding, Evolution*, Part A. Elsevier, New York pp 71-85
- Gadaleta A, Giancaspro A, Giove SL, Zacheo S, Mangini G, Simeone R, Signorile A, Blanco A (2009) Genetic and physical mapping of new EST-derived SSRs on the A and B genome chromosomes of wheat. *Theor Appl Genet* 118: 1015-1025
- Galiba G, Vagujfalvi A, Li C, Soltesz A, Dubcovsky J (2009) Regulatory genes involved in the determination of frost tolerance in temperate cereals. *Plant Sci* 176: 12-19
- Gao LF, Jing RL, Huo NX, Li Y, Li XP, Zhou HR, Chang XP, Tang JF, Ma ZY, Jia JZ (2004) One hundred and one new microsatellite loci derived from ESTs (EST-SSRs) in bread wheat. *Theor Appl Genet* 108: 1392-1400

- Gao LF, Tang J, Li H, Jia J (2003) Analysis of microsatellites in major crops assessed by computational and experimental approaches. *Mol Breed* 12: 245-261
- Gao W, Chen J, Yu JZ, Raska D, Kohel RJ, Womack JE, Stelly DM (2004) Wide-cross whole-genome radiation hybrid mapping of cotton (*Gossypium hirsutum* L.). *Genetics* 167: 1317-1329
- Garvin DF (2007) *Brachypodium*: a new monocot model plant system emerges. *J Sci Food Agric* 87: 1177-1179
- Garvin DF, McKenzie N, Vogel JP, Mockler TC, Blankenheim ZJ, Wright J, Cheema JJS, Dicks J, Huo N, Hayden DM, Gu Y, Tobias C, Chang JH, Chu A, Trick M, Michael TP, Bevan MW, Snape JW (2010) An SSR-based genetic map of the model grass *Brachypodium distachyon*. *Genome* 53: 1-13
- Gill BS, Friebe B (2002) Cytogenetics, phylogeny and evolution of cultivated wheats. In: Curtis BS, Rajaram S, Macpherson HG (eds) *Bread Wheat Improvement and Production*. FAO Plant Protection and Production Series 30: 71-88
- Gill BS, Friebe B, Endo TR (1991) Standard karyotype and nomenclature system for description of chromosome bands and structural aberrations in wheat (*Triticum aestivum*). *Genome* 34: 830-839
- Gill KS, Gill BS, Endo TR (1993a) A chromosome region-specific mapping strategy reveals gene-rich telomeric ends in wheat. *Chromosoma* 102: 374-381
- Gill KS, Gill BS, Endo TR, Mukai Y (1993b) Fine physical mapping of *Ph1*, a chromosome pairing regulator gene in polyploid wheat. *Genetics* 134: 1231-1236
- Gill KS, Gill BS, Endo TR, Boyko EV (1996a) Identification and high-density mapping of gene-rich regions in chromosome group 5 of wheat. *Genetics* 143: 1001-1012
- Gill KS, Gill BS, Endo TR, Taylor T (1996b) Identification and high-density mapping of gene-rich regions in chromosome group 1 of wheat. *Genetics* 144: 1883-1891
- Gomez MI, Islam-Faridi MN, Woo SS, Schertz KF, Czeschin Jr D, Zwick MS, Sing RA, Stelly DM, Price HJ (1997) FISH of a maize *sh2* selected sorghum BAC to chromosomes of *Sorghum bicolor*. *Genome* 40: 475-478
- Goss SJ, Harris H (1975) New method for mapping genes in human chromosomes. *Nature* 255: 680-684
- Goyal A (2008) Physical mapping of SSR markers in bread wheat. Ph. D. Thesis, Department of Genetics and Plant Breeding, Ch. Charan Singh University, Meerut, India.
- Goyal A, Bandopadhyay R, Sourdille P, Endo TR, Balyan HS, Gupta PK (2005) Physical molecular maps of wheat chromosomes. *Funct Integr Genomics* 5: 260-263

- Griffiths S, Sharp R, Foote TN, Bertin I, Wanous M, Reader S, Colas I, Moore G (2006) Molecular characterization of *Ph1* as a major chromosome pairing locus in polyploid wheat. *Nature* 439: 749-752
- Gu YQ, Ma Y, Huo N, Vogel JP, You FM, Lazo GR, Nelson WM, Soderlund C, Dvorak J Anderson OD, Luo M-C (2009) A BAC-based physical map of *Brachypodium distachyon* and its comparative analysis with rice and wheat. *BMC Genomics* 10: 496-508
- Gupta PK, Balyan HS, Edwards KJ, Isaac P, Korzun V, Roder MS, Gautier MF, Joudrier P, Schlatter AR, Dubcovsky J, De la Pena RC, Khairallah M, Penner G, Hayden MJ, Sharp P, Keller B, Wang RCC, Hardouin JP, Jack P, Leroy P (2002) Genetic mapping of 66 new microsatellite (SSR) loci in bread wheat. *Theor Appl Genet* 105: 413-422
- Gupta PK, Balyan HS, Goyal A, Mohan A, Kumar S (2008c) An integrated physical map of 2,072 SSR loci (gSSRs and EST-SSRs) in bread wheat. In: Appels R, Estwood R, Lagudah E, Langridge P, Mackay M, McIntyre L, Sharp P (eds) *Proceedings 11<sup>th</sup> International Wheat Genetic Symposium, Brisbane, Australia*, pp 333-335
- Gupta PK, Balyan HS, Sharma PC, Ramesh B (1996) Microsatellites in plants: a new class of molecular markers. *Curr Sci* 70: 45-54
- Gupta PK, Mir RR, Mohan A, Kumar J (2008b) Wheat genomics: present status and future prospects. *Int J Plant Genomics*, doi:10.1155/2008/896451
- Gupta PK, Rustgi S (2004) Molecular markers from the transcribed/expressed region of the genome in higher plants. *Funct Integr Genomics* 4: 139-162
- Gupta PK, Rustgi S, Mir RR (2008a) Array-based high-throughput DNA markers for crop improvement. *Heredity* 101: 1-14
- Gupta PK, Rustgi S, Sharma S, Singh R, Kumar N, Balyan HS (2003) Transferable EST-SSR markers for the study of polymorphism and genetic diversity in bread wheat. *Mol Genet Genomics* 270: 315-323
- Gupta PK, Varshney RK (2000) The development and use of microsatellite markers for genetic analysis and plant breeding with emphasis on bread wheat. *Euphytica* 113: 163-185
- Gupta PK, Varshney RK (2004) Cereal genomics: An overview. In: Gupta PK, Varshney RK (eds) *Cereal Genomics*, Kluwer Academic Publishers, Dordrecht, The Netherlands, pp 24-25
- Gupta PK, Varshney RK, Sharma PC, Ramesh B (1999) Molecular markers and their applications in wheat breeding. *Plant Breed* 118: 369-390
- Gupta SK, Charpe A, Koul S, Prabhu KV, Haque QMR (2005) Development and validation of molecular markers linked to an *Aegilops umbellulata*-derived leaf rust resistance gene, *Lr9*, for marker-assisted selection in bread wheat. *Genome* 48: 823-830
- Gustafson JP, McGuire PE, Qualset CO (2004) Genomic resources: moving wheat into the mainstream in plant genomics. *Genetics* 168: 583-584

- Guyomarc'h H, Sourdille P, Charmet G, Edwards KJ, Bernard M (2002) Characterization of polymorphic microsatellite markers from *Aegilops tauschii* and transferability to the D genome of bread wheat. *Theor Appl Genet* 104: 1164-1172
- Haaf T, Ward DC (1994) High resolution ordering of YAC contigs using extended chromatin and chromosomes. *Hum Mol Genet* 4: 629-633
- Hackauf B, Rudd S, Van der Voort JR, Miedaner T, Wehling P (2009) Comparative mapping of DNA sequences in rye (*Secale cereale* L.) in relation to the rice genome. *Theor Appl Genet* 118: 371-384
- Hackauf B, Wehling P (2002) Identification of microsatellite polymorphisms in an expressed portion of the rye genome. *Plant Breed* 121: 17-25
- Han Z, Wang C, Song X, Guo W, Gou J, Li C, Chen X, Zhang T (2006) Characteristics, development and mapping of *Gossypium hirsutum* derived EST-SSRs in allotetraploid cotton. *Theor Appl Genet* 106: 430-439
- Hanson RE, Zwick MS, Choi S, Islam-Faridi MN, McKnight TD, Wring RA, Price HJ, Stelly DM (1995) Fluorescent *in situ* hybridization of a bacterial artificial chromosome. *Genome* 38: 646-651
- Harker N, Rampling LR, Shariflou MR, Hayden MJ, Holton TA, Morell MK, Sharp PJ, Edwards KJ (2001) Microsatellites as markers for Australian wheat improvement. *Aust J Agric Res* 52: 1121-1130
- Harper LC, Cande WZ (2000) Mapping a new frontier; development of integrated cytogenetic maps in plants. *Funct Integr Genomics* 1: 89-98
- Hartl L, Mohler V, Zeller FJ, Hsam SKL, Schweizer G (1999) Identification of AFLP markers closely linked to the powdery mildew resistance genes *Pm1c* and *Pm4a* in common wheat. *Genome* 42: 322-329
- Hasterok R, Marasek A, Donnison IS, Armstead I, Thomas A, King IP, Wolny E, Idziak D, Draper J, Jenkins G (2006) Alignment of the genomes of *Brachypodium distachyon* and temperate cereals and grasses using bacterial artificial chromosome landing with fluorescence *in situ* hybridization. *Genetics* 173: 349-362
- Hayden MJ, Tabone T, Mather DE (2009) Development and assessment of simple PCR markers for SNP genotyping in barley. *Theor Appl Genet* 119: 939-951
- Hazen PS, Leroy P, Ward R (2002) AFLP in *Triticum aestivum* L. Patterns of diversity and genome distribution. *Euphytica* 125: 89-102
- Hodgkin J, Plasterk RHA, Waterston RH (1995) The nematode *Caenorhabditis elegans* and its genome. *Science* 270: 410-414
- Hohmann U, Endo TR, Gill KS, Gill BS (1994) Comparison of genetic and physical maps of group 7 chromosomes from *Triticum aestivum*. *Mol Gen Genet* 245: 644-653

- Hohmann U, Endo TR, Herrmann RG, Gill BS (1995a) Characterization of deletions in common wheat induced by an *Aegilops cylindrica* chromosome: detection of multiple chromosome rearrangements. *Theor Appl Genet* 91: 611-617
- Hohmann U, Graner A, Endo TR, Gill BS, Herrmann RG (1995b) Comparison of wheat physical maps with barley linkage maps for group 7 chromosomes. *Theor Appl Genet* 91: 618-626
- Holton TA, Christopher JT, McClure L, Harker N, Henry RJ (2002) Identification and mapping of polymorphic SSR markers from expressed sequences of barley and wheat. *Mol Breed* 9: 63-71
- Hossain GK, Riera-Lizararu O, Kalavacharla V, Vales MI, Maan SS, Kianian SF (2004a) Radiation hybrid mapping of the species cytoplasm specific (*scs<sup>ae</sup>*) gene in wheat. *Genetics* 168: 415-423
- Hossain GK, Riera-Lizararu O, Kalavacharla V, Vales MI, Maan SS, Kianian SF (2004b) Molecular cytogenetics characterization of an alloplasmic durum wheat line with a portion of chromosome 1D of *Triticum aestivum* carrying *scs<sup>ae</sup>* gene. *Genome* 47: 206-214
- Hossain KG, Kalavacharla V, Lazo GR, Hegstad J, Wentz MJ, Kianian PMA, Simons K, Gehlhar S, Rust KL, Syamala RR, Obeori K, Bhamidimarri S, Karunadharm P, Chao S, Anderson D, Qi LL, Echaliier B, Gill BS, Linkiewicz AM, Ratnasiri A, Dubcovsky J, Akhunov ED, Dvorak J, Miftahudin, Ross K, Gustafson JP, Radhawa HS, Dilbirli M, Gill KS, Peng JH, Lapitan NLV, Greene RA, Bermudez-Kandianis CE, Sorrells ME, Feril O, Pathan MS, Nguyen HT, Gonzalez-Hernandez JL, Conley EJ, Anderson JA, Choi DW, Fenton, Close TJ, McGuire PE, Qualset CO, Kianian SF (2004c) A chromosome bin map of 2148 expressed sequence tag loci of wheat homoeologous group 7. *Genetics* 168: 687-699
- Hsaio C, Chatterton NJ, Asay KH, Jensen KB (1994) Phylogenetic relationships of 10 grass species: an assessment of phylogenetic utility of the internal transcribed spacer region in nuclear ribosomal DNA in monocots. *Genome* 37: 112-120
- Huang L, Brooks SA, Li W, Fellers JP, Trick HN, Gill BS (2003) Map-based cloning of leaf rust resistance gene *Lr21* from the large and polyploid genome of bread wheat. *Genetics* 164: 655-664
- Huang XQ, Borner A, Roder MS, Ganai MW (2002) Assessing genetic diversity of wheat (*Triticum aestivum* L.) germplasm using microsatellite markers. *Theor Appl Genet* 105: 699-707
- Huang XQ, Coster H, Ganai MW, Roder MS (2003) Advanced backcross QTL analysis for the identification of quantitative trait loci alleles from wild relatives of wheat (*Triticum aestivum* L.). *Theor Appl Genet* 106: 1379-1389

- Huang XQ, Hsam SLK, Zeller FJ, Wenzel W, Mohler V (2000) Molecular mapping of the powdery mildew resistance gene *Pm24* and marker validation for molecular breeding. *Theor Appl Genet* 101: 407-414
- Huo N, Gu YQ, Lazo GR, Vogel JP, Coleman-Derr D, Luo M-C, Thilmony R, Garvin DF, Anderson OD (2006) Construction and characterization of two BAC libraries from *Brachypodium distachyon*, a new model for grass genomics. *Genome* 49: 1099-1108
- Huo N, Lazo GR, Vogel JP, You FM, Ma Y, Hayden DM, Coleman-Derr D, Hill TA, Dvorak J, Anderson OD, Luo MC, Gu YQ (2008). The nuclear genome of *Brachypodium distachyon*: analysis of BAC ends sequences. *Funct Integr Genomics* 8: 135-147
- Huo N, Vogel JP, Lazo GR (2009) Structural characterization of *Brachypodium* genome and its syntenic relationship with rice and wheat. *Plant Mol Biol* 70: 47-61
- International Rice Genome Sequencing Project (2005) The map-based sequence of the rice genome. *Nature* 436: 793-800
- Isidore E, Scherrer B, Chalhoub B, Feuillet C, Keller B (2005) Ancient haplotypes resulting from extensive molecular rearrangements in the wheat A genome have been maintained in species of three different ploidy levels. *Genome Res* 15: 526-536
- Jackson SA, Wang ML, Goodman HM, Jiang J (1998) Application of fiber-FISH in physical mapping of *Arabidopsis thaliana*. *Genome* 41: 566-572
- Jackson SA, Zhang P, Chen WP, Phillips RL, Friebe B, Muthukrishnan S, Gill BS (2001) High-resolution structural analysis of biolistic transgene integration into the genome of wheat. *Theor Appl Genet* 103: 56-62
- Janda J, Bartos J, Safar J, Kubalaková M, Valarik M, Cihalikova J, Simkova H, Caboche M, Sourdille P, Bernard M, Chalhoumb B, Dolezel J (2004) Construction of a subgenomic BAC library specific for chromosomes 1D, 4D and 6D of hexaploid wheat. *Theor Appl Genet* 109: 1337-1345
- Jiang J, Gill BS (1994) Non-isotopic *in situ* hybridization and plant genome mapping: the first 10 years. *Genome* 37: 717-725
- Jiang J, Gill BS (2006) Current status and the future of fluorescence *in situ* hybridization (FISH) in plant genome research. *Genome* 49: 1-12
- Jiang J, Gill BS, Wang G-L, Ronald PC, Ward D (1995) Metaphase and interphase fluorescence *in situ* hybridization mapping of the rice genome with bacterial artificial chromosomes. *Proc Natl Acad Sci USA* 92: 4487-4491
- Joshi AK, Mishra B, Chatrath R, Ferrara GO, Singh RP (2007) Wheat improvement in India: present status, emerging challenges and future prospects. *Euphytica* 157: 431-446
- Kalavacharla V, Hossain K, Gu Y, Riera-Lizarazu O, Isabel-Vales M (2006) High-resolution radiation hybrid map of wheat chromosome 1D. *Genetics* 173: 1089-1099

- Kam-Morgan LMW, Gill BS, Muthukrishnan S (1989) DNA restriction fragment polymorphisms: a strategy for genetic mapping of D genome of wheat. *Genome* 32: 724-732
- Kantety RV, La Rota M, Matthews DE, Sorrells ME (2002) Data mining for simple sequence repeats in expressed sequence tags from barley, maize, rice, sorghum and wheat. *Plant Mol Biol* 48: 501-510
- Kawahara T (2009) Molecular phylogeny among *Triticum-Aegilops* species and of the tribe Triticeae. *Breed Sci* 59: 499-504
- Kellogg EA (2001) Evolutionary history of the grasses. *Plant Physiol* 125: 1198-1205
- Khlestkina EK, Pshenichnikova TA, Röder MS, Salina EA, Arbusova VS, Börner A (2006a) Comparative mapping of genes for glume colouration and pubescence in hexaploid wheat (*Triticum aestivum* L.). *Theor Appl Genet* 113: 801-807
- Khlestkina EK, Salina EA (2006b) SNP markers: methods of analysis, ways of development and comparison on an example of common wheat. *Russian J Genet* 42: 585-594
- Khlestkina EK, Than MHM, Pestsova EG, Roder MS, Malyshev SV, Korzun V, Börner A (2004) Mapping of 99 new microsatellite-derived loci in rye (*Secale cereale* L.) including 39 expressed sequence tags. *Theor Appl Genet* 109: 725-732
- Khush GS, Rick CM (1967) Studies on the linkage map of chromosome 4 of tomato and on the transmission of induced deficiencies. *Genetics* 38:74-94
- Khush GS, Rick CM (1968) Cytogenetic analysis of the tomato genome by means of induced deficiencies. *Chromosoma* 23: 452-484
- Kihara H (1954) Considerations on the evolution and distribution of *Aegilops* species based on the analyser method. *Cytologia* 19: 336-357
- Kim H, San Miguel P, Nelson W, Collura K, Wissotski M, Walling JG, Kim JP, Jackson SA, Soderlund C, Wing RA (2007) Comparative physical mapping between *Oryza sativa* (AA genome type) and *O. punctata* (BB genome type). *Genetics* 176: 379-390
- Kim J-S, Childs KL, Islam-Faridi MN, Menz MA, Klein RR (2002) Integrated karyotyping of sorghum by *in situ* hybridization of landed BACs. *Genome* 45: 402-412
- Klein P, Klein RR, Cartinhour S, Ulanich P, Dong J, Obert J, Morishige DT, Schlueter S, Childs K, Ale M, Mullet JE (2000) A high-throughput AFLP-based method for constructing integrated genetic and physical maps: progress toward a sorghum genome map. *Genome Res* 10: 789-807
- Korzun V, Ebmeyer E (2003) Molecular markers and their applications in wheat breeding. In: Pogna NE, Romano M, Pogna EA, Galterio G (eds) *Proceedings 10<sup>th</sup> International Wheat Genetics Symposium, Istituto Sperimentale per la Cerealicoltura, Rome, Italy, Vol 1*, pp 140-143

- Korzun V, Roder MS, Wendehake K, Pasqualone A, Lotti C, Ganal MW, Blanco A (1999) Integration of dinucleotide microsatellites from hexaploid bread wheat into a genetic linkage map of durum wheat. *Theor Appl Genet* 98: 1202-1207
- Korzun V, Röder MS, Worland AJ, Borner A (1997a) Application of microsatellite markers to distinguish inter-varietal chromosome substitution lines of wheat (*Triticum aestivum* L.). *Euphytica* 95: 149-155
- Korzun V, Röder MS, Worland AJ, Borner A (1997b) Intra chromosomal mapping of genes for dwarfing (*Rht12*) and vernalization response (*Vrn1*) in wheat by using RFLP and microsatellite markers. *Plant Breed* 116: 227-232
- Kota R, Gill KS, Gill BS, Endo TR (1993) A cytogenetically based physical map of chromosome 1B in common wheat. *Genome* 36: 548-554
- Kota R, Rudd S, Facius A, Kolesov G, Thiel T, Zhang H, Stein N, Mayer K, Graner A (2003) Snipping polymorphisms from large EST collections in barley (*Hordeum vulgare* L.). *Mol Genet Genomics* 270: 24-33
- Kota R, Varshney RK, Theil T, Dehmer KJ, Graner A (2001) Generation and Comparison of EST-Derived SSRs and SNPs in barley (*Hordeum Vulgare* L.). *Hereditas* 135: 145-151
- Kulwal P, Kumar N, Kumar A, Gupta RK, Balyan HS, Gupta PK (2005a) Gene networks in hexaploid wheat: interacting quantitative trait loci for grain protein content. *Funct Integr Genomics* 5: 254-259
- Kulwal PL, Kumar N, Gaur A, Khurana P, Khurana JP, Tyagi AK, Balyan HS, Gupta PK (2005b) Mapping of a major QTL for pre-harvest sprouting tolerance on chromosome 3A in bread wheat. *Theor Appl Genet* 111: 1052-1059
- Kulwal PL, Singh R, Balyan HS, Gupta PK (2004) Genetic basis of pre-harvest sprouting tolerance using single-locus and two-locus QTL analyses in bread wheat. *Funct Integr Genomics* 4: 94-101
- Kumar A, Kumar J, Singh R, Garg T, Chhuneja P, Balyan HS, Gupta PK (2009) QTL analysis for grain colour and pre-harvest sprouting in bread wheat. *Plant Sci* 117:114-122
- Kumar N, Kulwal PL, Gour A, Khurana P, Khurana JP, Tyagi AK, Balyan HS, Gupta PK (2006) QTL analysis for grain weight in bread wheat. *Euphytica* 151: 135-144
- Kumar S, Mohan A, Balyan HS, Gupta PK (2009) Orthology between genomes of *Brachypodium*, wheat and rice. *BMC Res Notes* 2: 93
- Kunzel G, Korzun L, Meister A (2000) Cytologically integrated physical restriction fragment length polymorphism maps for the barley genome based on translocation breakpoints. *Genetics* 154: 397-412

- Kynast RG, Okagaki RJ, Galatowitsch MW, Granath SR, Jacobs MS, Stec AO, Rines HW, Phillips RL (2004) Dissecting the maize genome by using chromosome addition and radiation hybrid lines. *Proc Natl Acad Sci USA* 101: 9921-9926
- La Rota M, Kantety RV, Yu JK, Sorrells ME (2005) Nonrandom distribution and frequencies of genomic and EST-derived microsatellite markers in rice, wheat and barley. *BMC Genomics* 6: 23-29
- La Rota M, Sorrells ME (2004) Comparative DNA sequence analysis of mapped wheat ESTs reveals the complexity of genome relationships between rice and wheat. *Funct Integr Genome* 4: 34-46
- Lagudah ES, Moullet O, Appels R (1997) Map based cloning of a gene sequence encoding a nucleotide binding domain and leucine rich region at the *Cre3* nematode resistance locus of wheat. *Genome* 40: 659-665
- Langer-Safer P, Levine M, Ward DC (1982) Immunocytological method for mapping genes on *Drosophila* polytene chromosomes. *Proc Natl Acad Sci USA* 79: 4381-4385
- Langridge P, Lagudah ES, Holton TA, Appels A, Sharp P, Chalmers KJ (2001) Trends in genetic and genome analyses in wheat: a review. *Aust J Agric Res* 52: 1043-1077
- Larkin DM, Pape G, Donthu R (2009) Breakpoint regions and homologous synteny blocks in chromosomes have different evolutionary histories. *Genome Res* 19: 770-777
- Lawrence JB, Singer RH, McNeil JA (1990) Interphase and metaphase resolution of different distance within the human dystrophingene. *Science* 249: 928-932
- Lewis S, Faricelli ME, Appendino ML, Valarik M, Dubcovsky J (2008) The chromosome region including the earliness per se locus *Eps-Aml* affects the duration of early developmental phases and spikelet number in diploid wheat. *J Exp Bot* 59: 3595-3607
- Li S, Jia J, Wei X, Zhang X, Li L, Chen H, Fan Y, Sun H, Zhao X, Lei T (2007) An intervarietal genetic map and QTL analysis for yield traits in wheat. *Mol Breed* 20: 167-178
- Librado P, Rozas J (2009) DnaSP v5: A software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* 25: 1451-1452
- Lichter P, Tang CJC, Call K, Hermanson G, Evans GA, Housman D, Ward DC (1990) High-resolution mapping of human chromosome 11 by *in situ* hybridization with cosmid clones. *Science* 247: 64-69
- Ling HQ, Zhu Y, Keller B (2003) High resolution mapping of the leaf rust disease resistance gene *Lr1* in wheat and characterization of BAC clone from *Lr1* locus. *Theor Appl Genet* 106: 875-882

- Ling P, Chen X, Le DQ, Campbell KG (2005) Towards cloning of the *Yr5* gene for resistance to wheat stripe rust resistance. In: Plant and Animal Genome XIII Conference, Town & Country Convention Center, San Diego, CA, pp 74
- Linkiewicz AM, Qi LL, Gill BS, Echalié B, Chao S, Lazo GR, Hummel DD, Anderson OD, Akhunov ED, Dvorak J, Pathan MS, Nguyen HT, Peng JH, Lapitan NLV, Miftahudin, Gustafson JP, La Rota CM, Sorrells ME, Hossain KG, Kalavacharla V, Kianian SF, Sandhu D, Bondaeva SN, Gill KS, Conley EJ, Anderson JA, Fenton RD, Close TJ, McGuire PE, Qualset CO, Dubcovsky J (2004) A 2500-locus bin map of wheat homoeologous group 5 provides insights on gene distribution and colinearity with rice. *Genetics* 168: 665-676
- Litt M, Luty JA (1989) A hypervariable microsatellite revealed by *in vitro* amplification of a dinucleotide repeat within the cardiac muscle actin gene. *Am J Hum Genet* 44: 397-401
- Liu S, Pumphery MO, Zhang X, Gill BS, Stack RW, Gill KS, Dolezel J, Chalhouf B, Anderson JA (2005) Towards positional cloning of *Qfhs.ndsu-3BS*, a major QTL for fusarium head blight resistance in wheat. In: Proceedings Plant & Animal Genome XIII Conference, Town & Country Convention Center, San Diego, CA, pp 307
- Liu YG, Biyashev RM, Saghai-Maroo MA (1996) Development of simple sequence repeat DNA markers and their integration into a barley linkage map. *Theor Appl Genet* 93: 869-876
- Liu YG, Tsunewaki K (1991) Restriction fragment length polymorphism analysis of wheat II Linkage maps of the RFLP sites in common wheat. *Jpn J Genet* 66: 617-633
- Lu HJ, Fellers EJP, Friesen ETL, Meinhardt SW, Faris JD (2006) Genomic analysis and marker development for the *Tsn1* locus in wheat using bin-mapped ESTs and flanking BAC contigs. *Theor Appl Genet* 112: 1132-1142
- Lukaszewski AJ, Curtis CA (1993) Physical distribution of recombination in B-genome chromosomes of tetraploid wheat. *Theor Appl Genet* 86: 121-127
- Luo M.C., Deal K.R., Akhunov E.D., Akhunova A.R., Anderson O.D., Anderson J.A., Blake N., Clegg M.T., Coleman-Derr D., Conley E.J., Crossman CC, Dubcovsky J, Gill BS, Gu YQ, Hadam J, Heo HY, Huo N, Lazo G, Ma Y, Mathews DE, McGuire PE, Morrel PL, Qualset CO, Renfro J, Tabanao D, Talbert LE, Tian C, Toleno DM, Warburton ML, You FM, Zhang W, Dvorak J (2009) Genome comparisons reveal a dominant mechanism of chromosome number reduction in grasses and accelerated genome evolution in Triticeae. *Proc Natl Acad Sci USA* 106: 15780-15785
- Ma ZQ, Roder M, Sorrells ME (1996) Frequencies and sequence characteristics of di-, tri- and tetra- nucleotide microsatellites in wheat. *Genome* 39: 123-130
- Ma ZQ, Sorrells ME, Tanksley SD (1994) RFLP markers linked to powdery mildew resistance genes *Pm1*, *Pm2*, *Pm3* and *Pm4* in wheat. *Genome* 37: 871-875

- Malik R, Brown-Guedira GL, Smith CM, Harvey TL, Gill BS (2003) Genetic mapping of wheat curl mite resistance genes *Cmc3* and *Cmc4* in common wheat. *Crop Sci* 43: 644-650
- Maluszynska J (2002) *In situ* hybridization in plants - methods and application. In: Jain SM, Brar DS, Ahloowalia BS (eds) *Molecular Techniques in Crop Improvement*, Kluwer Academic Publishers, Dordrecht, The Netherlands, pp 299-326
- Marino CL, Nelson JC, Lu YH, Sorrells ME, Leroy P, Tauleen NA, Lopes CR, Hart GE (1996) Molecular genetic maps of group 6 chromosomes of hexaploid wheat (*Triticum aestivum* L em Thell). *Genome* 39: 359-366
- Matsumoto T, Wu JZ, Kanamori H, Katayose Y, Fujisawa M, Namiki N, Mizuno H, Yamamoto K, Antonio BA, Baba T, Sakata K, Nagamura Y, Aoki H, Arikawa K, Arita K, Bito T, Chiden Y, Fujitsuka N, Fukunaka R, Hamada M, Harada C, Hayashi A, Hijishita S, Honda M, Hosokawa S, Ichikawa Y, Idonuma A, Iijima M, Ikeda M, Ikeno M, Ito K, Ito S, Ito T, Ito Y, Iwabuchi A, Kamiya K, Karasawa W, Kurita K, Katagiri S, Kikuta A, Kobayashi H, Kobayashi N, Machita K, Maehara T, Masukawa M, Mizubayashi T, Mukai Y, Nagasaki H, Nagata Y, Naito S, Nakashima M, Nakama Y, Nakamichi Y, Nakamura M, Meguro A, Negishi M, Ohta I, Ohta T, Okamoto M, Ono N, Saji S, Sakaguchi M, Sakai K, Shibata M, Shimokawa T, Song JY, Takazaki Y, Terasawa K, Tsugane M, Tsuji K, Ueda S, Waki K, Yamagata H, Yamamoto M, Yamamoto S, Yamane H, Yoshiki S, Yoshihara R, Yukawa K, Zhong HS, Yano M, Sasaki T, Yuan QP, Shu OT, Liu J, Jones KM, Gansberger K, Moffat K, Hill J, Bera J, Fadrosch D, Jin SH, Johri S, Kim M, Overton L, Reardon M, Tsitrin T, Vuong H, Weaver B, Cieccko A, Tallon L, Jackson J, Pai G, Van Aken S, Utterback T, Reidmuller S, Feldblyum T, Hsiao J, Zismann V, Iobst S, de Vazeille AR, Buell CR, Ying K, Li Y, Lu TT, Huang YC, Zhao Q, Feng Q, Zhang L, Zhu JJ, Weng QJ, Mu J, Lu YQ, Fan DL, Liu YL, Guan JP, Zhang YJ, Yu SL, Liu XH, Zhang Y, Hong GF, Han B, Choisine N, Demange N, Orjeda G, Samain S, Cattolico L, Pelletier E, Couloux A, Segurens B, Wincker P, D'Hont A, Scarpelli C, Weissenbach J, Salanoubat M, Quetier F, Yu Y, Kim HR, Rambo T, Currie J, Collura K, Luo MZ, Yang TJ, Ammiraju JSS, Engler F, Soderlund C, Wing RA, Palmer LE, de la Bastide M, Spiegel L, Nascimento L, Zutavern T, O'Shaughnessy A, Dike S, Dedhia N, Preston R, Balija V, McCombie WR, Chow TY, Chen HH, Chung MC, Chen CS, Shaw JF, Wu HP, Hsiao KJ, Chao YT, Chu MK, Cheng CH, Hour AL, Lee PF, Lin SJ, Lin YC, Liou JY, Liu SM, Hsing YI, Raghuvanshi S, Mohanty A, Bharti AK, Gaur A, Gupta V, Kumar D, Ravi V, Vij S, Kapur A, Khurana P, Khurana JP, Tyagi AK, Gaikwad K, Singh A, Dalal V, Srivastava S, Dixit A, Pal AK, Ghazi IA, Yadav M, Pandit A, Bhargava A, Sureshbabu K, Batra K, Sharma TR, Mohapatra T, Singh NK, Messing J, Nelson AB, Fuks G, Kavchok S, Keizer G, Llaca ELV, Song RT, Tanyolac B, Young S, Il KH, Hahn JH, Sangsakoo G, Vanavichit A, de Mattos LAT, Zimmer PD, Malone G, Dellagostin O, de Oliveira AC, Bevan M, Bancroft I, Minx P, Cordum H, Wilson R, Cheng ZK, Jin WW, Jiang JM, Leong SA, Iwama H, Gojobori T, Itoh T, Niimura Y, Fujii Y, Habara T, Sakai H, Sato Y, Wilson G, Kumar K, McCouch S, Juretic N, Hoen D, Wright S,

- Bruskiewich R, Bureau T, Miyao A, Hirochika H, Nishikawa T, Kadowaki K, Sugiura M (2005) The map-based sequence of the rice genome. *Nature* 436: 793-800
- Mayers KFX, Martis M, Hedley PE, Simkova H, Liu H, Mirris JA, Steuernagel B, Taudien S, Roessner S, Gundlach H, Kubalaková M, Schanková P, Murat F, Felder M, Nussbaumer T, Graner A, Salse J, Endo T, Sakai H, Tanaka T, Itoh T, Sato K, Platzer M, Matsumoto T, Scholz U, Dolezel J, Waugh R, Stein N (2011) Unlocking the barley genome by chromosomal and comparative genomics. *Plant Cell* 23: 1249-1263
- McClintok B (1931) Cytological observations of deficiencies involving known genes, translocations and an inversion in *Zea mays*. *Res Bull Mo Agric Exp Sta* 163: 1-30
- McClintok B (1933) The association of non-homologous parts of chromosomes in the mid-prophase of meiosis in *Zea mays*. *Z Zellforsch Mikrosk Anat* 19: 191-237
- McClintok B (1934) The relation of a particular chromosomal element to the development of the nucleoli in *Zea mays*. *Z Zellforsch Mikrosk Anat* 21: 293-328
- McCouch SR, Chen X, Panaud O, Temnykh S, Xu Y, Cho YG, Hunag N, Ishii T, Blair M (1997) Microsatellite marker development and mapping and applications in rice genetics and breeding. *Plant Mol Biol* 35: 89-99
- McCouch SR, Teytelman L, Xu Y, Lobos KB, Clare K, Walton M, Fu B, Maghirang R, Li Z, Xing Y, Zhang Q, Kono I, Yano M, Fjellstrom R, DeClerck G, Schneider D, Cartinhour S, Ware D, Stein L (2002) Development and mapping of new 2240 SSR markers in rice (*Oryza sativa* L.). *DNA Res* 9: 257-275
- Messmer MM, Keller M, Zanetti S, Keller B (1999) Genetic linkage map of wheat spelt cross. *Theor Appl Genet* 98: 1163-1170
- Metzgar D, Bytof J, Wills C (2000) Selection against frameshift mutations limits microsatellite expansion in coding DNA. *Genome Res* 10: 72-80
- Michalak M, Riera-Lizarazu O, Gu Y, Paux E, Feuillet C, Kalavacharla V, Hossain K, Kumar A, Kumar S, Goyal A, Tiwari V, Balyan HS, Gupta PK, Dhaliwal HS, Maan SS, Kianian SF (2008) High-resolution radiation hybrid mapping in wheat: an essential tool for the construction of the wheat physical maps. In: *Proceedings Plant & Animal Genomes XVI Conference*, Town & Country Convention Center, San Diego, CA, pp 280
- Michelmore RW, Paran I, Kessel RV (1991) Identification of markers linked to disease-resistance genes by bulked segregant analysis: a rapid method to detect the markers in specific genomic regions by using segregating populations. *Proc Natl Acad Sci USA* 88: 9828-9832
- Mickelson-Young L, Endo TR, Gill BS (1995) A cytogenetic ladder-map of the wheat homoeologous group-4 chromosomes. *Theor Appl Genet* 90: 1007-1011

- Miftahudin, Ross K, Ma XF, Mahmoud AA, Layton J, Milla MAR, Chikmawati T, Ramalingam J, Feril O, Pathan MS, Momirovic GS, Kim S, Chema K, Fang P, Haule L, Struxness H, Birkes J, Yaghoubian C, Skinner R, McAllister J, Nguyen V, Qi LL, Echalié B, Gill BS, Linkiewicz AM, Dubcovsky J, Akhunov ED, Dvorak J, Dilbirligi M, Gill KS, Peng JH, Lapitan NLV, Bermudez-Kandianis CE, Sorrells ME, Hossain KG, Kalavacharla V, Kianian SF, Lazo GR, Chao S, Anderson OD, Gonzalez-Hernandez J, Colney EJ, Anderson JA, Choi D-W, Fenton RD, Close TJ, McGuire, Qualset CO, Nguyen HT, Gustafson JP (2004) Analysis of expressed sequence tag loci on wheat chromosome group 4. *Genetics* 168: 651-663
- Milla MAR, Gustafson JP (2001) Genetic and physical characterization of chromosome 4DL in wheat. *Genome* 44: 883-892
- Mohan A, Goyal A, Singh R, Balyan HS, Gupta PK (2007) Physical mapping of wheat and rye EST-SSRs on wheat chromosomes. *Crop Sci* 47: S1-S13
- Mohler V, Schwarz G (2004) Genotyping tools in plant breeding: From restriction fragment length polymorphisms to single nucleotide polymorphisms. In: Lorz H, Wenzel (eds) *Molecular Marker Systems in Plant Breeding and Crop Improvement*, Springer Publishers, pp 23-38
- Moore G, Devos KM, Wang Z, Gale MD (1995b) Cereal genome evolution - grasses, line up and form a circle. *Curr Biol* 5: 737-739
- Moore G, Foote T, Helentjaris T, Devos KM, Kurata N, Gale MD (1995a) Was there a single ancestral cereal chromosome? *Trends Genet* 11: 81-82
- Morgante M, Hanafey M, Powell W (2002) Microsatellites are preferentially associated with non repetitive DNA in plant genomes. *Nat Genet* 30: 194-200
- Morgante M, Olivieri AM (1993) PCR-amplified SSRs as markers in plant genetics. *Plant J* 3: 175-182
- Mozo T, Dewar K, Dunn P, Ecker JR, Fischer S, Kloska S, Lehrach H, Marra M, Martienssen R, Meier-Ewert S, Altmann T (1999) A complete BAC-based physical map of the *Arabidopsis thaliana* genome. *Nat Genet* 22: 271-275
- Mun J-H, Kim D-J, Choi H-K, Gish J, Debelle F, Mudge J, Denny R, Endre G, Saurat O, Dubez A-M, Kiss GB, Roe B, Young ND, Cook DR (2006) Distribution of microsatellites in the genome of *Medicago truncatula*: A resource of genetic markers that integrate genetic and physical maps. *Genetics* 172: 2541-2555
- Munkvold JD, Greene RA, Bermudez-Kandianis CE, La Rota CM, Edwards H, Sorrells SF, Dake T, Benscher D, Kantety R, Linkiewicz AM, Dubcovsky J, Miftahudin, Gustafson JP, Pathan MS, Nguyen HT, Matthews DE, Chao S, Lazo GR, Hummel DD, Anderson OD, Anderson JA, Gonzalez-Hernandez JL, Peng JH, Lapitan N, Qi LL, Echalié B, Gill BS, Hossain KG, Kalavacharla V, Kianian SF, Sandhu D, Erayman M, Gill KS, McGuire PE,

- Qualset CO, Sorrells ME (2004) Group 3 chromosome bin maps of wheat and their relationship to rice chromosome 1. *Genetics* 168: 639-650
- Naumov GI, Turakainen H, Naumova ES, Aho S, Korhola M (1990) A new family of polymeric genes in *Saccharomyces cerevisiae* -galactosidase genes *MELI-MEL7*. *Mol Gen Genet* 224: 119-128
- Nelson JC, Allen E, Van Deynze, Autrique E, Sorrells, Lu YH, Merlino M, Atkinson M, Leroy P (1995b) Molecular mapping of wheat, homoeologous group 2. *Theor Appl Genet* 38: 516-524
- Nelson JC, Van Deynze AE, Autrique E, Sorrel ME, Lu YH, Atkinson M, Bernard M, Leroy P, Faris JD, Anderson JA (1995d) Molecular mapping of wheat: major genes and rearrangements in homoeologous groups 4, 5 and 7. *Genetics* 141: 721-731
- Nelson JC, Van Deynze AE, Autrique E, Sorrells ME, Lu YH, Atkinson M, Bernard M, Leroy P, Faris JD, Anderson JA (1995a) Molecular mapping of wheat homoeologous group 1. *Genome* 38: 508-516
- Nelson JC, Van Deynze A, Autrique E, Sorrells M, Lu YH, Negre S, Bernard M, Leroy P (1995c) Molecular mapping of wheat homoeologous group 3. *Genome* 38: 525-533
- Nelson, JC, Singh RP, Autrique JE, Sorrells ME, Lu YH, Atkinson MK, Bernard M, Leroy P, Faris JD, Anderson JA (1997) Mapping genes conferring and suppressing leaf rust resistance in wheat. *Crop Sci* 37: 1928-1935
- Nicot N, Chiquet V, Gandon B, Amilhat L, Legeai F, Leroy P, Bernard M, Sourdille P (2004) Study of simple sequence repeat (SSR) markers from wheat expressed sequence tags (ESTs). *Theor Appl Genet* 109: 800-805
- Nitta M, Nasuda S (2008) Amplification profile of the SSR markers in cultivars of hexaploid wheat and relatives. In: Appels R, Estwood R, Lagudah E, Langridge P, Mackay M, McIntyre L, Sharp P (eds) *Proceedings 11<sup>th</sup> International Wheat Genetics Symposium, Brisbane, Australia*, pp 253-255
- Ogihara Y, Hasegawa K, Tsujimoto H (1994) High-resolution cytological mapping of the long arm of chromosome 5A in common wheat using a series of deletion lines induced by gametocidal (gc) of *Aegilops speltoides*. *Mol Gen Genet* 244: 253-259
- Paillard S, Schnurbusch T, Winzeler M, Messmer M, Sourdille P, Abderhalden O, Keller B, Schachermayr G (2003) An integrative genetic linkage map of winter wheat (*Triticum aestivum* L.). *Theor Appl Genet* 107: 1235-1242
- Paran I, Zamir D (2003) Quantitative traits in plants: beyond the QTL. *Trends Genet* 19: 303-306
- Pardue ML, Gall JG (1969) Molecular hybridization of radioactive DNA to DNA of cytological preparation. *Proc Natl Acad Sci USA* 64: 600-604

- Parida SK, Raj Kumar KA, Dalal V, Singh NK, Mohapatra T (2006) Unigene derived microsatellite markers for the cereal genomes. *Theor Appl Genet* 112: 808-817
- Parker GD, Chalmers KJ, Rathjen AJ, Langridge P (1999) Mapping loci associated with milling yield in wheat (*Triticum aestivum* L). *Mol Breed* 5: 561-568
- Paterson AH, Bowers JE, Bruggmann R, Dubchak I, Grimwood J, Gundlach H, Haberler G, Hellsten U, Mitros T, Poliakov A, Schmutz J, Spannag M, Tang H, Wang X, Wicker T, Bharti AK, Chapman J, Feltus FA, Gowik U, Grigoriev IV, Lyons E, Maher CA, Martis M, Narechania A, Otiillar RP, Penning BW, Salamov AA, Wang Y, Zhang L, Carpita NC, Freeling M, Gingle1 AR, Hash CT, Keller B, Klein P, Kresovich S, McCann MC, Ming R, Peterson DG, Mehboob-ur-Rahman, Ware D, Westhoff P, Mayer KFX, Messing J, Rokhsar DS (2009) The *Sorghum bicolor* Nature 457: 551-556
- Paterson AH, Bowers JE, Chapman BA (2004) Ancient polyploidization predating divergence of cereals and its consequences for comparative genomics. *Proc Natl Acad Sci USA* 101: 9903-9908
- Paterson AH, Lin YR, Li Z, Schertz KF, Doebley JF, Pinson SRM, Liu C-S, James W, Irvine JE (1995) Convergent domestication of cereal crops by independent mutations at corresponding genetic loci. *Science* 269: 1714-1718
- Paux E, Roger D, Badaeva E, Gay G, Bernard M, Sourdille P, Feuillet C (2006) Characterizing the composition and evolution of homoeologous genomes in hexaploid wheat through BAC-end sequencing on chromosome 3B. *Plant J* 48: 463-474
- Paux E, Sourdille P, Salse J, Saintenac C, Choulet F, Leroy P, Korol A, Michalak M, Kianian S, Spielmeyer W, Lagudah E, Somers D, Kilian A, Alaux M, Vautrin S, Bergès H, Eversole K, Appels R, Safar J, Simkova H, Dolezel J, Bernard M, Feuillet C (2008) A physical map of the 1-gigabase bread wheat chromosome 3B. *Science* 322: 101-104
- Peleg Z, Saranga Y, Suprunova T, Ronin Y, Roder MS, Kilian A, Korol AB, Fahima T (2008) High-density genetic map of durum wheat × wild emmer wheat based on SSR and DArT markers. *Theor Appl Genet* 117: 103-115
- Peng JH, Lapitan NLV (2005) Characterization of EST-derived microsatellites in the wheat genome and development of eSSR markers. *Funct Integr Genomics* 5: 80-96
- Peng JH, Zadeh H, Lazo GR, Gustafson JP, Chao S, Anderson OD, Qi LL, Echaliier B, Gill BS, Dilbirligi M, Sandhu D, Gill KS, Greene RA, Sorrells ME, Akhunov ED, Dvorák J, Linkiewicz AM, Dubcovsky J, Hossain KG, Kalavacharla V, Kianian SF, Mahmoud AA, Miftahudin, Conley EJ, Anderson JA, Pathan MS, Nguyen HT, McGuire PE, Qualset CO, Lapitan NLV (2004) Chromosome bin map of expressed sequence tags in homoeologous group 1 of hexaploid wheat and homoeology with rice and *Arabidopsis*. *Genetics* 168: 609-623

- Penner GA, Zirino M, Kruger S, Townley-Smith F (1998) Accelerated recurrent parent selection in wheat with microsatellite markers. In: Slinkard AE (eds) Proceeding 9<sup>th</sup> International Wheat Genetics Symposium, University Extension Press, University of Saskatchewan, Saskatoon, Canada, pp 131-134
- Pestsova E, Ganal MW, Roder MS (2000) Isolation and mapping of microsatellite markers specific for D genome of bread wheat (*Triticum aestivum* L). *Genome* 43: 689-697
- Peters JL, Cnudde F, Gerats T (2003) Forward genetics and map-based cloning approaches. *Trends Plant Sci* 8: 484-491
- Pillen K, Binder A, Kreuzkam B, Ramsay L, Waugh R, Forster J, Leon J (2000) Mapping new EMBL-derived barley microsatellites and their use in differentiating German barley cultivars. *Theor Appl Genet* 101: 652-660
- Plaschke J, Ganal M, Röder MS (1995) Detection of genetic diversity in closely related bread wheat using microsatellite markers. *Theor Appl Genet* 91: 1001-1007
- Powell W, Morgante M, Andre C, Hanafey M, Vogel J, Tingey S, Rafalski JA (1996) The comparison of RFLPs, RAPDs, AFLPs and SSR (microsatellite) markers for germplasm analysis. *Mol Breed* 2: 225-238
- Prasad M, Kumar N, Kulwal PL, Röder MS, Balyan HS, Dhaliwal HS, Gupta PK (2003) QTL analysis for grain protein content using SSR markers and validation studies using NILs in bread wheat. *Theor Appl Genet* 106: 659-667
- Prasad M, Varshney RK, Kumar A, Balyan HS, Sharma PC, Edwards KJ, Singh H, Dhaliwal HS, Roy JK, Gupta PK (1999) A microsatellite marker associated with a QTL for grain protein content on chromosome arm 2DL of bread wheat. *Theor Appl Genet* 99: 341-345
- Prasad M, Varshney RK, Roy JK, Balyan HS, Gupta PK (2000) The use of microsatellites for detecting DNA polymorphism, genotype identification and genetic diversity in wheat. *Theor Appl Genet* 100: 584-592
- Qi LL, Gill BS (2001) High-density physical maps reveal that the dominant male-sterile gene Ms3 is located in a genomic region of low recombination in wheat and is not amenable to map-based cloning. *Theor Appl Genet* 103: 998-1006
- Qi LL, Echalié B, Friebe B, Gill BS (2003) Molecular characterization of a set of wheat deletion stocks for use in chromosome bin mapping of ESTs. *Funct Integr Genomics* 3: 39-55
- Qi LL, Friebe B, Wu J, Gu Y, Qian C, Gill BS (2010) The compact *Brachypodium* genome conserve centromeric regions of a common ancestor with wheat and rice. *Funct Integr Genomics* 10: 477-492
- Qi, LL, Echalié B, Chao S, Lazo GR, Butler GE, Anderson OD, Akhunov ED, Dvorak J, Linkiewicz AM, Ratnasiri A, Dubcovsky J, Bermudez-Kandianis CE, Greene RA, Kantety

- R, La Rota CM, Munkvold JD, Sorrells SF, Sorrells SF, Dilbirligi M, Sidhu D, Erayman M, Randhawa HS, Sandhu D, Bondareva SN, Gill KS, Mahmoud AA, MA X-F Miftahudin, Gustafson JP, Conley AJ, Nduati V, Gonzalez-Hernandez JL, Anderson JA, . Peng JH, Lapitan NLV, Hossain KG, Kalavacharla V, Kianian SF, Pathan MS, Zhang DS, Nguyen HT, Choi DW, Fenton RD, Close TJ, McGuire PE, Qualset CO, Gill BS (2004) A chromosome bin map of 16,000 expressed sequence tag loci and distribution of genes among the three genomes of polyploidy wheat. *Genetics* 168: 701-712
- Quarrie SA, Steed A, Calestani C, Semikhodskii A, Lebreton C, Chinoy C, Steele N, Pljevljakusic D, Waterman E, Weyen J, Schondelmaier J, Habash DZ, Farmer P, Saker L, Clarkson DT, Abugalieva A, Yessimbekova M, Turuspekov Y, Abugalieva S, Tuberosa R, Sanguineti M-C, Hollington PA, Aragues R, Royo A, Dodig D (2005) A high-density genetic map of hexaploid wheat (*Triticum aestivum* L.) from the cross Chinese Spring SQ1 and its use to compare QTLs for grain yield across a range of environments. *Theor Appl Genet* 110: 865-880
- Rafalski JA (2002a) Novel genetic mapping tools in plants: SNPs and LD-based approaches. *Plant Sci* 162: 329-333
- Rafalski JA (2002b) Application of single nucleotide polymorphism in crop genetics. *Curr Opin Plant Biol* 5: 94-100
- Rakoczy-Trojanowska M, Bolibok H (2004) Characteristics and a comparison of three classes of microsatellite-based markers and their application in plants. *Cell Mol Biol Lett* 9: 221-238
- Ramsay L, Macaulay M, Degli Ivanissevich S, MacLean K, Cardle L, Fuller J, Edwards KJ, Tuvevsson S, Morgante M, Massari A, Maestri E, Marmiroli N, Sjakste T, Ganal M, Powell W, Waugh R (2000) A simple sequence repeat-based linkage map of barley. *Genetics* 156: 1997-2005
- Randhawa HS, Dilbirligi M, Sidhu D, Erayman M, Sandhu D, Bondareva S, Chao S, Lazo GR, Anderson OD, Miftahudin, Gustafson JP, Echaliier B, Qi LL, Gill BS, Akhunov ED, Dvorak J, Linkiewicz AM, Ratnasiri A, Dubcovsky J, Bermudez-Kandianis CE, Greene RA, Sorrells SF, Conley EJ, Anderson JA, Peng JH, Lapitan NLV, Hossain KG, Kalavacharla V, Kianian SF, Pathan MS, Nguyen HT, Endo TR, Close TJ, McGuire PE, Qualset CO and Gill BS (2004) Deletion mapping of homoeologous group 6- *Genetics* 168: 677-686
- Rayburn AL, Gill BS (1985) Use of biotin-labeled probes to map specific DNA sequences of wheat chromosomes. *J Hered* 76: 78-81
- Reif JC, Zhang P, Dreisigacker S, Warburton ML, Van Ginkel M, Hoisington D, Bohn M, Melchinger E (2005) Wheat genetic diversity trends during domestication and breeding. *Theor Appl Genet* 110: 859-864

- Reyes BGD, Khush GS, Brar DS (1998) Chromosomal location of eight isozyme loci in rice using primary trisomics and monosomic alien addition lines. *J Hered* 89: 164-168
- Rick CM (1971) Some cytogenetic features of the genome in diploid plant species. *Stadler Symp*, Vol 1 and 2: 153-174
- Rick CM, Khush GS (1966) Chromosomes engineering in *Lycopersicon*. *Heredity* 20: 8-20
- Riera-Lizarazu O, Rines HW, Phillips RL (1996) Cytological and molecular characterization of oat and maize partial hybrids. *Theor Appl Genet* 93: 123-135
- Riera-Lizarazu O, Vales MI, Ananiev EV, Rines HW, Philips RL (2000) Production and characterization of maize chromosome 9 radiation hybrids derived from an oat-maize addition line. *Genetics* 156: 327-339
- Roberts MA, Reader SM, Dalgliesh C, Miller TE, Foote TN, Fish LJ, Snape JW, Moore G (1999) Induction and characterization of *Ph1* wheat mutants. *Genetics* 153: 1909-1918
- Roberts PA (1965) Difference in the behavior of eu- and hetero-chromatin: crossing-over. *Nature* 205: 725-726
- Roder MS, Huang X-Q, Ganai MW (2004) Wheat microsatellites: potential and implications. In: Lorz H, Wenzel G (eds) *Molecular Marker Systems in Plant Breeding and Crop Improvement. Biotechnology in Agriculture and Forestry*, Vol 55, Springer, Berlin Heidelberg, pp 255-266
- Roder MS, Korzun V, Gill BS, Ganai MW (1998b) The physical mapping of microsatellite markers in wheat. *Genome* 41: 278-283
- Roder MS, Korzun V, Wendehake K, Plaschke J, Tixier MH, Leroy P, Ganai MW (1998a) A microsatellite map of wheat. *Genetics* 149: 2007-2023
- Roder MS, Plaschke J, Konig SU, Borner A, Sorrels ME (1995) Abundance, variability and chromosomal location of microsatellite in wheat. *Mol Gen Genet* 246: 327-333
- Rossetto M, McNally J, Henry RJ (2002) Evaluating the potential of SSR flanking regions for examining taxonomic relationships in the Vitaceae. *Theor Appl Genet* 104: 61-6
- Rostoks N, Mudie S, Cardle L, Russell J, Ramsay L, Booth A, Svensson JT, Wanamaker SI, Walia H, Rodriguez EM, Hedley PE, Liu H, Morris J, Close TJ, Marshall DF, Waugh R (2005) Genome-wide SNP discovery and linkage analysis in barley based on genes responsive to abiotic stress. *Mol Genet Genomics* 274: 515-527
- Roussel V, Koenig J, Beckert M, Balfourier F (2004) Molecular diversity in French bread wheat accessions related to temporal trends and breeding programmes. *Theor Appl Genet* 108: 920-930
- Roy JK, Prasad M, Varshney RK, Balyan HS, Blake TK, Dhaliwal HS, Singh H, Edwards KJ, Gupta PK (1999) Identification of a microsatellite on chromosome 6B and a STS on 7D of

- bread wheat showing an association with pre-harvest sprouting tolerance. *Theor Appl Genet* 99: 336-340
- Rustgi S, Bandopadhyay R, Balyan HS, Gupta PK (2009) EST-SNPs in bread wheat: discovery, validation, genotyping and haplotype structure. *Czech J Genet Breed* 45: 106-116
- Safar J, Bartos J, Janda J, Bellec A, Kubalaková M, Valarik M, Pateyron S, Weiserová J, Tusková R, Cihalíková J, Vrana J, Simková H, Faivre-Rampant P, Sourdille P, Caboche M, Bernard M, Dolezel J and Chalhoub B (2004) Dissecting large and complex genomes: flow sorting and BAC cloning of individual chromosome from bread wheat. *Plant J* 39: 960-968
- Saghai-Marooif MA, Biyashev RM, Yang GP, Zhang Q, Allard RW (1994) Extraordinary polymorphic microsatellite DNA in barley: species diversity, chromosomal locations and population dynamics. *Proc Natl Acad Sci USA* 91: 5466-5470
- Saghai-Marooif MA, Soliman KM, Jorgensen RA, Allard RW (1984) Ribosomal DNA spacer length polymorphism in barley: Mendelian inheritance, chromosomal location and population dynamics. *Proc Natl Acad Sci USA* 81: 8014-8018
- Saha MC, Rouf Mian MA, Eujayl I, Zwonitzer JC, Wang L, May GD (2004) Tall fescue EST-SSR markers with transferability across several grass species. *Theor Appl Genet* 109: 783-791
- Saintenac C, Falque M, Martin OC, Paux E, Feuillet C, Sourdille P (2009) Detailed recombination studies along chromosome 3B provide new insights on crossover distribution in wheat (*Triticum aestivum* L.). *Genetics* 181: 393-403
- Sakamura T (1918) Kurze Mitteilung über die Chromosomenzahlen und die Verwandtschaftsverhältnisse der *Triticum*-Arten. *Bot Mag* 32: 151-154
- Salse J, Bolot S, Throude M, Jouffe V, Piegue B, Quraishi UM, Calcagno T, Cooke R, Delseny M, Feuillet C (2008) Identification and characterization of shared duplications between rice and wheat provide new insight into grass genome evolution. *Plant Cell* 20: 11-24
- Salvi S, Tuberosa R (2005) To clone or not to clone plant QTLs: present and future challenges. *Trends Plant Sci* 10: 297-304
- Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular Cloning: A Laboratory Manual*, 2<sup>nd</sup> edition, Vol. 1. Cold Spring Harbor Laboratory Press, Plainview, New York, pp 1.18
- Sandhu D, Champoux AJA, Bondareva SN, Gill KS (2001) Identification and physical localization of useful genes and markers to a major gene-rich region on wheat group 1S chromosomes. *Genetics* 157: 1735-1747
- Sandhu D, Gill KS (2002a) Gene-containing regions of wheat and the other grass genomes. *Plant Physiol* 128: 803-811
- Sandhu D, Gill KS (2002b) Structural and functional organization of the '1S08 gene-rich region in the Triticeae. *Plant Mol Biol* 48: 791-804

- SanMiguel P, Tikhonov A, Jin YK, Motchoulskaia N, Zakharov D, Melake-Berhan A, Springer PS, Edwards KJ, Lee M, Avramova Z, Bennetzen JL (1996) Nested retrotransposons in the intergenic regions of the maize genome. *Science* 274: 765-768
- Sasaki T, Antonio BA (2004) Rice genome as a model system for cereals. In: Gupta PK, Varshney RK (eds) *Cereal Genomics*, Kluwer Academic Publishers, Dordrecht, The Netherlands, pp 535-557
- Sasaki T, Burr B (2000) International rice genome sequencing project: the effort to completely sequence the rice genome. *Curr Opin Plant Biol* 3: 138-141
- Schnable PS, Ware D, Fulton RS, Schnable PS, Ware D, Fulton RS, Stein JC, Wei F, Pasternak S, Liang C, Zhang J, Fulton L, Graves TA, Minx P, Reily AD, Courtney L, Kruchowski SS, Tomlinson C, Strong C, Delehaunty K, Fronick C, Courtney B, Rock SM, Belter E, Du F, Kim K, Abbott RM, Cotton M, Levy A, Marchetto P, Ochoa K, Jackson SM, Gillam B, Chen W, Yan L, Higginbotham J, Cardenas M, Waligorski J, Applebaum E, Phelps L, Falcone J, Kanchi K, Thane T, Scimone A, Thane N, Henke J, Wang T, Ruppert J, Shah N, Rotter K, Hodges J, Ingenthron E, Cordes M, Kohlberg S, Sgro J, Delgado B, Mead K, Chinwalla A, Leonard S, Crouse K, Collura K, Kudrna D, Currie J, He R, Angelova A, Rajasekar S, Mueller T, Lomeli R, Scara G, Ko A, Delaney K, Wissotski M, Lopez G, Campos D, Braidotti M, Ashley E, Golser W, Kim HR, Lee S, Lin J, Dujmic Z, Kim W, Talag J, Zuccolo A, Fan C, Sebastian A, Kramer M, Spiegel L, Nascimento L, Zutavern T, Miller B, Ambroise C, Muller S, Spooner W, Narechania A, Ren L, Wei S, Kumari S, Faga B, Levy MJ, McMahan L, Buren PV, Vaughn MW, Ying K, Yeh C-T, Emrich SJ, Jia Y, Kalyanaraman A, Hsia A-P, Barbazuk WB, Baucom RS, Brutnell TP, Carpita NC, Chaparro C, Chia J-M, Deragon L-M, Estill JC, Fu Y, Jeddelloh JA, Han J, Lee H, Li P, Lisch DR, Liu S, Liu Z, Nagel DH, McCann MC, Miguel PS, Myers AM, Nettleton D, Nguyen J, Penning BW, Ponnala L, Schneider KL, Schwartz DC, Sharma A, Soderlund C, Springer NM, Sun Q, Wang H, Waterman M, Westerman R, Wolfgruber TK, Yang L, Yu U, Zhang L, Zhou S, Zhu Q, Bennetzen, Dawe RK, Jiang J, Jiang N, Presting GG, Wessler SR, Aluru S, Martienssen RA, Clifton SW, McCombie WR, Wing RA, Wilson RK (2009) The B73 maize genome: complexity, diversity and dynamics. *Science* 326: 1112-1115
- Schnurbusch T, Sutton T, Paull J, Jamjod S, Rathjen A, Langridge P, Collins N (2005) Towards isolation of boron tolerance genes in wheat and barley. In: Plant and Animal Genome XIII Conference, Town & Country Convention Centre, San Diego CA, pp 5
- Scofield SR, Huang L, Brandt AS, Gill BS (2005) Development of a virus-induced gene silencing system for hexaploid wheat and its use in functional analysis of the *Lr21*-mediated leaf rust resistance pathway. *Plant Physiol* 138: 2165-2173
- Scott KD, Egger P, Seaton G, Rossetto M, Ablett EM, Lee LS, Henry RJ (2000) Analysis of SSRs derived from grape ESTs. *Theor Appl Genet* 100: 723-726

- Scotti I, Mangi F, Fink R, Powell W, Binelli G, Hedley PE (2000) Microsatellites are not randomly distributed within Norway spruce (*Picea abies* K.) expressed sequences. *Genome* 43: 41-46
- Sears ER (1954) The aneuploides of common wheat. *Missouri Agric Exp Sta Res Bull* 572: 1-59
- Sears ER (1966) Nullisomic-tetrasomic combinations in hexaploid wheat. In: Riley R, Lewis KR (eds) *Chromosome Manipulation and Plant Genetics*, Oliver and Boyd, Edinburgh, pp 29-45
- Sears ER, Sears LMS (1978) The telocentric chromosome of common wheat. In: Ramanujan S (eds) *Proceedings 5<sup>th</sup> International Wheat Genetics Symposium*, New Delhi, pp 3-6
- See DR, Brooks S, Nelson JC, Brown-Guedira G, Friebe B, Gill BS (2006) Gene evolution at the ends of wheat chromosomes. *Proc Natl Acad Sci USA*, 103: 4162-4167
- Semagn S, Bjornstad A, Skinnes H, Mary AG, Tarkegne Y, William M (2006) Distribution of DArT, AFLP, and SSR markers in a genetic linkage map of a doubled-haploid hexaploid wheat population. *Genome* 49: 545-555
- Sharopova N, McMullen MD, Schultz L, Schroeder S, Sanchez-Villeda H, Gardiner J, Bergstrom D, Houchins K, Melia-Hancock S, Musket T, Duru N, Polacco M, Edwards K, Ruff T, Register JC, Brouwer C, Thompson R, Velasco R, Chin E, Lee M, Woodman-Clíkeman W, Long MJ, Liscum E, Cone K, Davis G, Coe EH (2002) Development and mapping of SSR markers for maize. *Plant Mol Biol* 48: 463-481
- Shoemaker R, Grant D, Olson T, Warren WC, Wing R, Yu Y, Kim H, Cregan P, Joseph B, Futrell-Griggs M, Nelson W, Davito J, Walker J, Wallis J, Kremitski C, Scheer D, Clifton S, Graves T, Nguyen HT, Wu XL, Luo MC, Dvorak J, Nelson R, Cannon S, Thomkins J, Schmutz J, Stacey G, Jackson S (2008) Microsatellite discovery from BAC end sequences and genetic mapping to anchor the soybean physical and genetic maps. *Genome* 51: 294-302
- Shultz JL, Kazi S, Bashir R, Afzal JA, Lightfoot DA (2007) The development of BAC-end sequence-based microsatellite markers and placement in the physical and genetic maps of soybean. *Theor Appl Genet* 114: 1081-1090
- Sidhu D, Gill KS (2004) Distribution of genes and recombination in wheat and other eukaryotes. *Plant Cell Tissue Organ Cult* 79: 257-270
- Siedler H, Messmer MM, Schachermayr GM, Winzeler H, Winzeler M, Keller B (1994) Genetic diversity in European wheat and spelt breeding material based on RFLP data. *Theor Appl Genet* 88: 994-1003
- Sim SC, Yu JK, Jo Y, Sorrells ME, Jung G (2009) Transferability of cereal EST-SSR markers to ryegrass. *Genome* 52: 431-437

- Simmonds JR, Fish LJ, Leverington-Waite MA, Wang Y, Howell P, Snape JW (2008) Mapping of a gene (*Vir*) for a non-glaucous, viridescent phenotype in bread wheat derived from *Triticum dicoccoides*, and its association with yield variation. *Euphytica* 159: 333-341
- Simons KJ, Fellers JP, Trick HN, Zhang Z, Tai Y-S, Gill BS, Faris JD (2005) Molecular characterization of the major wheat domestication gene *Q*. *Genetics* 172: 547-555
- Singh H, Prasad M, Varshney RK, Roy JK, Balyan HS, Dhaliwal HS, Gupta PK (2001) STMS markers for grain protein content and their validation using near-isogenic lines in bread wheat. *Plant Breed* 120: 273-278
- Singh K, Multani DS, Khush GS (1996) Secondary trisomics and telotrisomics of rice: origin, characterization and use in determining the orientation of chromosome map. *Genetics* 143: 517-519
- Singh NK, Dalal V, Batra K, Singh BK, Chitra G, Singh A, Ghazi IA, Yadav M, Pandit A, Dixit R, Singh PK, Singh H, Koundal KR, Gaikwad K, Mohapatra T, Sharma TR (2007) Single-copy genes define a conserved order between rice and wheat for understanding differences caused by duplication, deletion, and transposition of genes. *Funct Integr Genomics* 7: 17-35
- Singh NK, Raghuvanshi S, Srivastava SK, Gaur A, Pal AK, Dalal V, Singh A, Ghazi IA, Bhargav A, Yadav M, Dixit A, Batra K, Gaikwad K, Sharma TR, Mohanty A, Bharti AK, Kapur A, Gupta V, Kumar D, Vij S, Vydiyanathan R, Khurana P, Sharma S, McCombie WR, Messing J, Wing R, Sasaki T, Khurana P, Mohapatra T, Khurana JP, Tyag AK (2004) Sequence analysis of the long arm of rice chromosome 11 for rice-wheat synteny. *Funct Integr Genomics* 4: 102-117
- Singh R, Kumar N, Bandopadhyay R, Rustgi S, Sharma S, Balyan HS, Gupta PK (2006) Development and use of anchored-SSRs to study DNA polymorphism in bread wheat (*Triticum aestivum* L.). *Mol Ecol Notes* 6: 296-299
- Singh S, Franks CD, Huang L, Brown-Guedira GL, Marshall DS, Gill BS, Fritz A (2004) *Lr41*, *Lr39*, and a leaf rust resistance gene from *Aegilops cylindrica* may be allelic and are located on wheat chromosome 2DS. *Theor Appl Genet* 108: 586-591
- Singleton WR (1939) Cytological observations on deficiencies produced by treating maize pollen with ultraviolet light. *Genetics* 24: 109-115
- Somers DJ, Isaac P, Edwards KJ (2004) A high density consensus map for bread wheat (*Triticum aestivum* L.). *Theor Appl Genet* 109: 1105-1114
- Somers DJ, Kirkpatrick R, Moniwa M, Walsh A (2003) Mining single-nucleotide polymorphisms from hexaploid wheat ESTs. *Genome* 46: 431-437
- Song QJ, Fickus EW, Cregan PB (2002) Characterization of trinucleotide SSR motifs in wheat. *Theor Appl Genet* 104: 286-293

- Song QJ, Shi JR, Singh S, Fickus EW, Costa JM, Lewis J, Gill BS, Ward R, Cregan PB (2005) Development and mapping of microsatellite (SSR) markers in wheat. *Theor Appl Genet* 110: 550-560
- Sorrells ME, La Rota M, Bermudez-Kandianis CE, Greene RA, Kantety R, Munkvold JD, Miftahudin , Mahmoud A, Ma X, Gustafson PJ, Qi LL, Echalié B, Gill BS, Matthews DE, Lazo GR, Chao S, Anderson OD, Edwards H, Linkiewicz AM, Dubcovsky J, Akhunov ED, Dvorak J, Zhang D, Nguyen HT, Peng J, Lapitan NLV, Gonzalez-Hernandez JL, Anderson JA, Hossain K, Kalavacharla V, Kianian SF, Choi D-W, Close TJ, Dilbirligi M, Gill KS, Steber C, Walker-Simmons MK, McGuire PE, Qualset CO (2003) Comparative DNA sequence analysis of wheat and rice genomes. *Genome Res* 13: 1818-1827
- Sourdille P, Cadalen T, Guyomarch H, Snape JW, Perretant MR, Charmet G, Boeuf C, Bernard S, Bernard M (2003) An update of the Courtot-Chinese Spring intervarietal molecular marker linkage map for the QTL detection of agronomic traits in wheat. *Theor Appl Genet* 106: 530-538
- Sourdille P, Guyomarch H, Baron C, Gandon B, Chiquet V (2001) Improvement of the genetic maps of wheat using new microsatellite markers. In: *Plant and Animal Genome IX Conference*, Town & Country Convention Centre, San Diego CA, pp 42
- Sourdille P, Singh S, Cadalen T, Gina L, Brown-Guedira, Gay G, Qi L, Gill BS, Dufour P, Murigneux A, Bernard M (2004) Microsatellite-based deletion bin system for the establishment of genetic-physical map relationships in wheat (*Triticum aestivum* L). *Funct Integr Genomics* 4: 12-25
- Srinivasachary, Dida MM, Gale MD, Devos KM (2007) Comparative analyses reveal high levels of conserved colinearity between the finger millet and rice genomes. *Theor Appl Genet* 115: 489-499
- Stadler LJ (1933) On the genetic nature of induced mutations in plants. *Res Bull Mo Agric Exp Sta* 204: 3-29
- Stadler LJ (1935) Genetic behavior of a haplo-viable internal deficiency in maize. *Amer Nat* 69: 80-81
- Stein N, Graner A (2004) Map-based isolation in cereal genomes. In: Gupta PK, Varshney RK (eds) *Cereal Genomics*, Kluwer Academic Publishers, Dordrecht, The Netherlands, pp 331-360
- Stephenson P, Bryan G, Kirby J, Collins A, Devos KM, Busso C, Gale MD (1998) Fifty new microsatellite loci for wheat genetic map. *Theor Appl Genet* 97: 946-949
- Svitashev SK, Somers DA (2002) Characterization of transgene loci in plants using FISH: A picture is worth a thousand words. *Plant Cell Tissue Organ Cult* 69: 205-214

- Tak LA, Song J, Macas J, Jiang J (2005) Sobo, a recently amplified satellite repeat of potato, and its implications for the origin of tandemly repeated sequences. *Genetics* 170: 1231-1238
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S (2011) MEGA5: Molecular Evolutionary Genetics Analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Bio Evol*, doi:10.1093/molbev/msr121
- Tao Q, Chang YL, Wang J, Chen H, Islam-Faridi MN, Scheuring C, Wang B, Stelly DM, Zhang HB (2001) BAC-based physical map of the rice genome constructed by restriction fingerprint analysis. *Genetics* 158: 1711-1724
- Tegelstrom H (1992) Detection of mitochondrial DNA fragments. In: Hoelzel AR (eds) *Molecular Genetic Analysis of Populations: A Practical Approach*, IRL Press, Oxford, pp 89-114
- The *Arabidopsis* Genome Initiative (2000) Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* 408: 796-815
- The International *Brachypodium* Initiative (2010) Genome sequencing and analysis of the model grass *Brachypodium distachyon*. *Nature* 463: 763-768
- Thiel T, Graner A, Waugh R, Grosse I, Close TJ, Stein N (2009) Evidence and evolutionary analysis of ancient whole-genome duplication in barley predating the divergence from rice. *BMC Evol Biol* 9: 209-226
- Thiel T, Michalek W, Varshney RK, Graner A (2003) Exploiting EST database for the development and characterization of gene derived SSR-markers in barley (*Hordeum vulgare* L). *Theor Appl Genet* 106: 411-422
- Torada A, Koike M, Mochida K, Ogihara Y (2006) SSR-based linkage map with new markers using an intraspecific population of common wheat. *Theor Appl Genet* 112: 1042-1051
- Tsuchiya T (1991) Chromosome mapping by means of aneuploid analysis in barley. In: Gupta PK, Tsuchiya T (eds) *Chromosome Engineering in Plants: Genetics, Breeding, Evolution*, Part A, Elsevier Publication, Amsterdam, pp 361-384
- Uauy C, Distelfeld A, Fahima T, Blechl A, Dubcovsky J (2006) A NAC gene regulating senescence improves grain protein, zinc, and iron content in wheat. *Science* 314: 1298-1301
- Van Deynze AE, Dubcovsky J, Gill KS, Nelson JC, Sorrells ME, Dvorak J, Gill BS, Lagudah ES, McCouch SR, Appels R (1995) Molecular-genetic maps for group 1 chromosomes of Triticeae species and their relation to chromosomes in rice and oat. *Genome* 38: 45-59
- anchored primers for the  
enhanced recovery of specific microsatellite markers in *Brassica napus* L. *Theor Appl Genet* 101: 115-119
- Varshney R, Graner A, Sorrells ME (2005) Genic microsatellite markers in plants: features and applications. *Trends Biotechnol* 23: 48-55

- Varshney RK, Prasad M, Roy JK, Kumar N, Singh H, Dhaliwal HS, Balyan HS, Gupta PK (2000) Identification of eight chromosomes and a microsatellite marker on 1A associated with QTL for grain weight in bread wheat. *Theor Appl Genet* 100: 1290-1294
- Vikal Y, Chhuneja P, Singh R, Dhaliwal HS (2004) Tagging of an *Aegilops speltoides* derived leaf rust resistance gene *Lr28* with a microsatellite marker in wheat. *J Plant Biochem Biotech* 13: 47-49
- Vogel JP, Garvin DF, Leong OM, Hayden DM (2006b) Agrobacterium-mediated transformation and inbred line development in the model grass *Brachypodium distachyon*. *Plant Cell Tissue Organ Cult* 85: 199-211
- Vogel JP, Gu YQ, Twigg P, Lazo GR, Laudencia-Chingcuanco D, Hayden DM, Donze TJ, Vivian LA, Stamova B, Coleman-Derr D (2006a) EST sequencing and phylogenetic analysis of the model grass *Brachypodium distachyon*. *Theor Appl Genet* 113: 186-195
- Vos P, Hogers R, Bleeker M, Reijans M, van de Lee T, Hornes M, Frijters A, Pot J, Peleman J, Kuiper M, Zabeau M (1995) AFLP: a new technique for DNA fingerprinting. *Nucleic Acids Res* 23: 4407-4414
- Wang D, Shi J, Carlson SR, Cregan PB, Ward RW, Diers BW (2003) A low-cost, high-throughput polyacrylamide gel electrophoresis system for genotyping with microsatellite DNA markers. *Crop Sci* 43: 1828-1832
- Wang Y, Tang X, Cheng Z, Mueller L, Giovannoni J, Tanksley SD (2006) Euchromatin and pericentromeric heterochromatin: comparative composition in the tomato genome. *Genetics* 172: 2529-2540
- Wardrop J, Fuller J, Powell W, Machray GC (2004) Exploiting plant somatic radiation hybrids for physical mapping of expressed sequence tags. *Theor Appl Genet* 108: 343-348
- Wardrop J, Snape JW, Powell W, Machray GC (2002) Constructing plant radiation hybrid panels. *Plant J* 31: 223-228
- Wei F, Coe E, Nelson W, Bharti AK, Engler F, Butler E, Kim H, Goicoechea JL, Chen M, Lee S, Fuks G, Sanchez-Villeda H, Schroeder S, Fang Z, McMullen M, Davis G, Bowers JE, Paterson AH, Schaeffer M, Gardiner J, Cone K, Messing J, Soderlund C, Wing RA (2007) Physical and genetic structure of the maize genome reflects its complex evolutionary history. *PLoS Genet* 3: e123
- Weng Y, Lazar MD (2002) Comparison of homoeologous group-6 short arm physical maps of wheat and barley reveals a similar distribution of recombinogenic and gene-rich regions. *Theor Appl Genet* 104: 1078-1085
- Weng Y, Tuleen NA, Hart GE (2000) Extended physical maps and a consensus physical map of the homoeologous group-6 chromosomes of wheat (*Triticum aestivum* L. em Thell). *Theor Appl Genet* 100: 519-527

- Werner JE, Endo TR, Gill BS (1992) Toward a cytogenetically based physical map of the wheat genome. *Proc Natl Acad Sci USA* 89: 11307-11311
- Wicker T, Buchmann JP, Keller B (2010) Patching gaps in plant genomes results in gene movement and erosion of colinearity. *Genome Res* 20: 1229-1237
- Wicker T, Yahiaoui N, Guyot R, Schlagenhauf E, Liu Z-D, Dubcovsky J, Keller B (2003) Rapid genome divergence at orthologous low molecular weight glutenin loci of the A and A<sup>(m)</sup> genomes of wheat. *Plant Cell* 15: 1186-1197
- Wicker T, Yahiaoui N, Keller B (2007) Contrasting rates of evolution in *Pm3* loci from three wheat species and rice. *Genetics* 177: 1207-1216
- Williams JGK, Kubelik ARK, Livak JL, Rafalski JA, Tingey SV (1990) DNA polymorphisms amplified by random primers are useful as genetic markers. *Nucl Acids Res* 18: 6531-6535
- Williams KJ, Fisher JM, Langridge P (1994) Identification of RFLP markers linked to the cereal cyst nematode resistance gene (*Cre*) in wheat. *Theor Appl Genet* 83: 919-924
- Wu C, Sun S, Nimmakayala P, Felipe A, Meksem K, Springman R, Ding K, Lightfoot DA, Zhang H-B (2004) A BAC- and BIBAC-based physical map of the soybean genome. *Genome Res* 14: 319-326
- Wu KS, Tanksley SD (1993) Abundance, polymorphism and genetic mapping of microsatellites in rice. *Mol Gen Genet* 241: 225-235
- Xie DX, Devos KM, Moore G, Gale MD (1993) RFLP-based genetic maps of the homoeologous group 5 chromosomes of bread wheat (*Triticum aestivum* L). *Theor Appl Genet* 87: 70-74
- Xue S, Zhang Z, Lin F, Kong Z, Cao Y, Li C, Yi H, Mei M, Zhu H, Wu J, Xu H, Zhao D, Tian D, Zhang C, Ma Z (2008) A high-density intervarietal map of the wheat genome enriched with markers derived from expressed sequence tags. *Theor Appl Genet* 117: 181-189
- Yahiaoui N, Srichumpa P, Dudler R, Keller B (2004) Genome analysis at different ploidy levels allows cloning of the powdery mildew resistance gene *Pm3b* from hexaploid wheat. *Plant J* 37: 528-538
- Yan L, Fu D, Li C, Blechl A, Tranquilli G, Bonafede M, Sanchez A, Valarik M, Yasuda S, Dubcovsky J (2006) The wheat and barley vernalization gene *VRN3* is an orthologue of *FT*. *Proc Natl Acad Sci USA* 103: 19581-19586
- Yan L, Loukoianov A, Blechl A, Tranquilli G, Ramakrishna W, SanMiguel P, Bennetzen JL, Echenique V, Dubcovsky J (2004) The wheat *VRN2* gene is a flowering repressor down-regulated by vernalization. *Science* 303: 1640-1644
- Yan L, Loukoianov A, Tranquilli G, Helguera M, Fahima T, Dubcovsky J (2003) Positional cloning of the wheat vernalization gene *VRN1*. *Proc Natl Acad Sci USA* 100: 6263-6268

- Yu J, Hu S, Wang J, Wong GK, Li S, Liu B, Deng Y, Dai L, Zhou Y, Zhang X, Cao M, Liu J, Sun J, Tang J, Chen Y, Huang X, Lin W, Ye C, Tong W, Cong L, Geng J, Han Y, Li L, Li W, Hu G, Huang X, Li W, Li J, Liu Z, Li L, Liu J, Qi Q, Liu J, Li L, Li T, Wang X, Lu H, Wu T, Zhu M, Ni P, Han H, Dong W, Ren X, Feng X, Cui P, Li X, Wang H, Xu X, Zhai W, Xu Z, Zhang J, He S, Zhang J, Xu J, Zhang K, Zheng X, Dong J, Zeng W, Tao L, Ye J, Tan J, Ren X, Chen X, He J, Liu D, Tian W, Tian C, Xia H, Bao Q, Li G, Gao H, Cao T, Wang J, Zhao W, Li P, Chen W, Wang X, Zhang Y, Hu J, Wang J, Liu S, Yang J, Zhang G, Xiong Y, Li Z, Mao L, Zhou C, Zhu Z, Chen R, Hao B, Zheng W, Chen S, Guo W, Li G, Liu S, Tao M, Zhu L, Wang J, Zhu L, Yuan L, Yang H (2002) A draft sequence of the rice genome (*Oryza sativa* L. ssp. *indica*). *Science* 296: 79-92
- Yu JK, Dake TM, Singh S, Benscher D, Li W, Gill BS, Sorrells ME (2004) Development and mapping of EST-derived simple sequence repeat markers for hexaploid wheat. *Genome* 47: 805-818
- Yu JK, La Rota M, Kantety V, Sorrells ME (2005) EST derived SSR markers for comparative mapping in wheat and rice. *Mol Genet Genomics* 271: 742-751
- Yuan Q, Liang F, Hsiao J, Zismann V, Benito MI, Quackenbush J, Wing R, Buell R (2000). Anchoring of rice BAC clones to the rice genetic map *in silico*. *Nucleic Acids Res* 28: 3636-3641
- Zane L, Bargelloni L, Patarnello T (2002) Strategies for microsatellite isolation: a review. *Mol Ecol* 11: 1-16
- Zeid M, Yu JK, Goldowitz I, Denton ME, Costich DE, Jayasuriya CT, Saha M, Elshire R, Benscher D, Breseghello DF, Munkvold J, Varshney RK, Belay G, Sorrells ME (2010) Cross-amplification of EST-derived markers among 16 grass species. *Field Crops Res* 118: 28-35
- Zhang H, Nasuda S, Endo TR (2000) Identification of AFLP markers on the satellite region of chromosome 1BS in wheat. *Genome* 43: 729-735
- Zhang LY, Bernard M, Leroy P, Feuillet C, Sourdille P (2005) High transferability of bread wheat EST-derived SSRs to other cereals. *Theor Appl Genet* 111: 677-687
- Zhang LY, Rvel C, Bernard M, Balfourier F, Leroy P, Feuillet C, Sourdille P (2006) Transferable bread wheat EST-derived SSRs can be useful for phylogenetic studies among the Triticeae species. *Theor Appl Genet* 113: 407-418
- Zhang P, Li W, Fellers J, Friebe B, Gill BS (2004) BAC-FISH in wheat identifies chromosome landmarks consisting of different types of transposable elements. *Chromosoma* 112: 288-299
- Zhou C, Xia G, Zhi D, Chen Y (2006) Genetic characterization of asymmetric somatic hybrids between *Bupleurum scorzoniferifolium* Willd and *Triticum aestivum* L.: potential application to the study of the wheat genome. *Planta* 223: 714-724

Zhou S, Wei F, Nguyen J, Bechner M, Potamouisis K, Goldstein S, Pape L, Mehan MR, Churas C, Pasternak S, Forrest DK, Wise R, Ware D, Wing RA, Waterman MS, Livny M, Schwartz DC (2009) A single molecule scaffold for the maize genome. *PLoS Genet* 5: e711