

CONTENTS

Chapter	Page No.
1. INTRODUCTION	14-18
2. REVIEW OF LITERATURE	19-42
2.1. EST and Genomic Sequences Databases	
2.2. Molecular Markers	
2.2.1. Sources and properties of DNA-based molecular markers	
2.2.2. Different classes of molecular markers	
2.2.2.1. Hybridization-based markers	
2.2.2.2. PCR-based markers	
2.2.3. Simple sequence repeats (SSRs): The markers of choice	
2.2.3.1. Expressed sequence tag-simple sequence repeats (EST-SSRs)	
2.2.3.2. Genomic SSRs (gSSRs)	
2.2.4. Uses of molecular markers	
2.2.4.1. Genetic mapping	
2.2.4.2. Physical mapping	
(A) Marker-based physical maps	
(a) Physical mapping using pseudodominance	
(b) Physical mapping using acrotrisomics and metatrisomics	
(c) Physical mapping using translocation break-points (TBs)	
(d) Physical mapping using <i>In-situ</i> hybridization (ISH)	
(e) Physical maps based on deletion stocks	
(f) Radiation hybrid mapping (RHM)	
(B) BAC-based physical maps for whole genome sequencing	
2.2.5. Use of genetic/physical maps	
2.2.5.1. Analysis of synteny and colinearity	
2.2.5.2. Identification of gene-rich and gene-poor regions	
2.2.5.3. Gene tagging	
2.2.5.4. Map-based cloning	
2.3. Wheat Genome Sequencing	

3. MATERIALS AND METHODS

43-57

3.1. Cytogenetic Stocks

3.2. Database Used

3.2.1. Sequence for physical mapping

3.2.2. SSR primer sequences for physical mapping

3.2.3. Sequences for the study of orthology

3.2.4. Sequences for synteny analysis

3.3. Available Mapped SSRs for Integrated Physical Map

3.4. Chemicals and Reagents

3.5. Solutions and Buffers

3.6. *In Silico* Approach

3.6.1. SSR mining and primer designing

3.6.2. Criteria used for sequence comparison in BLAST output

3.6.3. Assignment of putative function to orthologs

3.6.4. Construction of consensus maps of wheat genome

3.6.5. Identification of centromere location

3.6.6. Estimation of EST density

3.6.7. Estimation of nonsynonymous vs. synonymous substitution rates (Ka/Ks)

3.7. Wet-Lab Approach

3.7.1. Methodology for physical mapping of SSRs

3.7.2. DNA extraction, purification and quantification

3.7.3. PCR analysis

3.7.4. Polyacrylamide gel electrophoresis (PAGE)

3.7.5. Silver staining of polyacrylamide gel

3.8. Statistical Tests for Random Distribution of Mapped Loci

3.9. Comparison of Physical and Genetic Maps

4. RESULTS AND DISCUSSION

58-133

4.1. Physical Mapping of SSRs

4.1.1. Physical mapping of gSSRs and EST-SSRs

4.1.2. Distribution of genomic and EST-SSR loci in wheat genome

4.1.3. Transferability of brachypodium SSRs to wheat

- 4.2. Integrated Physical Map of SSRs
 - 4.2.1. Distribution of SSR loci on three sub-genomes and seven homoeologous groups
 - 4.2.2. Distribution of SSR loci on individual chromosomes and their arms
 - 4.2.3. Multilocus SSRs in integrated physical maps
 - 4.2.4. Genetic-physical map comparisons
 - 4.2.5. Perturbation of shared SSR loci in genetic and physical maps
 - 4.2.6. Genetic-physical map: recombination distribution
 - Group 1 chromosomes
 - Group 2 chromosomes
 - Group 3 chromosomes
 - Group 4 chromosomes
 - Group 5 chromosomes
 - Group 6 chromosomes
 - Group 7 chromosomes
- 4.3. Orthologous Relationship Among *Brachypodium*, Wheat and Rice
 - 4.3.1. Analysis of brachypodium EST contigs that matched mapped wheat EST contigs
 - 4.3.2. Orthologous relationship between bEST contigs and rice genome sequences
 - 4.3.3. Conserved orthologous sequences among brachypodium, wheat and rice
 - 4.3.4. Common SSRs among the three genomes
- 4.4. Synteny Analysis Between Wheat and *Brachypodium*
 - 4.4.1. Matching of mapped wESTs with brachypodium genome sequences
 - 4.4.2. Syntenic relationship of individual wheat consensus chromosomes with brachypodium chromosomes
 - Wheat consensus chromosome 1 (WC1)
 - Wheat consensus chromosome 2 (WC2)
 - Wheat consensus chromosome 3 (WC3)
 - Wheat consensus chromosome 4 (WC4)
 - Wheat consensus chromosome 5 (WC5)
 - Wheat consensus chromosome 6 (WC6)
 - Wheat consensus chromosome 7 (WC7)
 - 4.4.3. Colinear syntenic regions/blocks between wheat and brachypodium genomes

4.4.4. Consensus chromosome bins containing high density of ESTs showing homology with brachypodium	
4.4.5. Divergence between wheat and brachypodium homologs	
4.4.6. Comparison of wheat and brachypodium genomes in relation to the intermediate ancestral genome	
4.4.7. Wheat ESTs uniquely mapped to specific bins in wheat and their use for inferring relation with brachypodium genome	
4.4.8. Comparison of wheat-brachypodium synteny with previously reported wheat-rice synteny	
4.5. Future Prospects	
5. SUMMARY	134-138
6. REFERENCES	139-172
SUPPLEMENTARY TABLE 1. Details of the bin map position of ESTs on individual wheat chromosomes of seven homoeologous groups, their relationship with sequences of individual brachypodium chromosomes and bin map position in wheat consensus chromosomes	173-223
APPENDIX I. Details of cytogenetic stocks	223-230