

LIST OF TABLES

Table 1. A summary of the number of ESTs and GSSs belonging to genomes of some important plant species available in NCBI database

Table 2. Comparison of commonly used DNA-based molecular markers in crops

Table 3. Summary of the types and number of marker loci genetically mapped in bread wheat

Table 4. A summary of the available deletion based physical maps of bread wheat

Table 5. A summary of radiation hybrid mapping studies in plants

Table 6. A list of genes already cloned through map-based cloning in wheat

Table 7. A list of cytogenetic stocks used for physical mapping of SSRs in wheat

Table 8. Details of the SSR primers used for physical mapping during the present study

Table 9. Actual size of five individual brachypodium chromosomes

Table 10. List of chemicals and reagents used during the present study

Table 11. List of different solutions and buffers with their compositions

Table 12. Composition of reaction mixture and cycling profile used in PCR amplification reaction

Table 13. Details of the gSSRs and EST-SSRs used and the number of loci physically mapped in wheat

Table 14. Distribution of 128 gSSR and EST-SSR loci (mapped during the present study) according to their assignment to wheat chromosomes arranged in a two-way classification

Table 15. Distribution of 2,031 mapped gSSR and EST-SSR loci on the three sub-genomes and seven homoeologous groups in the integrated physical map of wheat

Table 16. SSR loci showing perturbations in their map positions between genetic and physical maps

Table 17. Detailed analysis of comparison between genetic and physical maps of wheat

Table 18. Distribution of the ortholoci according to their assignment to wheat chromosomes arranged in two-way classification

Table 19. Homology between brachypodium supercontigs and homoeologous groups of wheat

Table 20. Distribution of the orthologous loci on individual rice chromosomes

Table 21. Conserved brachypodium EST contigs, their location on wheat and rice chromosomes and their annotated functions

Table 22. List of primers for the 12 conserved SSRs used in wet-lab experiment

Table 23. Genome-wide homology between 8,210 mapped wheat EST sequences with brachypodium chromosomes

Table 24. Colinear syntenic blocks between wheat and brachypodium genomes

Table 25. Details of consensus wheat chromosome bins containing high number of ESTs showing homology with brachypodium

Table 26. Frequency distribution of nonsynonymous and synonymous substitutions for 153 wheat-brachypodium homologs

Table 27. Distribution of single bin wESTs on individual brachypodium chromosomes

Table 28. A summary of the prominent features of chromosome relationships among wheat, brachypodium and rice