

Chapter 7

REFERENCES

REFERENCE

- Abbasi, M., Termeh, F. and Khodaparast, S.A. (2004).** Host range and distribution of *Blumeria graminis* in Iran. *Rostaniha* **5**: 83-85.
- Adkins, S.W. and M.S., Sowerby. (1996).** Allelopathic potential of the weed, *Parthenium hysterophorus* L. in Australia. *Plant Protection Quarterly*, **11**: 20-23.
- Agrwal, G.P., Neema, K.G. and Beligram, R. (1959).** Fungi causing plant disease at Jabalpur (M.P.)-I. *Proc. Nat. Acad. Sci.* **43**:50.
- Akram, M. (1977).** Host specialization studies of *S. fuliginea* (Schlecht) *Poll. Act. Bot. I. nd.* **5**:87-88.
- Akram, M. and A.M., Khan. (1985).** Host range studies of *Sphaerotheca fuliginea* (Schlecht) *Poll. II. Journ. of Indian Bot. Soc.* **64(4)**:386-389.
- Akhundov. T.M. (1982).** Powdery mildew fungi in the Northeast Azerbazan *Bio. Abs.* **70**:6904.
- Alcorn, J.L. (1969).** Infection experiment with cucurbit powdery mildew. *Aust. J. Sci.* **31(1)**: 296-297.
- Amano, K. (1986).** Host range and geographical distribution of the powdery mildew fungi. *Jap. Sci. Soc. Press.* Japan.

Reference


- Anonymous. (1950).** List of common name of Indian plant disease. *Ind. J. Agr. Sci.* **20**:107-142.
- Anonymous. (1950).** Powdery mildew of cucumber pumpkins and related plants. *Agr. Gaz. New South Wales.* **41**:587-588.
- Arya, H.C. and M.S., Gremamat. (1953).** Occurrence of powdery mildew of wheat in the nighbour hood of Jodhpur. *Ind. Phytopath.* **6**:123-130.
- Bajwa, R., Safique, S., Anjum, T., and Safique, S. (2004)** International journal of agriculture and biology, **3**:511-516.
- Baker, K.F. and W.F., Locke. (1946).** Perithecia of powdery mildew on Zinnia . *Phytopath.* **36**:379-380.
- Bakshi, B.K., M.A.R., Ram., Y.N., Puri and S., Singh. (1972).** Survey of the disease of important FRIC, Dehradun, India. pp. 93-98.
- Ballantyne, B.J. (1963).** A preliminary not on the identity of cucurbit powdery mildew of the *Aust. J. Sci.* **25**:360-361.
- Bessey, E.A. (1943).** Notes on Hawaiian fungi papers *Mich. Acad. Sci. Arts. Letters.* **28**:3-8.
- Bessey, E.A. (1961).** Morphology and taxonomy of fungi. Hafner, New York.

Reference

- Bhagyanaryana, G.A., S., Rao and P., Ranachar. (1988).** Powdery mildew (Erysiphales) from Hyderabad (A.P.) Kavaka. **16:45-49.**
- Bhatnagar, et al. (1961).** *Indian J. Med. Res.* **49:807.**
- Bhatia, J. N. and Thakur D. P. (1989).** Field evaluation of systemic and non systemic fungicides against powdery mildew of different economic crops. *Indian Phytopath.* **42 (4): 573.**
- Blumer, S. (1922).** Die formen der *Erysiphe cichoracearum* D.C. *Cantabl Bakt* (Abt.2). **57:45-60.**
- Blumer, U. (1948).** Beitrag Zur. Kenntrisder Erysiphaceae Ber. Schweiz. Bot. Gaz. **58:61-68.**
- Blumer, S., (1967).** Eche Mehлтаupilge (Erysiphace) Veb. Gustav. Fischer. *Verlag. Jena. Pub.* pp. 334-403.
- Blumer, S. (1974).** Contribution to the knowledge of the Erysiphales *Bio. Abs.* **57:677.**
- Boesewinkle, H.J. (1979).** Observation on the host range of powdery mildews. *Phytopath. Z.* **94: 241-248.**
- Boesewinkel, H.J. (1980).** The morphology of imperfect states of powdery mildew (Erysiphaceae). *Bot. Rev.* **46:167-224.**

Reference

- Braun, U. (1980).** Morphological studies in the genus *Oidium* flora. **170**:77-90.
- Braun U. (1987).** A monograph of the *Erysiphales* (powdery mildews). Beihefte zur Nova Hedwigia. **89**: 1-700.
- Braun, U. (1995).** The Powdery Mildews (Erysiphales) of Europe. *Gustav Fischer-Verlag, Jena, Germany.*
- Braun, U. and Minnis, A.M. (2008).** The nomenclature of *Erysiphe clandestina* (*Uncinula clandestina*) revisited. *Schlechtendalia*. **17**: 55-57.
- Braun, U., Takamatsu, S., Heluta, V., Limkaisang, S., Divarangkoon, R., Cook, R. and Boyle, H. (2006).** Phylogeny and taxonomy of powdery mildew fungi of *Erysiphe* sect. *Uncinula* on *Carpinus* species. *Mycological Progress* **5**: 139–153.
- Braun, U., Cook, R.T., Inman, A.J. and Shin, H.D. (2002).** The taxonomy of the powdery mildew fungi. in RR Belanger *et al.* (eds.). The powdery mildews, A comprehensive treatise. The American Phytopathological Society, *St. Paul,, Minnesota, USA.* pp 13-55
- Bosnic, A.C. and C.J., Swanton. (1997).** Influence of barnyard grass (*Echinochola crus-galli*) time of emergence and density on corn (*Zea mays*). *Weed Sci.*, **5**: 276-282



Reference

- Bacigalova, K. and Markova, J. (2006).** *Erysiphe azaleae* (*Erysiphales*) – a new species of powdery mildew for Slovakia and further records from the Czech Republic. – *Czech Mycol.* **58**(3–4): 189–199.
- Braun U. (1987).** A monograph of the *Erysiphales* (powdery mildews). – Beih. Nova Hedwigia **89**: 1–700.
- Butlar, E.J. and Bisby, G.R. (1931).** The fungi of India. *Inp.* Council of Agric. Res. India, *Sci. Monogr.* I.XVIII+237pp.
- Chippendale, J.F. and F.D., Panetta. (1994).** The cost of *Parthenium* weed to the Queens land cattle industry. *Plant Protect.*, **9**: 73-76.
- Chopra, et al. (1958).** Wealth of India. pp. **320**.
- Chowdhury (1982).** *Sci. Cult.* **38**:358.
- Choi, H.W., Y.J., Choi, D.S., Kim, I. S., Hwang, D. S., Choi, N., H., Kim, D.H., Lee, H.D., Shin, J. Nam and B. K., Hwang. (2009).** First Report of Powdery Mildew Caused by *Erysiphe cruciferarum* on *Arabidopsis thaliana* in Korea. *Plant Pathol. J.* **25**(1): 86-90.
- Chauhan, S., A., Srivastava and K., Singh. (2010).** Study on the incidence of powdery mildew disease in agro climatic conditions. *ARPN Journal of Agricultural and Biological Science.* Vol.5. No. 3.
- Clare, B.G. (1958).** The identity of the cucurbit powdery mildew of south eastern Queensland *Aust. J. Sci.* **20**:273.

Reference

- Clare, B.G. (1964).** Erycephaceae of south eastern Queensland. Univ. Queensland papers. 4:111-144.
- Cole, J.S. (1966).** Powdery mildew of tobacco (*E. cichoracearum* D.C.) *Ann. App. Bio.* 57:201-209.
- Caroline, A.H., M.J., Renfrew and M.W., Woolridge. (2001).** Assessing the risks of pesticide residues to consumers: Recent and future developments. *Food Additives and Contaminants*, 18: 1124-1129.
- Das and Misra (1987).** *Indian J. For.* 10:301.
- Deslandes, J.A. (1954).** Studies and observations on lettuce powdery mildew. *Pl. Dis. Rep.* 38:560-562.
- Denchev, M.C. (2005).** New records of fungi, fungus-like organisms, and slime moulds from Europe and Asia. *Mycologia Balcanica.* 5: 93-96.
- Dey (1969).** *Indian For.* 95:551.
- Dhanwantri, B.N. and W.R., Jarvis. (1985).** Powdery mildew (*Erysiphe cichoracearum*) of green house lettuce on Ontario. *Pl. Dis.* 69:77.
- Dhar, et al. (1968).** *Indian J. exp. Biol.* 6:245.
- Diazfranco, A. (1985).** Effect of sunflower powdery mildew in the north of Tamaulipas. *Rev. Pl. Path.* 64:211.

Reference

- Drozdovaskaya, L.S. (1980).** Powdery mildew of medicinal species in Groundsel. *Bio. Abs.* **69**:582.
- Dhileepan, K., S.D., Setter and R.E., Mc Fadyen. (2000).** Impact of defoliation by the bio-control agent *Zygogramma bicoloration* the weed *Parthenium hysterophorus* in Australia. *Biocontrol.* **45**: 501-512.
- Dugan, F.M. and D.A., Glawe. (2007).** Powdery mildew on weeds in the Pacific Northwest: A miscellany of new records. *Pacific Northwest Fungi.* **2** (1): 1-7.
- Ellett, W.C. (1963).** Host range of the Erysiphaceae of *Ohio*. Vol. **66**. No.6. 570-581.
- Evans, C.H. (1997).** *Parthenium hysterophorus*: A review of its weed status and the possibilities for biological control. *Biocontrol/news and Information.* **18**: 389-398.
- Fallon, R.E., P.W., Southerland, and I.C., Hallet. (1989).** Morphology of *Erysiphe pisi* on leave of *Pisum sativum*. *Can. J. Bot.* **67**:3410-3416.
- Frank, M., Dugan. (2006).** First Report of Powdery Mildew on *Dipsacus sylvestris* Caused by *Sphaerotheca dipsacearum* in North America. *Plant Health Progress* doi:10.1094/PHP-2006-0607-02-BR.
- Glawe, D. A., Grove, G. G., and Nelson, M. (2003).** First report of powdery mildew of *Convolvulus arvensis* caused by *Erysiphe convolvuli* var.

Reference

- convolvuli* in the Pacific Northwest. Online. Plant Health Progress doi:10.1094/PHP-2006-0405-01-BR.
- Glawe, D. A., du Toit, L. J., and Pelter, G. Q. (2004).** First report of powdery mildew on potato caused by *Leveillula taurica* in North America. Online. Plant Health Progress doi:10.1094/PHP-2004-1214-01-HN.
- Glawe, D. A. and Dugan, F.M. (2006).** First report *Erysiphe flexusa* in Western North America. Pacific North West Fungi. **1**:1-11.
- Gardner, M.W. and C. E., Yarwood. (1974).** List of powdery mildew of California. *Calif. Agric. Exp. Sta. Leaflet No.* 217 p 31.
- Gorter, G.J.M.A. and A., Eicker. (1983).** First report of perfect stage of some powdery mildew fungi in south Africa *Rev. Pl. Path.* **62**:441.
- Gorter, G.J.M.A. and A., Eicker. (1986).** South African record of *Erysiphe* from *Transmal .Bio. Abs.* **81**: Ref. No. 79407.
- Gorter, G.J.M.A. (1989).** Identification of South African Erysiphaceae with a key to the species. *Rev. Pl. Path.* **68**:216.
- Graf and Martin, A. (1934).** Studies on powdery mildew of cereals. *Corn. Univ. Agr. Exp. Sta. Mem.* 157.
- Gritten, E.T. and Ebert, R.D. (1975).** Interaction of planting date and powdery mildew on pea plant performance. *Journal of the American Society for Horticultural Science* **100**, 137 – 42.

Reference

- Handa, S., B. Sahoo and V.K., Sharma. (2001).** Oral hyposensitisation in patients with contact dermatitis from *Parthenium hysterophorus*. *Contact Dermatitis.*, **44** : 279-82
- Hammett, K.R.N. (1977).** Taxonomy of Erysiphaceae in New Zeland. *N. Zeland. J. Bot.* **15**:687-711.
- Hammernton, J.L. (1981).** Weed problems and weed control in the commonwealth. *Caribbean Tropical Pest Manage.* **27**: 379-387.
- Hansford, C.G. (1961).** The Meliolineae. *Sydowia. Ser-II. Beih.* **2**:1-806.
- Hassan, S. (1974).** Host specialization of powdery mildew, *Erysiphe cichoracearum*, from *Chondrilla juncea*. *Aust. J. Agric. Res.* **25**(3): 459 – 465.
- Hartman, J. (2007).** Phlox powdery mildew. Plant pathology fact sheet.
- Hirata, K. (1955).** Host range and geographical distribution of the powdery mildew, Memoo. Niigata. *Univ. Japan.*
- Hirata, K. (1968).** Notes on host range and geographic distribution of the powdery mildew fungi. *Trans. Mycol. Soc. Japan.* **8**:73-88.
- Hopper et al. (1990).** *Eur. J. Med. Chem.* **25**:717.
- Husain, I.S. and M., Akram. (1997).** Distribution pattern and identity of powdery mildew of composities in the districts of Agra division. *Indian Phytopath.* **50**(2): 250-255.

Reference

- Ialongo, M.T. (1981).** A Population of *E.cichoracearum* D.C. ex. Merat. Specific to *Sonchous* Rev. Pl. Path. **59**:247.
- Ialongo, M.T. (1987).** Taxonomic study of some species of the genus *Erysiphe*. Mycotaxon **44**:251-256.
- Jankovitis, T. and O., Szentivanyi. (2006).** First report of powdery mildew on *Sedum alboroseum* in Europe. Plant Pathology. Volume **55**, Number **2**: 297-297.
- Javaid, A., T., Anjum and R., Bajwa. (2005).** Biological control of *Parthenium* II: Allelopathic effect of *Desmostachya bipinnata* on distribution and early seedling growth of *Parthenium hysterophorus* L. Int. J. Biol. Biotech. **2**: 459-463.
- Javaid, A., S., Shafique and S., Shafique. (2006).** *Parthenium* weed – an emerging threat to plant biodiversity in Pakistan. Int. J. Biol. Biotech. **3**(3): 619-622.
- Javaid, A., S., Shafique and S., Shafique. (2007).** Causes of rapid spread of *Parthenium hysterophorus* L. in Pakistan and possible control measures- A Review. Pak. J. Bot., **39**(7): 2611-2618.
- Jhooty, J.S. (1967).** Identity of powdery mildew of cucurbits in India. Pl. Dis. Rep. **51**:1079-1080.

Reference

- Jayachandra. (1971).** Parthenium weed in Mysore state and its control. *Current Sci.*, **40**: 568-569.
- Kable, P.F. and B.J., Ballantyne. (1963).** Observation on the cucurbits powdery mildew in the Ithaca district. *Pl. Dis. Rep.* **47**: 482.
- Kamal, Singh, S. and Singh, R.P. (1977).** Fungi of Gorakhpur-I. *Indian Phytopath.* **30**:186-188.
- Kamat, M.N. and M.K., Patel. (1948).** Some new hosts of *Oidiopsis taurica* (Lev.) Salmon in Bombay. **1** (2): 153-158.
- Kanchan, S.D. and Jayachandra. (1979).** Allelopathic effects of *Parthenium hysterophorus* L. exudation of inhibitors through roots. *Plant and Soil*, **53**: 27-35.
- Kanchan, S.D. and Jayachandra. (1980).** Allelopathic effects of *Parthenium hysterophorus* L. II. Leaching of inhibitors from aerial vegetative parts. *Plant and Soil*, **55**: 61-66.
- Kapoor, et al. (1992).** *J. Sci. Ind. Res.* **51**:8.
- Kapur. (1991).** *Indian Drugs.* **28**:210.
- Kasturi, et al. (1973).** *Indian J. Chem.* **11**:91.
- Keneth, R.G. and J., Palti. (1984).** The distribution of downy and powdery mildew and rust over tribes of Asteraceae. *Mycologia.* **76**: 705-718.

Reference

- Khan, B.M., M.A., Khan, N., Anwar, and A., Anees. (2008).** *Parthenium hysterophorus*: A Potential Source of Bioherbicide. *Pak. J. Bot.*, **40**(5): 1933-1942.
- Khan, M.W., K.A., Malik and A.M., Khan. (1975).** Perithecial stage of setrtain powdery mildews including some new records *III. Ind. Phytopath.* **28**: 199-210.
- Khan, M.W., M. Akram, A. Khan and A.M., Khan. (1977).** Occurrence of powdery mildew at Aligarh and adjoining areas. *Act. Bot. Ind.* **5**: 139-142.
- Khan, M.W., M., Akram and A.M., Khan. (1971).** Status of Cucurbit powdery mildew. I. Perithecial production in cucurbit powdery mildew in Northern India. *Indian Phytopath.* **23**:497-502.
- Khan, M.W., and E.A. Mussa. (1979).** Powdery mildw in the Lybian Jamahiriya. Identity of certain powdery mildew fungi. *Libiyan J. Agr.* **8**: 161-167.
- Khan, M.W. (1980).** Powdery mildew in the Libiyan Jamahiriya II. Identity of certain powdery mildew fungi. *Libiyan J. Agr.* **9**: 137-142.
- Khan, M.W. (1982).** Powdery mildew in the Libiyan Jamahiriya IV. Identity of certain powdery mildw fungi. *Libiyan J. Agr.* **11**: 127-132.
- Khan, M.W. and I.S., Faraj. (1982).** Powdery midew in the Libiyan Jamahiriya III. Identity of certain powdery mildew fungi. *Libiyan J. Agr.* **11**: 121-125.

Reference

- Khan, M.W. and G.K., Sharma. (1995).** Taxonomic evaluation of anamorph characters in identification of powdery mildew fungi on cucurbits. *Indian Phytopath.* **48:** 314-24.
- Khosla, S.N. and S.N., Sobti. (1979).** Parthenium: A national health hazard. its control and utility a review. *Pesticides*, **13:** 121-127.
- Khodaparast, S.A., Hedjaroude, G.A., Ershad, D., Zad, J., Termeh, F., Moosavi, M. (2001).** A Study on the identification of *Erysiphaceae* in Guilan province, Iran (II). *Rostaniha* (2): 45-52.
- Khosla, S.N. and S.N., Sobti. (1981).** Effective control of *Parthenium hysterophorus* L. *Pesticides*, **15:** 18-19.
- Khosla and Sobti. (1980).** *Herba. Hung.* **19** (2):73.
- Kiss, L. (2003).** A review of fungal antagonists of powdery mildew and their potential as biocontrol agents. *Pest Management Science.* **50:** 475-483.
- Klemm, H. (1986).** A notable injurious occurrence of powdery mildew (*Erysiphe cichoracearum* DC. Ex. Merat.) on Lettuce (*Lactuca sativa*) in Germany. *Rev. Pl. Path.* **65:** 51.
- Krasevicz, D.M. and Zitter, T.A. (1996).** Powdery mildew occurrence on green house tomato plants in New York. *Plant Disease.* **80:**709.

Reference

- Kumar, G. and N., Gautam. (2008).** Allelotoxicity of *Parthenium* leaf extracts on cytomorphological behaviour of sunflower (*Helianthus annuus*). *J. Environ. Biol.*, **29**: 243-247.
- Kumar, A., V.C., Verma, S.K., Gond, V., Kumar and R.N., Kharwar. (2009).** Bio-control potential of *Cladosporium* sp. against a noxious weed *Parthenium hysterophorus* L. *J. Environ. Biol.* **30**(2): 307-312.
- Kunkaliker, S. and Pdaganur, G.M. (1994).** *Euphorbia hirta* a new collateral host of *E. polygony*. *Indian Phytopathology.* **47**(4):440-441.
- Kunoh, H. (1995):** Host-parasite speci@city in powdery mildews. In: Kohmoto. K., U. S. Singh and R. P. Singh (eds), *Pathogenesis and Host Speci@city in Plant Diseases, Vol. II, Eukaryotes*, pp. 239±250. Pergamon, Elsevier Science, Oxford, UK.
- Lakshmanan, P. and A., Sudha. (2007).** *Solanum nigrum*, a new host for powdery mildew disease of *Capsicum annuum*. *Australasian Plant Disease Notes.* **2**: 97–98.
- Lakshmi, C. and C.R., Srinivas. (2007a).** *Parthenium* dermatitis caused by immediate and delayed hyper sensitivity. *Contact Dermatitis.* **57**: 64- 65.
- Lakshmi, C. and C.R., Srinivas. (2007b).** Type I hypersensitivity to *Parthenium hysterophorus* in patients with *Parthenium* dermatitis. *Ind. J. Dermatol Veneriol. Leprol.* **73**: 103-105.

Reference

- Lazzeri, L., D., Rongai and C., Cerato.(2009).** A natural fungicide for the control of *Erysiphe betae* and *Erysiphe cichoracearum*. *Eur. J. Plant Pathol.* **124**:613–619.
- Liu, T.J. and Y.Z., Shang. (2008).** *Erysiphe alashanensis* sp. Nov. from Inner Mangolia, China. *Nova Hedvigia.* **86**: 255-258.
- Lebeda, A. and J., Buczkwski. (1986).** Occurrence of *Erysiphe cichoracearum* perithicia on wild *Lactuca* speceis. *Phytopath. Z.* **115**: 21-28.
- Lebeda, A., Sedlarova, M., Jankovski, L., and Shin, H. D. (2006).** First report of rhododendron powdery mildew on *Rhododendron* spp. in Czech Republic. – New Disease Reports 14.
- Lebeda, A., E., Krrioastkova, V., Rybka and P., Havraanek. (2001).** *J. Phytopathology* **149**: 207-212.
- Lebeda, A., Mieslerova, B. and Dolizalova, I. (2002).** The first record and characterization of powdery mildew (*Erysiphe pachypodiae* sp. Nov.) on *Pachypodium lamerei* (Apocynaceae). *J. Phytopathology.* **150**: 149–154.
- Leyvkh. (1940).** Chemical method in the control of powdery mildew of tobacco. pp. 78-96.
- Liabach, F. (1930).** Ueber Die Bedingungen Der Perithezienbildung Bei Den Erysipheen. *Jahrub. Wiss. Bot.* **72**:106-136.
- Linnaeus, C. (1753).** Species Plantarum. **11**:1186.

Reference

- Lonkar, A., J.C., Mitchell and C.D., Calna. (1974).** Contact dermatitis from *Parthenium hysterophorus*. Transactions of the St. John's Dermatological.. **60:** 43-49.
- Mathur, R.L., B.L., Mathur and L.P., Bhargava, (1971).** New host records of powdery mildew from Rajasthan. *Indian Phytopath.* **24:** 63-66.
- McDonald, J.A. (1939).** Powdery mildew on Cinenarias. *Res. Abs. Mycol.* **29:** 459.
- McKeen, W.E., R., Smith and N., Mitchell. (1966).** The houstorium of *E. cichoraciarum* and host parasite interface on *Helianthus annuus*. *Can. J. Bot.* **44:** 1299-1305.
- Mahadevappa, M. (1997).** Ecology, distribution, menace, and management of *Parthenium*. In: *Proc. First International Conference on Parthenium Management*, UAS, Dharwad, **1:** 1-12.
- Maharshi, R.P. (1984).** *Leveillula taurica* on *Peristrophe bicalyculata*. *Indian Phytopath.* **37:** 587.
- Mamulk, O.F., H.C. Weltzien, (1973).** Untersuchungen über die Hauptfruchtform des Echten Rübenmehltaus, *Erysiphe betae* (Vanha) Weltzien. I. Fruchtkörperbildung in Abhängigkeit von Umweltfaktoren. *Phytopathol. Z.* **76:** 221-252.
- Matheron, M.E. and M., Porchas. (2007).** Evaluation of Fungicides for Management of

Reference

Powdery Mildew on Lettuce. Vegetable Report (P-152).

Meloide, L.P. and D.A., Glawe. (2007). North American records of anomorphic powdery mildew fungi (Erysephale) parasitizing species of *Leucothoe* and *Lemnanthis*. *Pacific North West Fungi*. **2**:1-6.

Minev, K. (1957). Investigation of the biology of tobacco powdery mildew *E. cichoracearum* D.C. *Fac. Agr. and forestry Univ. Scopie*. **10**: 5-72.

Milovtzoa, M.O. (1937). New species of fungi on the medicinal and essential oil plants of Ukraine. *Trans. Institute Bot. Institute of Kharkoff*. **2**:7.

Mittal, S. and M., Akram. (1984). Reaction of different cultivated umbeliferous plants against culture of powdery mildew *Erysiphe umbeliferarum* D E Bary. (Abs.). *Sci. Cong. No. 76*. Part. III.

Mittel, J.H. and Tondon, R.N. (1930). Fungus flora of Allahabad. *J. Ind. Bot. Soc.* **9**:190-198.

Mohiddin, et al. (1991). *Int. J. Pharmacogn.* **29**:252.

Morrison, R.M. (1961). A study of *E. cichoraciarum* D.C. *Ex. Merat*. On detached leaf culture *Dis. Abs.* **21**: 2448-2449.

Moyer, C. and N.A., Peres. (2008). Evaluation of Biofungicides for Control of Powdery Mildew of *Gerbera Daisy*. *Proc. Fla. State Hort. Soc.* **121**:389–394.

Reference

- Moyer, C., N. A. Peres, L. E. Datnoff, E. H. Simonne, and Z. Deng. (2004).**
Evaluation of Silicon for Managing Powdery Mildew on Gerbera Daisy.
Journal of Plant Nutrition, **31**: 2131–2144.
- More, P.R., V.P., Vadlamudi and M.I., Qureshi. (1982).** Note on the toxicity of
Parthenium
hysterophorus in livestock. *Ind. J. Ani. Sci.* **52**: 456-457.
- Munjal, R. L., V. V., Chenulu, and I. S., Hora, (1963).** Assessment of losses
due to powdery mildew *Erysiphe polygoni* on pea. *Indian Phytopath.*, **16**:
260-267.
- Nagaraja, T.J. and S.M., Deshmukh. (2009).** Phytotoxic effect of *Andrographis*
paniculata nees on metabolism of *Parthenium hysterophorus*. *L. Journal of*
Biopesticides, **2(2)**: 165-167.
- Nandakumar and Shetty. ((1989).** *Geobios.* **16**:8.
- Nath, R. (1988).** *Parthenium hysterophorus* L. A review. *Agricultural Reviews*, **9**:
171-179.
- Njoroge, J.M. (1986).** New weeds in Kenya coffee: A short communication.
Kenya Coffee, **51**: 333-335.
- Nour, M.A. (1958).** Studies on *Leveillula taurica* (Lev) Arn. and other powdery
mildew. *Transaction of the British Mycological Society*.**41**:17-38.

Reference

- Oudhia, P. (1998).** Parthenium: A curse for biodiversity of Chattisgarh (India) plains. Abstract. National Research Seminar on Bio-chemical changes. An Impact on Environments, R.D. Govt. P.G. College, Mandala M.P. p. 26.
- Panday and Dubey. (1989).** *Proc. Indian Acad. Sci. (Plant Science)*. **99**:51.
- Patil, S.D. (1964).** Genus *Sphaerotheca* Lev. In Maharashtra. *J. Univ. Poona. Sci. Tech.* **28**:33-34.
- Patel, M.K., Kamat, M.N. and Bhide, V.P. (1949).** Fungi of Bombay Supplement. I. *Indian Phytopath.* **2**:142-155.
- Prasad, S.K. and Rani, P. U. (1981).** Occurrence of powdery mildew on Parthenium caused by *Oidium parthenii*. *Nov. Curr. Sci.* **50**:1081-1082.
- Parwez, M.S. and M., Akram. (1987).** Powdery mildew on *Vernonia cineria* Schreb. A new host from India. *Act. Bot. Ind.* **15**: 319.
- Parwez, M.S. and M., Akram. (1989).** Occurrence of powdery mildew at Aligarh and its adjoining areas. *Acta. Botanica. Indica.* **17**(2): 267.
- Pathak, K.D. and L.M., Joshi. (1972).** Chemical control of powdery mildew of wheat. *Ind. Phytopath.* **25**: 139-141.
- Patwardhan, P.G. (1996).** Some new records of powdery mildew fungi. *Pl. Dis. Reprtr.* **50**:709-710.
- Patel, R. and Sharma, N.D. (1986).** Taxonomy of some powdery mildew fungi III. *Advances in Plant Science Research. Vol. IV*: 1-22.
- Patel, R. and Sharma. (1996).** *Advances in Plant Science Research-IV*: 1-22.

Reference

- Paul, Y.S. and R.L., Munjal. (1982).** New host record of powdery mildew from India. *Indian Phytopath.* **35** (1): 170-171.
- Paul, Y.S. and Thakur, V.K. (2006).** *Indian Erysiphaceae.* pp.134.
- Pawar, P.V., Chudhari, G.K. and Deshmukh, S.A. (2009).** Occurrence of severity of powdery mildew on cucurbits. *International Research Journal.* Vol. II. 0974-2832.
- Phillip, T., Gupta V., G., Bajpai, A.K. and Datta, R.K. (1994).** Disease of mulberry in India-Research priorities and management strategies. *Int. J. Trop. plant Dis.* **12**: 1-21.
- Piatek, M. (2002).** *Erysiphe flexuosa*, a new for Poland powdery mildew causing disease of *Aesculus hippocastanum*. – *Phytopathol. Pol.* **24**: 67–71.
- Plenk, A., U., Holzer and S., Sedlan. (1992).** Powdery mildew fungi in the domestic garden. *Rev. Pl. Path.* **71**: 255.
- Putnam, M.L., and Glawe, D.A. (2007).** New North American records of anamorphic powdery mildew fungi (Erysiphales) parasitizing species of *Leucothoë* and *Limnanthes*. *Pacific Northwest Fungi* **2**:1-6.
- Puzanova, L.A., (1992).** Powdery mildew fungi on plants in Krasnodarkii Kari. *Rev. Pl. Path.* **71**: 101.
- Piatek, M. (2003).** *Chionanthus* (Oleaceae), a new host genus for the powdery mildew *Phyllactinia fraxini* (Erysiphaceae). - *Nova Hedwigia* **77**: 379-381.

Reference

- Rai, et al. (1982).** *Bull. Bot. Surv. India.* **24:87.**
- Rajan, L. (1973).** Growth inhibitor(s) from *Parthenium hysterophorus*. *Curr. Sci.*, **42:** 729-730.
- Rao, P.N. (1961).** *Sphaerotheca* on two members of Euphorbiaceae from Hyderabad (India). *Curr. Sci.* **30:**433-435.
- Rao, R.S. (1956).** Parthenium a new records for India. *J. Bombay Nat. Hist. Soc.*, **54,** 218-220
- Reis, A., Boiteux, L.S., M.L.P., Lema. (2007).** Powdery mildew of ornamental species caused by *Oidiopsis haplophylli* in Brazil. *Summa Phytopathol., Botucatu*, v. **33,** n. **4,** p. 405-408.
- Rassonw, D.J. (1959).** The effect of temperature and humidity on tobacco powdery mildew fungus. *S. Afr. J. Agr. Sci.* **2:** 19-31.
- Rector, B.G., V., Harizanova, R., Safora, T., Wedmer and R.N., Wiedenmann. (2006).** Prospects for biological control of teasels, *Dipsacus* spp., a new target in the United States. *Biological control.* **36:** 1-14.
- Reiling, T.P. (1984).** Powdery mildew. D. J. Hajidon. ed. Compendium of pea disease. The American Phytopathological Society. *St. Paul.* pp. 21-22.
- Romero, A., Carrion, G. and Rico-Gray, V. (2001).** Fungal latent pathogens and endophytes from leaves of *Parthenium hysterophorus* (Asteraceae). *Fungal Diversity* **7:** 81-87.
- Rudloff and Sood (1969).** *Perfume essent oil Rec.* **60:**303.

Reference

- Sachan, S.N. (1977).** A new record of *Uncinula aspera* from India. *Indian Phytopath.* **30**: 126-127.
- Salmon, E.S. (1900).** A monograph of the Erysiphaceae. *Mem. Torrey. Bot. Club.* **91**: 1- 292.
- Sawada, K. (1914).** On the classification on Erysiphaceae by the conidial state. *Formosa. Agr. Exp. Sta. Spec. Rep.* **9**: 1-120.
- Saxena, S. and M., Kumar. (2007).** Mycoherbicidal Potential of *Alternaria alternata* ITCC4896 for the control of *Parthenium hysterophorus*. *Journal of plant protection research.* Vol. **47**, No. 2.
- Saxena, A.K. and S.B., Saksena. (1981).** Powdery mildew disease in Madhya Pradesh. *J. Ind. Bot. Soc.* 60-17.
- Schmitt, J.A. (1955).** The host specialization of *E. cichoraciarum* from *Zinnia*, *Phlox* and Cucurbits. *Mycologia.* **47**: 688.
- Schnathorst, W.C. (1959).** Spread and life cycle of the lettuce powdery mildew fungus. *Phytopath.* **49**: 464-468.
- Schnathorst, W.C. (1990).** Effect of temperature and moisture stress on the lettuce powdery mildew fungus. *Phytopath.* **50**: 304-308.
- Schnathorst, W.C. (1965).** Environmental relationship in the powdery mildew. *Ann. Rev. Phytopath.* **3**: 343-366.
- Sert, H., H., Sumbul and Sterflinger, K. (2006).** New host of powdery mildew in Turkey. *Phytoparasitica.* **34** (5): 474-476.

Reference

- Sethi, et al. (1987).** Phytochemistry. **26**:3359.
- Shabbir, A. and Bajwa, R. (2007).** *Parthenium* invasion in Pakistan- A threat still un-recognized. *Pak. J. Bot.*, **39**(7): 2519-2526.
- Sharma, N.D. and Khare, C.P. (1992).** Morphology of anomorph of some powdery mildew fungi of Jabalpur division. *Acta. Botanica. Indica.* **20**: 269-277.
- Sharma and Bhutani. (1988).** *Planta. Med.* **54**:120.
- Sharma, V.K., G., Sethuraman and R., Bhat. (2005).** Evolution of clinical pattern of *Parthenium dermatitis*: A study of 74 cases. *Contact Dermatitis.* **53**: 84-88.
- Shin, H.D., Park, M.J., J.G., Han and J.Y., Kim, (2000).** Powdery Mildew Outbreak on Kale Caused by *Erysiphe cruciferarum*. *Plant Pathol. J.* **26**(2): 205
- Shukla, R. and Pandey, A. K. (2008).** Pathogenic diversity of *Sclerotium rolfsii* isolates, a potential biocontrol agent against *Parthenium hysterophorus* L. *African Journal of Environmental Science and Technology* Vol. **2** (5), pp. 124-126.
- Sohi, H.S. and S.K., Nayar. (1969).** Some new records of fungi from India-I. *Indian Phytopath.* **22**: 410-421.
- Somani, R.B., Sapkal, K.N., Wangikar, P.D. and Kodmilwar, R.V. (1976).** A new host of *Sphaerotheca fuliginea*. *Indian Phytopath.* **29**:216.

Reference

- Srinivas, C.R. (2006).** Parthenium dermatitis treated with azathioprine weekly pulse doses. *Ind. J. Dermatol. Venereol. Leprol.* **72**: 234-238.
- Srinivas, C.R. (2005).** Transmission of Parthenium dermatitis by clothing. *Arch. Dermatol.* **141** : 1605.
- Singh, K.S. and R.P., Singh. (1977).** Fungi of Gorakhpur-1. *Ind. Phytopath.* **30**: 186-187.
- Singh, S., A., Yadav, R.S., Balyan, R.K., Malik and M., Singh, (2004).** Control of ragweed parthenium (*Parthenium hysterophorus*) and associated weed. *Weed Technol.* **18**: 658-664.
- Sinha and Singh. (1990).** *Sci. Cult.* **56**:169.
- Suriachandraselvan and Narayansamy. (1987).** *Madras. Agric. J.* **74**:154.
- Swaminathan, S., R.S., Rai and K.K. Smesh. (1990).** Allelopathic effects of *Parthenium hysterophorus* L. on germination and growth of a few multipurpose trees and arable crops. *Int. Tree Crops J.*, **6**: 143-150.
- Tang, K., A., Samed, M.A., A.H., Anyul, Azhar, S. B. and Absar, N. (2006).** Nutritional changes of four varieties of mulberry leaves infected with fungus (*Phyllactinia corylea*). *Pak J. Biol. Sci.* **9**(3): 355-359.
- Takamatsu, S. and K., Pastircakova. (2007).** *Erysiphe arcuata* found on European hornbeam (*Carpinus betulus*) is reported from Slovakia for the first time. *J. Phytopathology* **156**: 597–601.

Reference

- Takamatsu, S., Heluta, V., Havrylenko, M. and Divarangkoon, R. (2009).** Four powdery mildew species with catenate conidia infect Galium: molecular and morphological evidence. *Mycol. Res.* 113:117-129.
- Talwar and Kalsi. (1989).** *Phytochemistry.* 28:1091.
- Tamado, T. L., W., Schutz and P., Milberg. (2002b).** Germination of ecology of the weed *Parthenium hysterophorus* in eastern Ethiopia. *Ann. Appl. Biol.*, 140: 263-270.
- Taye, T. and M., Gossmann. (2007).** Exploration of Fungal Pathogens Associated with *Parthenium hysterophorus* in Ethiopia. *Eth. J. of Weed Mgt.* 1(1):79 – 86.
- Tiwari K.R., Panner G.A., Warkentin T.D., Rashid K.Y. (1997).** Pathogenic variation in *Erysiphe pisi* the causal organism of powdery mildew in pea. *Can. J. Plant Pathol.*, 19: 267–271.
- Towers, et al. (1977).** *J. Sci. Ind. Res.* 36:672.
- Trdan, S., Valic, N., Jerman, J., Ban, D., Znidarcic, D. (2004).** Efficacy of Three Natural Chemicals to Reduce the Damage of *Erysiphe cichoracearum* on Chicory in Two Meteorologically Different Growing Seasons. *J. Phytopathol.*, 152, 10: 567-574.
- Trelese, S.F. and H.M., Trelease. (1928).** Susceptibility of wheat to powdery mildew as influenced by salt nutrition. *Bulletin of the Torrey botanical Club.* 55:41-68.

Reference

- Utkhede, R. S., Koch, C. A., Menzies, J. G. and Ehret, D. L. (2001).** Host range of a powdery mildew (*Erysiphe orontii*) on tomato. *Can. J. Plant Sci.* **81**: 179–182.
- Uppal, B.N., Patel, M.K. and Kamat, M.N. (1935).** The fungi of Bombay- VIII. 1-56.
- Verma, K.K., A., Bansal and G., Sethuraman. (2006).** Parthenium dermatitis treated with azathioprine weekly pulse doses. *Ind. J. Dermatol. Venereol. Leprol.* **72**: 24-27
- Voityuk, S.O., Heluta, V.P. and Nevo, E. (2004):** *Neoerysiphe cumminsiana* (Erysiphales, Eumycota), a new powdery mildew fungus in Israel. – *Flora Mediterranea* **14**: 267-273.
- Voityuk, S.O., Heluta, V.P., and Nevo, E. (2004):** *Neoerysiphe galii* (Erysiphales), a powdery mildew fungus new for Israel. *Mycologia Balcanica* **1**: 135–137.
- Wajid, K., M., Akram, M. and Abrar, M., Khan. (1972).** Perithecial stage of some powdery mildew including some new records. *Indian Phytopath.* **25(2)**:220-224.
- Wajid, K., M., Akram, M., K., Azmatullah and Abrar, M., Khan. (1977).** Occurrence of powdery mildews at Aligarh and adjoining areas. *Acta Botanica Indica.* **5**:139-142.

Reference

- Wolcan, S.M. and Y., Sato. (2007).** First report of powdery mildew on *Aloysia* Spp. (Verbinaceae). *Australasian Plant Disease Notes*. **2(1)**: 113–114.
- Walczanska, A. (2006).** First report of *Erysiphe carpinicola* (perfect state) in Poland. *New Disease Report*. **14**: 21.
- Walczanska, A. (2008).** New records of Erysiphales and Uredinales from Poland. *Acta. Mycologica*. **43** (1): 71–75.
- Weltzein, H.C. (1963).** *Erysiphe bitae* (Vanha) *Comb. Nov.* The powdery mildew of beets. *Phytopath. Z.* **47**: 123-128.
- Williams, J.D. and R.H. Groves. (1980).** The influence of temperature and photoperiod on growth and development of *Parthenium hysterophorus* L. *Weed Res.*, **20**: 47-52.
- Yaduraja, T.N. (2005).** Campaign Launched for Biological Control of a Dangerous Weed. *The Hindu*, September, 4.
- Yarwood, C.E. (1952).** Apricot powdery mildew from rose and peach. *Calif. Dept. Agric. Monthly Bull.* **41**:19-25.
- Yarwood, C.E. (1957).** Powdery mildew. *Bot. Rev.* **23**: 235-300.
- Yarwood, C.E. (1973).** Pyrenomycetes: Erycophales. In “The Fungi” eds. G.C. Anisworth, F.K. Sparrow and A.S. Susman Academic Press New York. pp. 71-86.
- Yarwood, C.E. and Gardner, M.W. (1974).** A list of powdery mildews of California. *Calif. Agric. Exp. Sta. Ext. Serv.* **31** pp.

Reference

- Yarwood, C. E. (1978).** History and taxonomy of powdery mildews. In: Spencer, D. M. (ed.), *The Powdery Mildews*. Academic Press, London, UK. pp. 1-38.
- Zaracovitis, C. (1965).** Attempts to identify of powdery mildew fungi by conidial characters. *Trans. Brit. Myco. Soc.* **48**: 533-558.