

ACKNOWLEDGEMENTS

I take this opportunity to record my deep sense of gratitude to Dr. Mohd. Shafi for his constant guidance and encouragement, and also for his painstaking attempts to teach me Physics. It is a great pleasure to express my gratefulness to Prof. Rais Ahmad, Head Deptt. of Physics, Aligarh Muslim University, Aligarh for his personal interest in the work and also for providing to me all sorts of facilities in the Department. I am highly indebted to my teachers, Prof. M.Z.R. Khan, Dr. J.H. Naqvi and Dr. Israr Ahmad who have been a constant source of inspiration.

I am thankful to all members of the Nuclear Emulsion Group and in particular to Mr. A.R. Ansari, Dr. K.B. Lal and Dr. A.H. Naqvi for the very fruitful and pleasant collaboration. The advice and critical suggestions of Messrs Zubair Ahsan, Zaheer Ahmad Khan, M. Zafar and Rashid Hasan were always available and have been of great help. The completion of this work owes a great deal to their strenuous efforts in hunting of references and carrying out of computations.

I am grateful to Dr. Jaliluddin, Dr. Khaliq Raza and Mr. M. Sajid for their active cooperation and various other helps. The help of Messrs M. Laiq and A.K. Agrawal is gratefully acknowledged. I express my thanks to Messrs S.A. Jafri, S.A. Ahmad and H. Rahman for careful scanning. My thanks are also due to Mr. S. Munawar Ali for typing this thesis.

Finally, the financial assistance from the Council of Scientific and Industrial Research, Government of India, is thankfully acknowledged.


(Mohammad Irfan)