

### List of Figures

<b>Figure1.1</b>	Crystal structure of rutile (left) and anatase (right); Ti: grey and O: red	21
<b>Figure1.2</b>	Crystal Structure of ZrO <sub>2</sub> (baddeleyite); Zr: grey and O: red	21
<b>Figure1.3</b>	Phase transformation of ZrO <sub>2</sub> at different temperatures	24
<b>Figure1.4</b>	Three phases of hafnia and zirconia; a: cubic, b: tetragonal, c: monoclinic	25
<b>Figure1.5</b>	Types of displacive transformation	26
<b>Figure1.6</b>	Pictorial representation of the two phases in zirconia	27
<b>Figure1.7</b>	Hysteresis in martensitic transformation	28
<b>Figure1.8</b>	Mechanism of Photo-oxidation by charge-carriers for TiO <sub>2</sub>	30
<b>Figure1.9</b>	Formation of YSZ in a step-wise manner	35
<b>Figure1.10</b>	System of charges arranged in a particular manner	47
<b>Figure1.11</b>	Electric Quadrupole Interaction and splitting of energy levels	51
<b>Figure1.12</b>	Decay of parent (P) to daughter (D) nuclei and subsequent emission of cascade $\gamma$ -rays from the daughter atom	54
<b>Figure1.13</b>	Splitting of Intermediate level with $I = 5/2$ in presence of external field	62
<b>Figure1.14</b>	Effect of $\eta$ on perturbation function for $I=5/2$ state	75
<b>Figure1.15</b>	Decay schemes of <sup>181</sup> Hf/ <sup>181</sup> Ta (left) and <sup>111</sup> In/ <sup>111</sup> Cd probes	90
<b>Figure2.1</b>	Fast-Slow coincidence circuit diagram for TDPAC measurement	110
<b>Figure2.2</b>	The energy spectra obtained with LaBr <sub>3</sub> (Ce) detector <sup>133</sup> Ba, <sup>152</sup> Eu and <sup>106</sup> Ru	115
<b>Figure2.3</b>	The energy response curve for LaBr <sub>3</sub> (Ce) detector up to 1.4 MeV	116
<b>Figure2.4</b>	Prompt for 511-511 cascade of <sup>22</sup> Na source	117
<b>Figure2.5</b>	The variation of time resolution of the detector at different bias voltages	118
<b>Figure3.1</b>	TDPAC spectra with cosine transform for anatase and rutile with <sup>181</sup> Ta probe	128
<b>Figure3.2</b>	Representative TDPAC spectra with cosine transform for HfO <sub>2</sub> and ZrO <sub>2</sub>	129

<b>Figure3.3</b>	TDPAC spectra with cosine transform for rutile TiO <sub>2</sub> with <sup>111</sup> In probe	130
<b>Figure3.4</b>	Representative TDPAC spectrum for HfO <sub>2</sub> and ZrO <sub>2</sub> with <sup>111</sup> In probe	131
<b>Figure3.5</b>	TDPAC spectrum for 5% Mn-doped anatase TiO <sub>2</sub>	134
<b>Figure3.6</b>	TDPAC spectrum for 5% Mn-doped rutile TiO <sub>2</sub>	135
<b>Figure3.7</b>	TDPAC spectrum for 1% Zr-doped rutile TiO <sub>2</sub>	135
<b>Figure3.8</b>	TDPAC spectra for 5% (left) and 10% (right) Zr -doped rutile TiO <sub>2</sub>	136
<b>Figure3.9</b>	TDPAC spectrum with cosine transform for $\gamma$ -irradiated rutile TiO <sub>2</sub>	141
<b>Figure3.10</b>	XRD spectra of the HfO <sub>2</sub> fiber annealed at different temperatures	145
<b>Figure3.11</b>	SEM picture of HfO <sub>2</sub> fiber annealed at (a) 1173K and (b) 1673K	145
<b>Figure3.12</b>	TDPAC spectra for the HfO <sub>2</sub> fiber annealed at different temperatures	146
<b>Figure3.13</b>	HfO <sub>2</sub> crystal doped with Ta	148
<b>Figure3.14</b>	Lowest panel: Plot of Total Density of states (DOS) vs energy for Ta doped HfO <sub>2</sub> . Middle panel: Plot of Atom (Ta) projected DOS vs energy, Upper panel: Plot of projected DOS vs energy of probe Ta in HfO <sub>2</sub> .	150
<b>Figure3.15</b>	Lowest panel: Plot of Density of states vs energy for pure HfO <sub>2</sub> . Upper panel: Plot of atom projected density of states vs energy of nearest neighbour of probe Ta.	151
<b>Figure4.1</b>	Size distribution of the TiO <sub>2</sub> nanoparticles	156
<b>Figure4.2</b>	XRD spectra of TiO <sub>2</sub> annealed at different temperatures	157
<b>Figure4.3</b>	TDPAC spectra of nano TiO <sub>2</sub> samples annealed at different temperatures	159
<b>Figure4.4</b>	UV-Vis absorption spectra for Ag and Ag@TiO <sub>2</sub> nanoparticles.	166
<b>Figure4.5</b>	TEM picture of Ag@TiO <sub>2</sub> core-shell nanoparticles.	166
<b>Figure4.6</b>	XRD patterns at different annealing temperature for Ag@TiO <sub>2</sub> nanoparticles	167

<b>Figure4.7</b>	XRD patterns at different annealing temperature for pure TiO <sub>2</sub> nanoparticles	167
<b>Figure4.8</b>	Nature of grain growth with annealing temperature	169
<b>Figure4.9</b>	TDPAC spectra of Ag@TiO <sub>2</sub> at different annealing temperatures	169
<b>Figure4.10</b>	TDPAC spectra of pure TiO <sub>2</sub> and Ag@TiO <sub>2</sub> nanoparticles at 1073K	170
<b>Figure4.11</b>	Schematic of particle growth in pure TiO <sub>2</sub> and Ag@TiO <sub>2</sub> nanoparticles	171
<b>Figure4.12</b>	Schematic of Thin film preparation by ion beam sputter deposition	177
<b>Figure4.13</b>	RBS spectrum for unannealed Hf thin film on Si (111) substrate	178
<b>Figure4.14</b>	TDPAC spectra of thin film annealed under 10mbar O <sub>2</sub> pressure	179
<b>Figure4.15</b>	TDPAC spectra of the thin film annealed at 1273K under 10mbar O <sub>2</sub> pressure for different durations	180
<b>Figure4.16</b>	TDPAC spectrum of the thin film annealed directly at 1273K under 10mbar O <sub>2</sub> pressure for different durations	180
<b>Figure4.17</b>	Evolution of monoclinic phase under different annealing conditions	181
<b>Figure4.18</b>	AFM pictures of thin film as prepared (left) and annealed at 1273K (right)	181