

## LIST OF FIGURES

- Figure I.1. Summary of environmental influences on the establishment of naturally regenerated forests.
- Figure I.2. Major causes and consequences of patterns of natural regeneration of forest trees.
- Figure II.1. Rainfall, relative humidity and temperature data for the study area during January to December (mean values for the years 1997,1998,1999 & 2000).
- Figure III.1. Density of tree species in various girth classes in four forest stands.
- Figure III.2. Species richness of tree species among various girth classes in four forest stands experiencing different degrees of disturbance.
- Figure III.3. Population structure of some important trees growing with *Elaeocarpus ganitrus*.
- Figure IV.1. Density of tree, sapling and seedling populations of Rudraksh in four forest stands experiencing different degrees of disturbance.
- Figure IV.2. Population structure of Rudraksh in four forest stands.
- Figure IV.3. Number of cut stumps and percentage of sprouting stumps in different natural and planted forest stands.
- Figure V.1. Dispersal curves of Rudraksh in different forest stands.
- Figure V.2. Percentage of disappeared and dormant/rotten nuts at different distance from parent tree of Rudraksh in different forest stands.
- Figure V.3. Disappearance of fruits of Rudraksh as influenced by their ripeness and pulp in the undisturbed (UND), mildly disturbed (MID), moderately disturbed (MOD) and highly disturbed (HID) forest stands.
- Figure V.4. Schematic diagram of general trend of different events in Rudraksh.
- Figure VI.1. Frequency distribution of 10,000 Rudraksh nuts having different number of locules or 'faces'.
- Figure VI.2. Frequency distribution of seeds per nut (n= 405).
- Figure VII.1. Seedling growth in terms of shoot elongation, collar diameter, leaf number and leaf area under nursery conditions.
- Figure VII.2. Growth rate of seedlings in three forest stands; open canopy (stand I), sparse canopy (stand II) and dense canopy (stand III).
- Figure VII.3. Growth rate of seedlings in terms of shoot length, collar diameter, leaf number and leaf area in stand I, stand II and stand III after one year period of transplantation.