

## LIST OF TABLES

		<b>Page no.</b>
Table 2.1	Summary of the gene effects controlling different agronomic characters including yield and fiber attributes in cotton	12
Table 2.2	Summary of the studies on correlation analysis of seed cotton yield with other important traits in cotton	18
Table 2.3	Summary of the recent studies involving the application of molecular markers in diversity analysis, fingerprinting, seed and genetic purity analysis in cotton	23
Table 2.4	Summary of the studies on the use of molecular markers in linkage studies in cotton	26
Table 2.5	Summary of the significant studies on QTL analysis in cotton	32
Table 3.1	Details of cotton genotypes used in biometrical studies	34
Table 3.2	ANOVA of diallel cross analysis	38
Table 3.3	ANOVA for combining ability analysis	38
Table 3.4	Details of the cotton genotypes used for molecular diversity analysis	46
Table 4.1	Analysis of variance of $6 \times 6$ full diallel for yield and quality contributing traits in cotton	56
Table 4.2	Genetic components of variance for yield and quality contributing traits in cotton	58
Table 4.3	GCA effects of the parental lines for yield and quality contributing traits in cotton	60
Table 4.4	SCA effects for yield and quality contributing traits in direct and reciprocal $F_1$ hybrids in cotton	62
Table 4.5	Mean values for yield and quality contributing traits of 6 parental genotypes of a diallel cross in cotton	66
Table 4.6	Mean values of direct $F_1$ hybrids ( $F_{1s}$ ) and reciprocal $F_1$ hybrids ( $RF_{1s}$ ) and percent heterosis over better parent and best parent for plant height and monopodial branches in $6 \times 6$ diallel crosses in cotton	68
Table 4.7	Mean values of direct $F_1$ hybrids ( $F_{1s}$ ) and reciprocal $F_1$ hybrids ( $RF_{1s}$ ) and percent heterosis over better parent and best parent for sympodial branches and seed cotton yield in $6 \times 6$ diallel crosses in cotton	70
Table 4.8	Mean values of direct $F_1$ hybrids ( $F_{1s}$ ) and reciprocal $F_1$ hybrids ( $RF_{1s}$ ) and percent heterosis over better parent and best parent for lint index and ginning out turn in $6 \times 6$ diallel crosses in cotton	71
Table 4.9	Mean values of direct $F_1$ hybrids ( $F_{1s}$ ) and reciprocal $F_1$ hybrids ( $RF_{1s}$ ) and percent heterosis over better parent and best parent for boll number and boll weight in $6 \times 6$ diallel crosses in cotton	73

		<b>Page no.</b>
Table 4.10	Mean values of direct F <sub>1</sub> hybrids (F <sub>1</sub> s) and reciprocal F <sub>1</sub> hybrids (RF <sub>1</sub> s) and percent heterosis over better parent and best parent for seed index and 2.5% span length in 6 × 6 diallel crosses in cotton	75
Table 4.11	Mean values of direct F <sub>1</sub> hybrids (F <sub>1</sub> s) and reciprocal F <sub>1</sub> hybrids (RF <sub>1</sub> s) and percent heterosis over better parent and best parent for uniformity ratio and fiber strength in 6 × 6 diallel crosses in cotton	76
Table 4.12	Mean values of direct F <sub>1</sub> hybrids (F <sub>1</sub> s) and reciprocal F <sub>1</sub> hybrids (RF <sub>1</sub> s) and percent heterosis over better parent and best parent for micronaire value and fiber elongation in 6 × 6 diallel crosses in cotton	77
Table 4.13	Mean performance for yield and quality contributing traits of six generations involving parents P56-4 (P <sub>1</sub> ) and RS2013 (P <sub>2</sub> )	79
Table 4.14	Scaling test for generation means for the cross P56-4 (P <sub>1</sub> ) × RS2013 (P <sub>2</sub> ) for yield and quality contributing traits	83
Table 4.15	Genetic parameters for yield and quality contributing traits following joint scaling test	84
Table 4.16	Phenotypic correlation coefficient values among yield and fibre quality contributing traits in cotton	87
Table 4.17	Direct and indirect effects of yield and fiber quality traits on seed cotton yield	88
Table 4.18	Direct and indirect effects of yield and fiber quality traits on fiber strength in cotton	89
Table 4.19	Details of number of alleles, polymorphic information content (PIC) and heterozygosity percentage of 38 SSRs used for diversity analysis	91
Table 4.20	Estimates of Intra- and Inter-cluster distances among three clusters involving six cotton genotypes obtained due to Mahalanobis D <sup>2</sup> analysis	94
Table 4.21	Extent of polymorphism revealed between the parental genotypes P56-4 and RS2013 using 32 AFLP primer combinations	95
Table 4.22	Marker-trait associations detected through simple linear regression analysis in different generations	97
Table 4.23	Marker-trait associations detected in different generations through multiple regression analysis	99
Table 5.1	Summary of the best parents, best crosses, best general combiners and best specific combinations (not more than two in each case) in a 6×6 diallel cross for 14 traits in cotton	116
Table 5.2	Summary of the marker–trait associations detected in different generations through simple and multiple regression analyses	118