

- Achari R, Mayersohn M, Conrad KA (1983) HPLC analysis of creatinine in human plasma and urine. *J. Chromatogr. Sci.* **21**: 278-281.
- Akamatsu S, Kanai Y (1951) Bacterial decomposition of creatinine I. Creatinomutase. *Enzymologia.* **15**: 122-125.
- Ambrose RT, Ketchum DF, Smith JW (1983) Creatinine determined by "High-Performance" liquid chromatography. *Clin. Chem.* **29**: 256-259.
- Antunes AM, Ferreira MMC, Melgo MS, Volpe PLO (1999) Potassium transport through liquid membranes using spectral and chemometric methods. *J. Mol. Strut.* **481**: 563-567.
- Arya SK, Solanki PR, Singh RP, Pandey MK, Datta M, Malhotra BD (2006) Application of octadecanethiol self-assembled monolayer to cholesterol biosensor based on surface plasmon resonance technique. *Talanta* **69**: 918-926.
- Atkinson SA, Halton JM, Bradley C, Wu B, Barr RD (1998) Bone and mineral abnormalities in childhood acute lymphoblastic leukemia: influence of disease, drugs and nutrition. *Int. J. Cancer. Suppl.* **11**: 35-39.
- Bagnoud MA, Reymand JP (1993) Interference of metamizol (Dipyrone) on the determination of creatinine with the Kodak dry chemistry slide. Comparison with the enzymatic method from Boehringer. *Eur. J. Clin. Chem. Clin. Bio. Chem.* **31**: 753-757.
- Barr DB, Wilder LC, Caudill SP, Gonzalez AJ, Needham LL, Pirkle JL (2005) Urinary creatinine concentrations in the U.S. population: Implications for urinary biologic monitoring measurements. *Environ. Health Perspect.* **113**: 192-200.
- Battilotti M, Colapicchioni C, Giannini I, Porcelli F (1989) Characterization of biosensors based on membranes containing a conducting polymer. *Anal. Chim. Acta.* **221**: 157-161.
- Bauer J, Brooks C, Burch R (1982) Clinical appraisal of creatinine clearance as a measurement of glomerular filtration rate. *Am. J. Kidney Dis.* **2**: 337-346.
- Benedict SB, Behre JA (1936) Some applications of a new color reaction for creatinine. *J. Biol. Chem.* **114**: 515-532.
- Benkert A, Scheller F, Schossler W, Hentschel C, Micheel B, Behrsing O, Scharte G, Stocklein W, Warsinke A (2000a) Development of a creatinine ELISA and an amperometric antibody-based creatine sensor with a detection limit in the nanomolar range. *Anal. Chem.* **72**: 916-921.

- Benkert A, Scheller FW, Schoessler W, Micheel B, Warsinke A (2000b) Size exclusion redox-labeled immunoassay (SERI): A new format for homogeneous amperometric creatinine determination. *Electroanalysis* **12**: 1318-1321.
- Bergam A, Ohman G (1980) Effect of detergent on kinetic Jaffe-method assay of creatinine. *Clin. Chem.* **26**: 1729-1732.
- Berger GM, Cowlin J, Turner TJ (1976) Amylase:creatinine clearance ratio and urinary excretion of lysozyme in acute pancreatitis and acute duodenal perforation. *S. Air. Med. J.* **50**: 1559-1561.
- Besbas N, Karpman D, Landau D, Loirat C, Proesmans W, Remuzzi G, Rizzoni G, Taylor CM, Van de Kar N, Zimmerhackl LB (2006) A classification of hemolytic uremic syndrome and thrombotic thrombocytopenic purpura and related disorders. *Kidney Int.* **70**: 423-431.
- Beyer C, Hoenderdos A, Mairuhu WM, Stadius-Van Eps LW, Van-Den-Ende A (1984) Evaluative and comparative study of an enzymatic method using creatine kinase for the determination of urinary creatine. *Clin. Chim. Acta.* **136**: 263-270.
- Bjorkhem I, Bergman A, Falk O, Kallner A, Lantto O, Svensson L, Akerlof E, Blomstrand R (1981) Accuracy of some routine method used in clinical chemistry as judged by isotope dilution-mass spectrometry. *Clin. Chem.* **27**: 733-735.
- Bjorkhem I, Blomstrand R, Ohman G (1977) Mass fragmentography of creatinine proposed as a reference method. *Clin. Chem.* **23**: 2114-2121.
- Bjorkhem I, Blomstrand R, Ohman G (1979) Accuracy of a fully enzymatic method for creatinine as judge by isotope dilution-mass fragmentography. *Clin. Chem.* **25**: 1666-1667.
- Bjornsaon TD, Cocchetto DM, McGowan FX, Vergheae CP, Sedor F (1983) Nomogram for estimating creatinine clearance. *Clin. Pharmacokinet.* **8**: 365-369.
- Blijenberg B, Brower H, Kuller T, Leeneman R, Van-Lecuwen C (1994) Improvement in creatinine methodology: A critical asseement. *Eur. J. Clin. Chem. Clin. Biochem.* **32**: 529-537.
- Bloch K, Schoenheimer R (1939) Studies in protein metabolism. XI. The metabolic relation of creatine and creatinine studied with isotopic nitrogen. *J. Biol. Chem.* **131**: 111-119.

- Bohn H, Haun F (1933) Untersuchungen zum Mechanismus des blassen Hochdrucks. Blutdrucksteigernde Stoffe im Harn, insbesondere beim blassen und beim roten Hochdruck. *Z. klin. Med.* **125**: 558-584.
- Bonvicini P, Ceriotti G, de-Besi T (1982) Kinetic enzymatic determination of creatinine by a rapid semiautomated procedure. *J. Clin. Chem. Clin. Biochem.* **20**: 185-189.
- Borsook H, Dubnoff JW (1947) The hydrolysis of phosphocreatine and the origin of urinary creatinine. *J. Biol. Chem.* **168**: 493-510.
- Bourdillon C (2006) Enzymatic electrolysis, spatial and kinetic coupling between electrochemical reaction and enzymatic catalysis. *Enz. Engg.* **672**: 195-212.
- Bowers LD, Wong ET (1980) Kinetic sum creatinine assay II: A critical evaluation and review. *Clin. Chem.* **26**: 555-561.
- Brahim S, Narinesingh D, Guseppi-Elie A (2002) Polypyrrole-hydrogel composites for the construction of clinically important biosensors. *Biosens. Bioelectron.* **17**: 53-59.
- Bristow TT, Stokes PP, O'Connor GG (2005) Quantitative Fourier transform ion cyclotron resonance mass spectrometry--the determination of creatinine by isotope dilution mass spectrometry. *Rapid Commun. Mass Spectrom.* **19**: 375-380.
- Brogan KL, Walt DR (2005) Optical fiber-based sensors: application to chemical biology. *Curr. Opin. Chem. Biol.* **9**: 494-500.
- Brown GB (1966) Séparation chromatographique en couche mince de la créatine, la créatinine et de leurs dérivés N-phosphorylés. *J. Chromatogr.* **25**: 161-162.
- Campanella L, Mazzei F, Sammartino MP, Tomassetti M (1990b) New enzyme sensors for urea and creatinine analysis. *Bioelectrochem. Bioenerg.* **23**: 195-202.
- Campanella L, Sammartino MP, Tomassetti M (1990a) Suitable potentiometric enzyme sensors for urea and creatinine. *Analyst* **115**: 827-830.
- Chang MC, Chang CC, Chang JC (1992) Cloning of a creatinase gene from *Pseudomonas putida* in *Escherichia coli* by using an indicator plate. *Appl. Environ. Microbiol.* **58**: 3437-3440.
- Chasson AL, Grady HJ, Stanley MA (1961) Determination of creatinine by means of automatic analysis. *Am. J. Clin. Pathol.* **35**: 83-88.
- Chaubey A, Malhotra BD (2002) Mediated biosensors. *Biosens. Bioelectron.* **17**: 441-454.
- Chen CH, Lin MS (2012) A novel structural specific creatinine sensing scheme for the determination of the urine creatinine. *Biosens. Bioelectron.* **31**: 90-94.

- Chen JC, Kumar AS, Chung HH, Chien SH, Kuo MC, Zen JM (2006) An enzymeless electrochemical sensor for the selective determination of creatinine in human urine. *Sens. Actuat. B* **115**: 473-480.
- Chen SA, Hwang GWJ (1994) Synthesis of Water-Soluble Self-Acid-Doped Polyaniline. *Am. Chem. Soc.* **116**: 7939-7940.
- Chinn KSK (1967) Prediction of muscle and remaining tissue protein in man. *J. Appl. Physiol.* **23**: 713-715.
- Chiou W, Pu F, Prueksaritanont T (1983) Creatinine: XIII. Micro high-performance liquid chromatography assay of creatinine in biological using a fixed or variable wavelength UV-detector. *J. Chromatogr.* **277**: 436-438.
- Chiou WL, Gadalla MAF, Peng GW (1978) Simple, rapid, and micro high-pressure liquid chromatographic determination of endogenous "true" creatinine in plasma, serum, and urine. *J. Pharm. Sci.* **67**: 182-187.
- Choi SH, Lee SD, Shin JH, Ha J, Nam H, Cha GS (2002) Amperometric biosensors employing an insoluble oxidant as an interference-removing agent. *Anal. Chim. Acta.* **461**: 251-260.
- Chou JC, Sun TP, Hsiung SK (2009) All solid-state potentiometric biosensors for creatinine determination based on pH and ammonium electrodes. *Sensor* **9**: 665-672.
- Christopher B, Jacobs M, Jennifer-Peairs B, Venton J (2010) Review: carbonnanotube based electrochemical sensors for biomolecules. *Anal. Chim. Acta.* **662**: 105-127.
- Clark EA, Fanguy JC, Henry CS (2001) High-throughput multi-analyte screening for renal disease using capillary electrophoresis. *J. Pharm. Biomed. Anal.* **25**: 795-801.
- Clark LC, Lyons C (1962) "Biosensors - A Practical Approach." *J. Amn. N. Y. Acad. Sci.* **102**: 29-45.
- Cockroft G, Gault M (1976) Prediction of creatinine clearance from serum creatinine. *Nephron* **16**: 31-41.
- Coll M, Knof SH, Ohga Y, Messerschmidt A, Huber R, Moellering H, Russmann L, Schumacher G (1990) Enzymatic mechanism of creatine amidinohydrolase as deduced from crystal structures. *J. Mol. Biol.* **214**: 597-610.
- Collison ME, Meyerhoff ME (1987) Continuous-flow enzymatic determination of creatinine with improved on-line removal of endogenous ammonia. *Anal. Chim. Acta.* **200**: 61-72.
- Conn Jr RB (1960) Fluorometric determination of creatine. *Clin. Chem.* **6**: 537-548.

- Cook JGH (1971) Creatinine assay in the presence of protein. *Clin. Chem. Acta.* **32**: 485-486.
- Cook JGH (1975) Factors influencing the assay of creatinine. *Ann. Clin. Biochem.* **12**: 219-232.
- Cooper JM, Biggs HG (1961) An evaluation of four methods of measuring urinary creatinine. *Clin. Chem.* **7**: 665-673.
- D’Orazio P (2003) Biosensors in clinical chemistry. *Clin. Chim. Acta.* **334**: 41-69.
- D’Souza SF (2001) Microbial biosensors. *Biosens Bioelectron.* **16**: 337-353.
- Dai X, Fang X, Shao M, Li M, Huang Z, Li H, Jiang Y, Song D, He Y (2011) Certified reference materials (GBW09170 and 09171) of creatinine in human serum. *J. Chromatogr. B: Analyt. Technol. Biomed. Life Sci.* **879**: 429-435.
- Dash AK, Sawhney A (2002) A simple LC method with UV detection for the analysis of creatine and creatinine and its application to several creatine formulations. *J. Pharm. Biomed. Anal.* **29**: 939-945.
- Delaney TP, Mirsky VM, Wolfbeis OS (2002) Capacitive creatinine sensor based on a photografted molecularly imprinted polymer. *Electroanalysis* **3**: 221-224.
- Delanghe J, De Slypere JP, De Buyzere M, Robbrecht J, Wieme R, Vermeulen (1989) A Normal reference values for creatine, creatinine and carnitine are lower in vegetarians (PDF). *Clin. Chem.* **35**: 1802-1803.
- Devi R, Thakur M, Pundir CS (2011a) Construction and application of an amperometric xanthine biosensor based on zinc oxide nanoparticles–polypyrrole composite film. *Biosens. Bioelectron.* **26**: 3420-3426.
- Devi R, Yadav S, Pundir CS (2011b) Electrochemical detection of xanthine by xanthine oxidase immobilized on carboxylated multiwalled carbon nanotubes/polyaniline composite film. *Biochem. Engineering J.* **58-59**: 148-153.
- Dodder NG, Tai SS, Sniegoski LT, Zhang NF, Welch MJ (2007) Certification of creatinine in a human serum reference material by GC-MS and LC-MS. *Clin. Chem.* **53**: 1694-1699.
- Du W, Zhao F, Zeng B (2009) Novel multiwalled carbon nanotubes–polyaniline composite film coated platinum wire for headspace solid-phase microextraction and gas chromatographic determination of phenolic compounds. *J. Chromato. A.* **1216**: 3751-3757.

- Edward KDG, Whyte HM (1958) The measurement of creatinine in plasma and urine. *Aust. J. Exp. Biol. Med.* **36**: 383-394.
- Elmosallamy MAF (2006) New potentiometric sensors for creatinine. *Anal. Chim. Acta.* **564**: 253-257.
- Ennor AH, Morrison JF (1958) Biochemistry of the phosphagens and related guanidines. *Physiol. Rev.* **38**: 631-674.
- Erlenkotter A, Fobker M, Chemnitz GC (2002) Biosensors and flow-through system for the determination of creatinine in hemodialysate. *Anal. Bioanal. Chem.* **372**: 284-292.
- Facila L, Nunez J, Bodi V, Sanchis J, Bertomeu-Gonzalez V, Consuegra L, Pellicer M, Ferrero A, Sanjuan R, Llacer A (2006) Prognostic value of serum creatinine in non ST elevation acute coronary syndrome. *Rev. Esp. Cardiol.* **59**: 209-216.
- Feng JJ, Zhao G, Xu JJ, Chen HY (2005) Direct electrochemistry and electrocatalysis of heme proteins immobilized on gold nanoparticles stabilized by chitosan. *Anal. Biochem.* **342**: 280-286.
- Fink JC, Burdick RA, Kurth SJ, Blahut SA, Armistead NC, Turner MS, Shickle LM, Light PD (1999) Significance of serum creatinine values in new end-stage renal disease patients. *Am. J. Kidney Disease.* **34**: 694-701.
- Fischl J, Segal S, Yulzari Y (1964) Separation of creatine and creatinine by "small-volume" electrophoresis. *Clin. Chim. Acta.* **10**: 73-75.
- Fitch CD, Shield RP (1966) Creatine metabolism in skeletal muscle. I. Creatine movement across muscle membranes. *J. Biol. Chem.* **241**: 3611-3614.
- Folin O (1905) Laws governing the chemical composition of urine. *Am. J. Physiol.* **13**: 66-115.
- Folin O (1914) On the determination of creatinine and creatine in blood, milk and tissues. *J. Biol. Chem.* **17**: 475-481.
- Fossati P, Ponti M, Passoni G, Tarengi G, d'Eril G, Prencip L (1994) A step forward in enzymatic measured of creatinine. *Clin. Chem.* **40**: 130-137.
- Fossati P, Prencipe L, Bertl G (1983) Enzymic creatinine assay: A new colorimetric method based on hydrogen peroxide measurement. *Clin. Chem.* **29**: 1494-1496.
- Fridolin I, Jerotskaja J, Lauri K, Uhlin F, Luman M (2010) A new optical method for measuring creatinine concentration during dialysis. *IFMBE Proc.* **29**: 379-382.
- Friedman RB, Anderson RE, Entine SM, Hirshberg SB (1980) Effect of diseases on clinical laboratory tests. *Clin. Chern.* **26**: ID-476D.

- Fujii T, Kawabe S, Horike T, Taguchi T, Ogata M (1999) Simultaneous determination of the urinary metabolites of toluene, xylene and styrene using high-performance capillary electrophoresis: Comparison with high-performance liquid chromatography. *J. Chromatogr. B* **730**: 41-47.
- Fung YS, Wong CCW (2002) Determination of carbon monoxide in ambient air using piezoelectric crystal sorption detection. *Anal. Chim. Acta.* **456**: 227-239.
- Gao B, Li Y, Zhang Z (2010) Preparation and recognition performance of creatinine-imprinted material prepared with novel surface-imprinting technique. *J. Chromatogr. B.* **878**: 2077-2086.
- Garcia CD, Henry CS (2004) Direct detection of renal function markers using microchip CE with pulsed electrochemical detection. *Analyst* **129**: 579-584.
- Gatti R, Lazzarotto V, De Palo CB, Cappellin E, Spinella P, De Palo EF (1999) A rapid urine creatinine assay by capillary zone electrophoresis. *Electrophoresis* **20**: 2917-2921.
- Geigy (1971) Scientific tables, 7th Ed. Basle, Ciba-Geigy Ltd. 572, 665.
- Goren M, Osborne S, Wright R (1986) A peroxidase coupled kinetic enzymatic procedure evaluated for measurement of serum and urinary creatinine. *Clin. Chem.* **32**: 548-551.
- Gospodinova N, Terlemezyan L (1998) Conducting polymers prepared by oxidative polymerization: Polyaniline. *Prog. Polym. Sci.* **23**: 1443-1484.
- Grabowska I, Chudy M, Dybko A, Brzozka Z (2005) Determination of creatinine in clinical samples based on flow-through microsystem. *Anal. Chim. Acta.* **540**: 181-185.
- Grabowska I, Sajnoga M, Juchniewicz M, Chudy M, Dybko A, Brzozka Z (2007) Microfluidic system with electrochemical and optical detection. *Microelectron. Engineer.* **84**: 1741-1743.
- Granot E, Basnar B, Cheglakov Z, Katz E, Willner I (2006) Enhanced bioelectrocatalysis using single-walled carbon nanotubes (SWCNTs)/polyaniline hybrid systems in thin-film and microrod structures associated with electrodes. *Electroanalysis* **18**: 26-34.
- Greenwald I, Gross J (1924) The chemistry of Jaffe's reaction for creatinine. A red tautomer of creatinine pictrate. *J. Biol. Chem.* **59**: 601-612.
- Greenwald I, McGuire G (1918) The estimation of creatinine and of creatine in the blood. *J. Biol. Chem.* **34**: 103-118.

- Guder WG, Hoffmann GE, Hubbuch A, Poppe WA, Siedel J, Price CP (1986) Multicentre evaluation of an enzymatic method for creatinine determination using a sensitive colour reagent. *J. Clin. Chem. Clin. Biochem.* **24**: 889-902.
- Guilbault GG, Chen SP, Kuan SS (1980) A creatinine specific enzyme electrode. *Anal. Lett.* **13**: 1607-1624.
- Guilbault GG, Coulet PR (1983) Creatinine-selective enzyme electrodes. *Anal. Chim. Acta.* **152**: 223-228.
- Guo MD, Guo HX (2005) Voltammetric behaviour study of creatinine at phosphomolybdic-polypyrrole film modified electrode. *J. Electroanal. Chem.* **585**: 28-34.
- Guterres MV, Volpe POL, Ferreira MMC (2004) Multiway calibration for creatinine determination in human serum using the Jaffe' reaction. *Appl. Spectro.* **58**: 54-60.
- Gutierrez M, Alegret S, del Valle M (2008) Bioelectronic tongue for the simultaneous determination of urea, creatinine and alkaline ions in clinical samples. *Biosens. Bioelectron.* **23**: 795-802.
- Guzman NA, Berck CM, Hernandez L, Advis JP (1990) Capillary electrophoresis as a diagnostic tool: Determination of biological constituents present in urine of normal and pathological individuals. *J. Liq. Chromatogr.* **13**: 3833-3848.
- Haeckel R (1981) Assay of creatinine in serum, with use of Fuller's Earth to remove interferences. *Clin. Chem.* **27**: 179-183.
- Hahn A, Meyer G (1928) On the mutual transformation of creatine and creatinine. *Z. Biol.* **78**: 111-115.
- Harita N, Hayashi T, Sato KK, Nakamura Y, Yoneda T, Endo G, Kambe H (2009) Lower serum creatinine is a new risk factor of type 2 diabetes: the Kansai healthcare study. *Diabetes Care* **32**: 424-426.
- Harlan R, Clarke W, Di-Bussolo JM, Kozak M, Straseski J, Li-Meany D (2010) An automated turbulent flow liquid chromatography–isotope dilution mass spectrometry (LC–IDMS) method for quantitation of serum creatinine. *Clin. Chim. Acta.* **411**: 1728-1734.
- Hartmann L, Doree G, Ambert JP (1983) Gas–liquid-chromatography reference method for the assay of urinary creatinine. *J. Chromatogr.* **273**: 235-243.

- Hassan SSM, Eman ME, Ayman HKM (2005) Novel biomedical sensors for flow injection potentiometric determination of creatinine in human serum. *Electroanalysis* **17**: 2246-2253.
- Hayashi S, Suzuki M, Nakamura S (1983) Steady-state kinetics and spectral properties of *Corynebacterium* sarcosine oxidase. *Biochim. Biophys. Acta.* **742**: 630-636.
- He ZK, Fuhrmann B, Spohn U (2000) Calibrationless determination of creatinine and ammonia by coulometric flow titration. *Anal. Biochem.* **283**: 166-174.
- Hetu PO, Gingras ME, Vinet B (2010) Development and validation of a rapid liquid chromatography isotope dilution tandem mass spectrometry (LC-IDMS/MS) method for serum creatinine. *Clin. Biochem.* **43**: 1158-1162.
- Hicks J, Josefoohn M, Levis S (1979) Evaluation of the beckman creatinine analyzer. *Clin. Chem.* **25**: 1005-1008.
- Hirst J, Duff JLC, Jameson GNL, Kemper MA, Burgess BK, Armstrong FA (1998) Kinetics and mechanism of redox-coupled, long-range proton transfer in an iron-sulfur protein. Investigation by fast-scan protein-film voltammetry. *J. Am. Chem. Soc.* **120**: 7085-7094.
- Hjelmesath J, Roislien J, Nordstrand N, Hofso D, Hager H, Hartmann A (2010) Low serum creatinine is associated with type 2 diabetes in morbidly obese women and men: A cross-sectional study. *BMC Endocrine Disorders* **10**: 1-6.
- Ho WO, Krause S, McNeil CJ, Pritchard JA, Armstrong RD, Athey D, Rawson K (1999) Electrochemical sensor for measurement of urea and creatinine in serum based on ac impedance measurement of enzyme-catalyzed polymer transformation. *Anal. Chem.* **71**: 1940-1946.
- Hoeffken HW, Knof SH, Bartlett PA, Huber R (1988) Crystal structure determination, refinement and molecular model of creatine amidinohydrolase from *Pseudomonas putida*. *J. Mol. Biol.* **204**: 417-433.
- Horbaczewski J (1885) Neue synthese des Kreatins. *Weiner Med. Jahrbdcher*, p459, 1885; cited from *Jahresber. Thier. Chem.* **15**: 86-87.
- Horber FF, Scheidegger J, Frey FJ (1985) Overestimation of renal function in glucocorticosteroid treated patients. *Eur. J. Clin. Pharmacol.* **28**: 537-541.
- Horsburgh AM, Mardlin DP, Turner NL, Henkler R, Strachan N, Glover LA, Paton GI, Killham K (2002) On-line microbial biosensing and fingerprinting of water pollutants. *Biosens. Bioelectron.* **17**: 495-501.

- Hsiue GH, Lu PL, Chen JC (2004) Multienzyme-immobilized modified polypropylene membrane for an amperometric creatinine biosensor. *J. Appl. Poly. Sci.* **92**: 3126-3134.
- Hsiung SK, Chou JC, Sun TP, Pan CW, Chou NH (2010) Dual type potentiometric biosensor. US patent, No. 7758733B2.
- Husdan H, Rapoport A (1968) Estimation of creatinine by the Jaffe reaction: A comparison of three methods. *Clin. Chem.* **14**: 222-238.
- Huskova R, Chrastina P, Adam T, Schneiderka P (2004) Determination of creatinine in urine by tandem mass spectrometry. *Clin. Chim. Acta* **350**: 99-106.
- Ibraheem A, Campbell RE (2010) Designs and applications of fluorescent protein-based biosensors. *Curr. Opin. Chem. Biol.* **14**: 30-36.
- Iijima S (1991) Helical microtubules of graphitic carbon. *Nature* **354**: 56-58.
- Inouye Y, Matsuda Y, Naid T, Arai S, Hashimoto Y, Asano K, Ozaki M, Nakamura S (1986) Purification and characterization of creatinine amidohydrolase of *Alcaligenes* origin. *Chem. Pharm. Bull.* **34**: 269-274.
- Inversen BS, Menne C, White MA, Kristiansen J, Christensen JM, Sabbioni E (1998) Inductively coupled plasma mass spectrometric determination of molybdenum in urine from a Danish population. *Analyst* **123**: 81-85.
- Isildak I, Cubuk O, Altikatoglu M, Yolcu M, Erci V, Tinkilic N (2012) A novel conductometric creatinine biosensor based on solid-state contact ammonium sensitive PVC-NH₂ membrane. *Biochem. Engg. J.* **62**: 34-38.
- Iyengar MR, Coleman DW, Butler TM (1985) Phosphocreatinine, a high-energy phosphate in muscle, spontaneously forms phosphocreatine and creatinine under physiological conditions. *J. Biol. Chem.* **260**: 7562-7567.
- Jacobsen F, Christensen C, Mangensen CE (1980) Evaluation of kidney function after meals. *Lancet* **315**: 319.
- Jaffe M (1886) Ueber den niederschlag welchen pikrinsaure in normalen harn erzeugt und uber eine neue reaction des kreatinins. *Z. Physio. Chem.* **10**: 391-400.
- Jelliffe R (1971) Estimation of creatinine clearance when urine cannot be collected. *Lancet* **297**: 975-976.
- Jelliffe R (1973) Creatinine clearance: Bedside estimate. *Ann. Int. Med.* **79**: 604-605.
- Juan A, Maeder M, Martinez M, Tauler R (2000) Combining hard- and soft-modelling to solve kinetic problems. *Chemom. Intell. Lab. Syst.* **54**: 123-141.

- Julia M, Magalhaes CS, Adelio AS, Machado C (2002) Array of potentiometric sensors for the analysis of creatinine in urine samples. *Analyst* **127**: 1069-1075.
- Jurkiewicz M, Alegret S, Almirall J, Garcia M, Fabregas E (1998b) Development of a biparametric bioanalyser for creatinine and urea. Validation of the determination of biochemical parameters associated with hemodialysis. *Analyst* **123**: 1321-1327.
- Jurkiewicz M, Alegret S, Fabrega E (1998a) Comparison of flow injection analytical biosystems based on open-tube and packed-bed enzyme reactors. *Anal. Chim. Acta.* **370**: 47-58.
- Kampman J, Siersbaek-Nielsen K, Kristensen M, Hansen J (1971) Alderbertingede variationer i urinkreatinine og endogen kreatininclearance. *Ugeskr. Laeg.* 133: 2369-2372.
- Kanakam CC, Raj KP (2011) A simple device to determine creatinine in the blood. *Canad. J. Biomed. Engineer. Technol.* **2**: 1-4.
- Kaplan A, Naugler D (1974) Creatinine hydrolase and creatine amidinohydrolase. I. Presence in cell-free extracts of *Arthrobacter ureafaciens*. *Mol. Cell. Biochem.* **3**: 9-15.
- Kaplan A, Szabo LL (1974) Creatinine hydrolase and creatine amidinohydrolase. II. Partial purification and properties. *Mol. Cell. Biochem.* **3**: 17-25.
- Karon BS, Daly TM, Scott MG (1998) Mechanisms of dopamine and dobutamine interference in biochemical tests that use peroxide and peroxidase to generate chromophore. *Clin. Chem.* **44**: 155-160.
- Karyakin AA, Karyakina EE, Gorton L (1996) Prussian-Blue-based amperometric biosensors in flow-injection analysis. *Talanta* **43**: 1597-1606.
- Kazmlerczak SC, Lente FV, Hodges ED (1991) Diagnostic and prognostic utility of phospholipase: A activity in patients with acute pancreatitis: comparison with amylase and lipase. *Clin. Chem.* **37**: 356-360.
- Keevil BG, Tierney DP, Cooper DP, Morris MR, Machaal A, Yonan N (2002) Simultaneous and rapid analysis of cyclosporine A and creatinine in finger prick blood samples using liquid chromatography tandem mass spectrometry and its application in C2 monitoring. *Ther. Drug Monit.* **24**: 757-767.
- Kennedy JF (1985) Principle of immobilization of enzyme. In: Wiseman A (Ed) *Handbook of enzyme biotechnology*, Wiley, New York: 147-275.

- Khadro B, Sanglar C, Bonhomme A, Errachid A, Jaffrezic-Renault N (2010) Molecularly imprinted polymers (MIP) based electrochemical sensor for detection of urea and creatinine. *Procedia. Engineer.* **5**: 371-374.
- Khan GF, Wernet W (1997) A highly sensitive amperometric creatinine sensor. *Anal. Chim. Acta.* **351**: 151-158.
- Kihara K, Yasukawa E (1986) Determination of creatinine with a sensor based on immobilized glutamate dehydrogenase and creatinine deiminase. *Anal. Chim. Acta.* **183**: 75-80.
- Killard AJ, Malcolm SR (2000) Creatinine biosensors: Principles and designs. *Trends Biotechnol.* **28**: 433-437.
- Kim EJ, Haruyama T, Yanagida Y, Kobatake E, Aizawa M (1999) Disposable creatinine sensor based on thick-film hydrogen peroxide electrode system. *Anal. Chim. Acta.* **394**: 225-231.
- Kinoshita H, Torimura M, Kana K, Ikeda T (1997) Peroxidase-based amperometric sensor of hydrogen peroxide generated in oxidase reaction: application to creatinine and creatine assay. *Electroanalysis* **9**: 1234-1238.
- Klaus R, Fisher W, Hauck HE (1991) Qualitative and quantitative analysis of uric acid, creatine and creatinine together with carbohydrates in biological material by HPTLC. *Chromatographia.* **32**: 307-316.
- Knapp ML, Mayne PD (1987) Development of an automatic kinetic Jaffe method designed to minimise bilirubin interference in plasma creatinine assays. *Clin. Chim. Acta* **168**: 239-246.
- Kochansky C, Koziol S, Strein T (2001) Electrophoretically mediated microanalysis with small molecules: The Jaffé method for creatinine carried out in a capillary tube. *Electrophoresis* **22**: 2518-2525.
- Kostir JV, Rabek V (1950) The detection of Jaffe-positive substances in blood serum by partition chromatography. *Biochim. Biophys. Acta.* **5**: 210-223.
- Kostir JV, Sonka T (1952) Creatinine estimation in blood serum: A new method. *Biochim. Biophys. Acta.* **8**: 86-89.
- Kozitsina AN, Shalygina ZhV, Dedeneva SS, Rusinov GL, Tolshchina SG, Verbitskiy EV, Brainina KhZ (2009) Catalytic systems based on the organic nickel (II) complexes in chronoamperometric determination of urea and creatinine. *Russ. Chem. Bullet.* **58**: 1119-1125.

- Kratochvila J, Friedecky B, Budina M, Sperlingova I (2007) Creatinine determination in urine from the point of view of reference values. *Accred. Qual. Assur.* **12**: 146-150.
- Krolewski AS, Warram JH, Christlieb AR, Busick Ed, Kahn CR (1985) The changing natural history of nephropathy in the type 1 diabetes. *Am. J. Med.* **78**: 785-794.
- Kroll MH, Roach NA, Poe B, Elin RJ (1987) Mechanism of interference with the Jaffe reaction for creatinine. *Clin. Chem.* **33**: 1129-1132.
- Kubo I, Karube I (1986) Immobilization of creatinine deiminase on a substituted poly(methylglutamate) membrane and its use in a creatinine sensor. *Anal. Chim. Acta.* **187**: 31-37.
- Kubo I, Karube I, Suzuki S (1983) Amperometric determination of creatinine with a biosensor based on immobilized creatininase and nitrifying bacteria. *Anal. Chim. Acta.* **151**: 371-376.
- Kugimiya A, Takeuchi T (1999) Molecularly imprinted polymer-coated quartz crystal microbalance for detection of biological hormone. *Electroanalysis* **11**: 1158-1160.
- Kumar N, Ananthi A, Mathiyarasu J, Joseph J, Phani KL, Yegnaraman V (2011) Enzymeless creatinine estimation using poly(3,4-ethylenedioxythiophene)- β -cyclodextrin. *J. Electroanal. Chem.* **661**: 303-308.
- Lad U, Khokhar S, Kale GM (2008) Electrochemical creatinine biosensors. *Anal. Chem.* **80**: 7910-7917.
- Lee EK, Cho YM, Kim JT, Koo BK, Cho HY, Ku YH, Park KS, Jang HC, Kim SY, Lee HK (2009) Derivation of a new equation for estimating creatinine clearance by using fat-free mass and serum creatinine concentration in Korean patients with type 2 diabetes mellitus. *Diabetes Res. Clin. Practice* **83**: 44-49.
- Lee MH, Tsaib TC, Thomasc JL, Linb HY (2008) Recognition of creatinine by poly(ethylene-co-vinylalcohol) molecular imprinting membrane. *Desalination* **234**: 126-133.
- Lempert C (1959) The chemistry of the glycoamidines. *Chem. Rev.* **59**: 667-736.
- Levene PA, Kristeller L (1909) Factors regulating the creatinine output in man. *Am. J. Physiol.* **24**: 45-65.
- Levey A, Bosch J, Breyer-Lewis J, Greene T, Rogers N, Roth D (1999) A more accurate method to estimate glomerular filtration rate from serum creatinine: A new prediction equation. *Ann. Int. Med.* **130**: 2369-2372.

- Levey A, Coresh J, Greene T, Marsh J, Stevens LA, Kusek JW, Van Lente F (2007) Expressing the Modification of Diet in Renal Disease study equation for estimating glomerular filtration rate with standardized serum creatinine values. *Clin. Chem.* **53**: 766-772.
- Li Y, Wang S, Zhong N (1992) Simultaneous determination of pseudouridine and creatinine in urine of normal children and patients with leukaemia by high performance liquid chromatography. *Biomed. Chromatogr.* **6**: 191-193.
- Liebig J (1847) Kreatin und Kreatinin, Bestandtheile des Hams der Menschen. *J. Prakt. Chem.* **40**: 288-292.
- Lim CK, Richmond W, Robinson DP, Brown SS (1978) Towards a definitive assay of creatinine in serum and urine: Separation by high-performance liquid chromatography. *J. Chromatogr.* **145**: 41-49.
- Lin Z, Chen Z, Liu Y, Wang J, Chen G (2008) An electrochemiluminescent biosensor for uric acid based on the electrochemiluminescence of bis-[3,4,6-trichloro-2-(pentylloxycarbonyl)-phenyl] oxalate on an ITO electrode modified by an electropolymerized nickel phthalocyanine film. *Analyst* **133**: 797-801.
- Lindback B, Bergman A (1989) A new commercial method for the enzymatic determination of creatinine in serum and urine evaluated: Comparison with kinetic Jaffe method and isotope dilution-mass spectrometry. *Clin. Chem.* **35**: 835-837.
- Linnet K, Bruunshuus I (1991) HPLC with enzymatic detection as a candidate reference method for serum creatinine. *Clin. Chem.* **37**: 1669-1675.
- Liotta E, Gottardo R, Bonizzato L, Pascali JP, Bertaso A, Tagliaro F (2009) Rapid and direct determination of creatinine in urine using capillary zone electrophoresis. *Clin. Chim. Acta.* **409**: 52-55.
- Liu L, Zhou HY, He L, Song JW, Nie WX, Su ZY (2006) Measurement of urinary S100B protein levels and lactate/creatinine ratio in early detection of neonatal hypoxic-ischemic encephalopathy. *World J. Pediatr.* **2**: 270-275.
- Lolekha P, Taksinamanee R (1980) Evaluation of serum creatinine measurements by the direct acidification method for errors contributed by non-creatinine chromogens. *Clin. Chim. Acta.* **107**: 97-104.
- Lustgarten JA, Wenk RE (1972) Simple, rapid, kinetic method for serum creatinine measurement. *Clin. Chem.* **18**: 1419-1422.

- Lvova L, Martinelli E, Dini F, Bergamini A, Paolesse R, Di-Natale C, D'Amico (2009) A Clinical analysis of human urine by means of potentiometric electronic tongue. *Talanta* **77**: 1097-1104.
- Madaras MB, Buck RP (1996) Miniaturized biosensors employing electropolymerized permselective films and their use for creatinine assays in human serum. *Anal. Chem.* **68**: 3832-3839.
- Madaras MB, Popescu IC, Ufer S, Buck RP (1996) Microfabricated amperometric creatine and creatinine biosensors. *Anal. Chim. Acta.* **319**: 335-345.
- Magalhaes JMCS, Machado AASC (2002) Array of potentiometric sensors for the analysis of creatinine in urine samples. *Analyst* **127**: 1069-1075.
- Malhotra BD, Chaubey A, Singh SP (2006) Prospects of conducting polymers in biosensors. *Anal. Chim. Acta.* **578**: 59-74.
- Mascini M, Fortunati S, Moscone D, Palleschi G (1985) Ammonia abatement in an enzymatic flow system for the determination of creatinine in blood sera and urine. *Anal. Chim. Acta.* **171**: 175-184.
- Mascini M, Palleschi G (1982) Determination of creatinine in clinical samples with a creatininase reactor and an ammonia probe. *Anal. Chim. Acta.* **136**: 69-76.
- Masson P, Ohlsson P, Bjorkhem I (1981) Combined enzymic-Jaffe method for determination of creatinine in serum. *Clin. Chem.* **27**: 18-21.
- Matharu Z, Sumana G, Arya SK, Singh BP, Gupta V, Malhotra BD (2007) Polyaniline Langmuir–Blodgett film based cholesterol biosensor. *Langmuir* **23**: 13188-13192.
- Matsuda Y, Wakamatsu N, Inouye Y, Uede S, Hashimoto Y, Asano K, Nakamura S (1986) Purification and characterization of creatine amidinohydrolase of *Alcaligenes* origin. *Chem. Pharm. Bull.* **34**: 2155-2160.
- Mayerhoff M, Rechnitz GA (1976) An activated enzyme electrode for creatinine. *Anal. Chim. Acta.* **85**: 277-285.
- McLellan M, Gallwas J, Hendrixson M (1973) Evaluation of an automated creatinase creatinine procedure. *Clin. Chem.* **19**: 623-625.
- Mehrvar M, Bis C, Scharer JM, Moo-Young M, Loung JH (2000) Fiber optic biosensors- Trends and advances. *Anal. Sci.* **16**: 677-691.
- Miller BF, Dubos R (1937) Determination by a specific enzymatic method of the creatinine content of blood and urine from normal and nephritic individuals. *J. Biol. Chem.* **121**: 457-464.

- Mitchell RJ (1973) Improved method for specific determination of creatinine in serum and urine. *Clin. Chem.* **19**: 408-410.
- Miyake M, Shibukawa A, Nakagawa T (1991) Simultaneous determination of creatinine and uric acid in human plasma and urine by micellar electrokinetic chromatography. *J. High Resolut. Chromatogr.* **14**: 181-185.
- Miyoshi K, Taira A, Yoshida K, Tamura K, Uga S (1980a) Abnormalities of creatinase in skeletal muscle of patients with Duchenne muscular dystrophy. *Proc. Jpn. Acad.* **56**: 99-101.
- Miyoshi K, Taira A, Yoshida K, Tamura K, Uga S (1980b) Presence of creatinase and sarcosine dehydrogenase in human skeletal muscle. Proposal for creatine-urea pathway. *Proc. Jpn. Acad.* **56**: 95-98.
- Monler JL, Barton SD, Blouin RA, Cowen DL, Flanigan RC (1986) The evaluation of creatinine clearance in spinal cord injury patients. *J. Urol.* **136**: 366-369.
- Morgan D, Dillon S, Payne R (1978) The assessment of glomerular function: Creatinine clearance or plasma creatinine? *Postgrad. Med. J.* **54**: 302-310.
- Morrison J, Nayiager T, Webber CE, Sala A, Barr R (2011) Creatinine as a measure of lean body mass during treatment of acute lymphoblastic leukemia in childhood. *J. Pediatr. Hematol. Oncol.* **33**: e13-e16.
- Morrison JF, Ennor AH (1960) N-phosphorylated guanidines. In: *The Enzymes* (2nd Ed.), edited by P. D. Boyer, H. Lardy, and K. Myrback. New York: Academic, 89-109.
- Moss GA, Bondar RJL, Buzzeli DM (1975) Kinetic enzymatic method for determining serum creatinine. *Clin. Chem.* **21**: 1422-1426.
- Mottaghitlab V, Xi B, Spinks GM, Wallace GG (2006) Polyaniline fibres containing single walled carbon nanotubes: Enhanced performance artificial muscles. *Synth. Metals* **156**: 796-803.
- Munkhomm C (1999) Fluorescent polymeric sensor for the detection of creatinine. US patent, No. 5,958,786.
- Murray WB, Mackay C (1977) The amylase creatinine clearance ratio in acute pancreatitis. *Br. J. Surg.* **64**: 189-191.
- Murzin AG (1993) Can homologous proteins evolve different enzymatic activities? *Trends Biochem. Sci.* **18**: 403-405.

- Myers VC, Fine MS (1919) Comparative distribution of urea, creatinine, uric acid and sugar in blood and spinal fluid. *J. Biol. Chem.* **37**: 239-244.
- Nguyen VK, Wolff CM, Seris JL, Schwing JP (1991) Immobilized enzyme electrode for creatinine determination in serum. *Anal. Chem.* **63**: 611-614.
- Nishiya Y, Toda A, Imanaka T (1998) Gene cluster for creatinine degradation in *Arthrobacter* sp. TE1826. *Mol. Gen. Genet.* **257**: 581-586.
- Ogata M, Taguchi T (1988) Simultaneous determination of urinary creatinine and metabolites of toluene, xylene, styrene, ethylbenzene and phenol by automated high performance liquid chromatography. *Int. Arch. Occup. Environ. Health.* **61**: 131-140.
- Ohira SI, Kirk AB, Dasgupta PK (2009) Automated measurement of urinary creatinine by multichannel kinetic spectrophotometry. *Anal. Biochem.* **384**: 238-244.
- Osaka T, Komaba S, Amano A (1998) Highly sensitive microbiosensor for creatinine based on the combination of inactive polypyrrole with polyion complexes. *J. Electrochem. Soc.* **145**: 406-408.
- Osaka T, Komaba S, Amano A, Fujino Y, Mori H (2000) Electrochemical molecular sieving of the polyion complex film for designing highly sensitive biosensor for creatinine. *Sens. Actuat. B* **65**: 58-63.
- Osborne MD, Girault HH (1995) The micro water/1,2-dichloroethane interface as a transducer for creatinine assay. *Mikrochim. Acta.* **117**: 175-185.
- Owen LJ, Wear JE, Keevil BG (2006) Validation of a liquid chromatography tandem mass spectrometry assay for serum creatinine and comparison with enzymatic and Jaffe methods. *Ann. Clin. Biochem.* **43**: 118-123.
- Pacakava V, Stulik K, Kluodova M, Jelinkine K (1993) High-performance liquid chromatographic determination of creatinine in serum, and a correlation of the results with those of the Jaffe and enzymic methods. *J. Chromatogr. B: Biomed. Sci. Appl.* **614**: 221-226.
- Panasyuk-Delaney T, Mirsky VM, Otto WS (2002) Capacitive creatinine sensor based on a photografted molecularly imprinted polymer. *Electroanalysis* **14**: 221-224.
- Pandey PC, Mishra AP (2004) Novel potentiometric sensing of creatinine. *Sens. Actuat. B* **99**: 230-235.
- Pardue HL, Bacon BL, Nevius MG, Skoug JW (1987) Kinetic study of the Jaffe reaction for quantifying creatinine in serum: I. Alkalinity controlled with NaOH. *Clin. Chem.* **33**: 278-285.

- Parekh A, Cook S, Sims C (1976) A new method for the determination of serum creatinine based on reaction with 3,5-dinitrobenzyl chloride in an organic medium. *Clin. Chem. Acta.* **73**: 221-231.
- Parekh A, Sims C (1977) Serum creatinine assay by use 3,5-dinitrobenzoate: A critique. *Clin. Chem.* **23**: 2066-2071.
- Park EK, Watanabe T, Gee SJ, Schenker MB, Hammock BD (2008) Creatinine measurements in 24 h urine by liquid chromatography-tandem mass spectrometry. *J. Agric. Food Chem.* **56**: 333-336.
- Paroni R, Arcelloni C, Fermo I, Bonini PA (1990) Determination of creatinine in serum and urine by a rapid liquid-chromatographic method. *Clin. Chem.* **36**: 830-836.
- Paroni R, Fermo I, Cighetti G, Ferrero CA, Carobene A, Ceriotti F (2004) Creatinine determination in serum by capillary electrophoresis. *Electrophoresis* **25**: 463-468.
- Paulmann W (1894) Beitrage zur Kenntniss des Sarkosins. *Arch. Pharm.* **232**: 601-639.
- Perakis N, Wolff CM (1984) Kinetic approach for the enzymic determination of creatinine. *Clin Chem.* **30**: 1792-1796.
- Perrone RD, Madias NE, Levey AS (1992) Serum creatinine as an index of renal function: New insights into old concepts. *Clin. Chem.* **38**: 1933-1953.
- Pieroni L, Delanaye P, Boutten A, Bargnoux AS, Rozet E, Delatour V, Carlier MC, Hanser AM, Cavalier E, Froissart M, Cristol JP (2011) A multicentric evaluation of IDMS-traceable creatinine enzymatic assays. *Clin. Chim. Acta.* **412**: 2070-2075.
- Pobozy E, Radomska A, Koncki R, Glab S (2003) Determination of dialysate creatinine by micellar electrokinetic chromatography. *J. Chromatogr. B* **789**: 417-424.
- Polyak B, Bassis E, Novodvoretz A, Belkin S, Marks RS (2001) Bioluminescent whole cell optical fiber sensor to genotoxicants: system optimization. *Sens. Actuat. B* **74**: 18-26.
- Pookaiyaudom P, Seelanan P, Lidgey FJ, Hayatleh K, Toumazou C (2011) Measurement of urea, creatinine and urea to creatinine ratio using enzyme based chemical current conveyor (CCCII⁺). *Sens. Actuat. B* **153**: 453-459.
- Premanode B, Toumazou C (2007) A novel, low power biosensor for real time monitoring of creatinine and urea in peritoneal dialysis. *Sens. Actuat. B* **120**: 732-735.
- Puig JG, Anton FM, Grande C, Pallardo LF, Arnalich F, Gil A, Vazquez JJ, Montero-Garcia A (1981) Relation of kidney size to kidney function in early insulin dependent diabetes. *Diabetologia.* **21**: 363-367.

- Quillard S, Louarn G, Lefrant S, Macdiarmad AG (1994) Vibrational analysis of polyaniline: A comparative study of leucoemeraldine, emeraldine, and pernigraniline bases. *Phys. Rev. B* **50**: 12496-12508.
- Quinn A, Gemperline PJ, Baker B, Zhu M, Walker DS (1999) Fiber-optic UV/visible composition monitoring for process control of batch reactions. *Chemom. Intell. Lab. Syst.* **45**: 199-214.
- Radomska A, Bodenszac E, Glab S, Koncki R (2004a) Creatinine biosensor based on ammonium ion selective electrode and its application in flow-injection analysis. *Talanta* **64**: 603-608.
- Radomska A, Koncki R, Pyrzyńska K, Glab S (2004b) Bioanalytical system for control of hemodialysis treatment based on potentiometric biosensors for urea and creatinine. *Anal. Chim. Acta.* **523**: 193-200.
- Rahman MM, Muhammad JA, Shiddiky, Rahman MA, Shim YB (2009b) A lactate biosensor based on lactate dehydrogenase/nicotinamide adenine dinucleotide (oxidized form) immobilized on a conducting polymer/multiwall carbon nanotube composite film. *Anal. Biochem.* **384**: 159-165.
- Rahman MM, Umar A, Sawada K (2009a) Development of amperometric glucose biosensor based on glucose oxidase co-immobilized with multi-walled carbon nanotubes at low potential. *Sens. Actuat. B* **137**: 327-333.
- Ramanavicius A (2007) Amperometric biosensor for the determination of creatine. *Anal. Bioanal. Chem.* **378**: 1899-1906.
- Rapoport A, Husdan H (1968) Endogenous creatinine clearance and serum creatinine in the clinical assessment of kidney function. *Can. Med. Assoc. J.* **99**: 149-156.
- Rasmussen CD, Andersen JET, Zachau-Christiansen B (2007) Improved performance of the potentiometric biosensor for the determination of creatinine. *Anal. Lett.* **40**: 39-52.
- Ravel R (1978) *Clinical Laboratory Medicine, Clinical Application of Laboratory Data*, 3rd Ed. Year Book Medical Publishers, Inc., Chicago, IL, 136.
- Razumas V, Kanapieniene J, Nylander T, Engstrom S, Larsson K (1994) Electrochemical biosensors for glucose, lactate, urea, and creatinine based on enzymes entrapped in a cubic liquid crystalline phase. *Anal. Chim. Acta.* **289**: 155-162.
- Rehberg PB (1926) Studies on kidney function. I. The rate of filtration and reabsorption in the human kidney. *Biochem. J.* **20**: 447-460.
- Renkin E, Robinson R (1974) Glomerular filtration. *New Engl. J. Med.* **290**: 785-792.

- Rikitake K, Oka I, Ando M, Yoshimoto T, Tsuru D (1979) Creatinine amidohydrolase (creatininase) from *Pseudomonas putida*. Purification and some properties. *J. Biochem.* **86**: 1109-1117.
- Ripamonti M, Mosca A, Rovida E, Luzzan M, Luzzi L, Ceriotti F, Cottini F, Rossi-Bernardi L (1984) Urea, creatinine and glucose determined in plasma and whole blood by differential pH technique. *Clin. Chem.* **30**: 556-559.
- Robert R, Alexander N, Kelner M (1988) Definitive liquid chromatographic demonstration that N-ethylglycine I sthe metabolite of lidocaine that interferes in the Kodak sarcosine oxidase-coupled method for creatinine. *Clin. Chem.* **34**: 2569-2572.
- Roberts JJ, Walker JB (1985) Higher homolog and N-ethyl analog of creatine as synthetic phosphagen precursors in brain, heart, and muscle, repressors of liver amidinotransferase, and substrates for creatine catabolic enzymes. *J. Biol. Chem.* **260**: 13502-13508.
- Romer J (1975) Evaluation of a kinetic method for creatinine. *Lab. Med.* **6**: 15-18.
- Rosano T, Brown H (1982) Analytical and biological variability of serum creatinine and creatinine clearance: Implication for interpretation. *Clin. Chem.* **28**: 2330-2331.
- Rosano TG, Ambrose RT, Wu AH, Swift TA, Yadegari P (1990) Candidate reference method for determining creatinine in serum: Method development and inter laboratory validation. *Clin. Chem.* **36**: 1951-1955.
- Roscoe MH (1953) The estimation of creatinine in serum. *J. Clin. Path.* **6**: 201-207.
- Rudnick ME, Bastl CP, Elfinbein IB, Narins HG (1983) The differential diagnosis of acute renal failure. In: Brenner BM, Lazarus JM, Eds. Acute renal failure. Philadelphia: WB Saunders, 176-222.
- Rui CS, Ogawa HI, Sonomoto K, Kato Y (1993b) Multifunctional flow-injection biosensor for the simultaneous measurement of creatinine, glucose and urea. *Biosci. Biotech. Biochem.* **57**: 191-194.
- Rui CS, Sonomoto K, Kato Y (1992) Amperometric flow-injection biosensor system for the simultaneous determination of urea and creatinine. *Anal. Sci.* **8**: 845-850.
- Rui CS, Sonomoto K, Ogawa HI, Kato Y (1993a) A flow-injection biosensor system for the amperometric determination of creatinine: Simultaneous compensation of endogenous interferents. *Anal. Biochem.* **210**: 163-171.

- Ruiz-Jimenez J, Mata-Granados JM, Luque de Castro MD (2007) On-line automatic SPE–CE coupling for the determination of biological markers in urine. *Electrophoresis* **28**: 789-798.
- Sadilek L (1965) Creatinine determination in the urine of alcaptonuric patients. *Clin Chim. Acta.* **12**: 436-439.
- Sakai T, Ohta H, Ohno N, Imai J (1995) Routine assay of creatinine in newborn baby urine by spectrophotometric flow-injection analysis. *Anal. Chim. Acta.* **308**: 446-450.
- Sakslund H, Hammerich O (1992) Effects of pH, temperature and reaction products on the performance of an immobilized creatininase-creatinase-sarcosine oxidase enzyme system for creatinine determination. *Anal. Chim. Acta.* **268**: 331-345.
- Sant W, Pourciel-Gouzy ML, Launay J, Do Conto T, Colin R, Martinez A, Temple-Boyer P (2004) Development of a creatinine-sensitive sensor for medical analysis. *Sens. Actuat. B* **103**: 260-264.
- Scheller FW, Schubert F, Neumann B, Pfeiffer D, Hintsche R, Dransfeld I (1991) Second generation biosensors. *Biosens. Bioelectron.* **6**: 245-253.
- Schenk PW, Cransberg K, Wolff ED, de Rijke YB (2007) Point-of-care creatinine testing in children at risk for sudden deterioration of renal function. *Clin. Chem. Lab. Med.* **45**: 1536-1541.
- Schneider J, Grhdig B, Renneberg R, Cammann K, Madaras MB, Buck RP, Vorlop KD (1996) Hydrogel matrix for three enzyme entrapment in creatine/creatinine amperometric biosensing. *Anal. Chim. Acta.* **325**: 161-167.
- Schoucci R, Poliot M (1977) Some observations on the kinetics of the Jaffe reaction for creatinine. *Clin. Chem.* **23**: 1527-1530.
- Schumann R (1931) Uber das Vorkommen von Kreatinin und Kreatin im menschlichen Schweibe. *Z. Ges. Exp. Med.* **79**: 145-152.
- Schwartz G, Brion L, Spitzer A (1987) The use of plasma creatinine concentration for estimating glomerular filtration rate in infants, children and adolescents. *Pediatr. Clin. North Am.* **34**: 571-590.
- Schwartz GJ, Kwong T, Erway B, Warady B, Sokoll L, Hellerstein S, Dharnidharka V, Furth S, Munoz A (2009) Validation of creatinine assays utilizing HPLC and IDMS traceable standards in sera of children. *Pediatr. Nephrol.* **24**: 113-119.
- Scott PH (1992) High-performance liquid chromatography measurement of plasma creatinine in newborns. *Clin. Chem.* **38**: 101-103.

- Shih YT, Huang HJ (1999) A creatinine deiminase modified polyaniline electrode for creatinine analysis. *Anal. Chim. Acta.* **392**: 143-150.
- Shimizu S, Kim JM, Shinmen Y, Yamada H (1986) Evaluation of two alternative metabolic pathways for creatinine degradation in microorganisms. *Arch. Microbiol.* **145**: 322-328.
- Shin JH, Choi YS, Lee HJ, Choi SH, Ha J, Yoon IJ, Nam H, Cha GS (2001) A planar amperometric creatinine biosensor employing an insoluble oxidizing agent for removing redox-active interferences. *Anal. Chem.* **73**: 5965-5971.
- Shirao MK, Suzuki S, Kobayashi J, Nakazawa H, Mochizuki E (1997) Analysis of creatinine, vanilmandelic acid, homovanillic acid and uric acid in urine by micellar electrokinetic chromatography. *J. Chromatogr. B* **693**: 463-467.
- Siekmann L (1985) Determination of creatinine in human serum by isotope dilution-mass spectrometry. *J. Clin. Chem. Clin. Biochem.* **23**: 137-44.
- Sims C, Parekh AC (1977) Determination of serum creatinine by reaction with methyl-3,5-dinitrobenzoate in methyl sulfoxide. *Ann. Clin. Biochem.* **14**: 227-232.
- Sims EAR, Krantz KE (1958) Serial studies of renal function during pregnancy and the puerperium in normal women. *J. Clin. Invest.* **37**: 1764-1774.
- Slot C (1965) Plasma creatinine determination. A new specific Jaffe reaction method. *Scand. J. Clin. Lab. Invest.* **17**: 381-387.
- Soldatkin AP, Montoriol J, Sant W, Martelet C, Jaffrezic-Renault N (2002a) Development of potentiometric creatinine-sensitive biosensor based on ISFET and creatinine deiminase immobilised in PVA/SbQ photopolymeric membrane. *Mat. Sci. Engineer. C* **21**: 75-79.
- Soldatkin AP, Montoriol J, Sant W, Martelet C, Jaffrezic-Renault N (2002b) Creatinine sensitive biosensor based on ISFETs and creatinine deiminase immobilised in BSA membrane. *Talanta* **58**: 351-357.
- Soldin S, Henderson L, Hill J (1978) The effect of bilirubin and ketones on reaction rate methods for the measurements of creatinine. *Clin. Biochem.* **12**: 18-21.
- Soldin S, Wan B, Cherian A (1981) The measurement of creatinine: A comparison between the Beckman Creatinine Analyzer II and the Selective Analyzer GSA IID. *Clin. Biochem.* **14**: 165-168.
- Soldin SJ, Hill JG (1978) Micromethod for determination of creatinine in biological fluids by high-performance liquid chromatography. *Clin. Chem.* **24**: 747-750.

- Sonntag O (1991) Die Bestimmung der Creatinin-Konzentration in Serum und Urin. Kritische Übersicht der Routine-Bestimmungsmethoden. *D.G. Klin. Chem. Mitteilungen*. **22**: 235-251.
- Spencer K (1986) Analytical reviews in clinical biochemistry: The estimation of creatinine. *Ann. Clin. Biochem.* **23**: 1-25.
- Spencer K, Price CP (1980) A review of non-enzyme mediated reactions and their applications to centrifugal analysers. In: Price C.P., Spencer K. Eds. *Centrifugal Analysers in Clinical Chemistry*, Easrbourne, Praeger: 231-253.
- Squretti DJ, Price RL, Murphy MJ (2002) Rapid and specific detection of bacteria using bioluminescence. *Anal. Chim. Acta.* **457**: 109-114.
- Sreenivasan K, Sivakumar R (1997) Interaction of molecularly imprinted polymers with creatinine. *J. Appl. Poly. Sci.* **66**: 2539-2542.
- Stefan RI, Bokretson RG, van Staden JF, Aboul-Enein HY (2003) Simultaneous determination of creatine and creatinine using amperometric biosensors. *Talanta* **60**: 1223-1228.
- Stefan-van Staden RI, Bokretson RG, van Staden JF, Aboul-Enein HY (2006) Simultaneous detection of creatine and creatinine using a sequential injection analysis/biosensor system. *Prep. Biochem. Biotechnol.* **36**: 287-296.
- Stokes P, O'Connor G (2003) Development of a liquid chromatography–mass spectrometry method for the high-accuracy determination of creatinine in serum. *J. Chromatogr. B* **794**: 125-136.
- Stosch R, Henrion A, Schiel D, Guttler B (2005) Surface-enhanced Raman scattering based approach for quantitative determination of creatinine in human serum. *Anal. Chem.* **77**: 7386-7392.
- Su X, Dai C, Zhang J, O'Shea SJ (2002) Quartz tuning fork biosensor. *Biosens. Bioelectron.* **17**: 111-117.
- Subrahmanyam S, Piletsky SA, Piletska EV, Chen B, Karim K, Turner APF (2001) 'Bite-and-Switch' approach using computationally designed molecularly imprinted polymers for sensing of creatinine. *Biosens. Bioelectron.* **16**: 631-637.
- Sugita O, Uchyama K, Yamada T, Sato T, Okada M, Takeuchi K (1992) Reference values of serum and urine creatinine and creatinine clearance by a new enzymatic method. *Ann. Clin. Biochem.* **29**: 523-528.

- Sullivan M, Irreverre F (1958) A highly specific test for creatinine. *J. Biol. Chem.* **233**: 530-533.
- Suzuki H, Arakawa H, Karube I (2001) Fabrication of a sensing module using micromachined biosensors. *Biosens. Bioelectron.* **16**: 725-733.
- Suzuki H, Matsugi Y (2004) Microfabricated flow system for ammonia and creatinine with an air-gap structure. *Sens. Actuat. B* **98**: 101-111.
- Suzuki H, Matsugi Y (2005) Integrated microfluidic system for the simultaneous determination of ammonia, creatinine, and urea. *Sens. Actuat. B* **108**: 700-707.
- Suzuki K, Sagai H, Sugiyama M, Imamura S (1993) Molecular cloning and high expression of the Bacillus creatinase gene in Escherichia coli. *J. Ferment. Bioeng.* **76**: 77-81.
- Tabata M, Kido T, Totani M, Murachi T (1983) Automated assay of creatinine in serum as simplified by the use of immobilized enzymes creatinine deaminase and glutamate dehydrogenase. *Anal. Chem.* **134**: 44-49.
- Takatsu A, Nishi S (1993) Determination of serum creatinine by isotope dilution method using discharge-assisted thermospray liquid chromatography/mass spectrometry. *Biol. Mass Spectrom.* **22**: 643-646.
- Tanganelli E, Prencipe L, Bassi D, Cambiaghi S., Murador E (1982) Enzymic assay of creatinine in serum and urine with creatinine iminohydrolase and glutamate dehydrogenase. *Clin. Chem.* **28**: 1461-1464.
- Tausky H (1954) A microcolorimetric determination of creatinine in urine by Jaffe reaction. *J. Clin. Chem.* **208**: 853-861.
- Tausky H (1956) A procedure for increasing the specificity of the Jaffe reaction for the determination of creatine and creatinine in urine and plasma. *Clin. Chim. Acta.* **1**: 210-224.
- Tedesco FJ, Harter HR, Alpers DH (1976) Serum amylase determinations and amylase to creatinine clearance ratios in patients with chronic renal insufficiency. *Gastroenterology* **71**: 594-598.
- Thevenot DR, Toth K, Durs RA, Wilson GS (2001) Electrochemical biosensors: recommended definitions and classification. *Biosens. Bioelectron.* **16**: 121-131.
- Thienpont LM, Van-Landuyt KG, Stock D, De-Leenheer AP (1995) Candidate reference method for determining serum creatinine by isocratic HPLC: Validation with isotope

- dilution gas chromatography-mass spectrometry and application for accuracy assessment of routine test kits. *Clin. Chem.* **41**: 995-1003.
- Thienpont LM, Van-Nieuwenhove B, Stockl D, Reinauer H, De-Leenheer AP (1996) Determination of reference method values by isotope dilution-gas chromatography/mass spectrometry: a five years' experience of two European Reference Laboratories. *Eur. J. Clin. Chem. Clin. Biochem.* **34**: 853-860.
- Thompson H, Rechnitz G (1974) Ion electrode based enzymatic analysis of creatinine. *Anal. Chem.* **46**: 246-249.
- Tietz NW (1986) Tietz Textbook of Clinical Chemistry, 1st Ed.; W. B. Saunders: Philadelphia, PA, 1810-1857.
- Tiwari A, Dhakate SR (2009) Chitosan-SiO₂-multiwall carbon nanotubes nanocomposite: A novel matrix for the immobilization of creatine amidinohydrolase. *Int. J. Biol. Macromolecul.* **44**: 408-412.
- Tiwari A, Shukla SK (2009) Chitosan-g-polyaniline: A creatine amidinohydrolase immobilization matrix for creatine biosensor. *eXPRESS Poly. Letts.* **3**: 553-559.
- Toffaletti J, Blosser N, Hall T, Smith S, Tompkin D (1983) An automated dryslide enzymatic method evaluated for measurement of creatinine in serum. *Clin. Chem.* **29**: 684-687.
- Tombach B, Schneider J, Matzkies F, Schaefer RM, Chemnitz GC (2001) Amperometric creatinine biosensor for hemodialysis patients. *Clin. Chim. Acta.* **312**: 129-134.
- Tomson R, Uhlin F, Holmar J, Lauri K, Luman M, Fridolin I (2011) Development of a method for optical monitoring of creatinine in the spent dialysate. *Eston. J. Engineer.* **17**: 140-150.
- Trojanowicz M, Lewenstam A, Vel-Krawczyk TK, Lahdesmaki I, Szczepk W (1996) Flow injection amperometric detection of ammonia using a polypyrrole-modified electrode and its application in urea and creatinine biosensors. *Electroanalysis* **9**: 233-243.
- Tsai HA, Syu MJ (1995) Synthesis of creatinine-imprinted poly(β -cyclodextrin) for the specific binding of creatinine. *Biomater.* **26**: 2759-2766.
- Tsai HA, Syu MJ (2004) Synthesis and characterization of creatinine imprinted poly(4-vinylpyridine-co-divinylbenzene) as a specific recognition receptor. *Anal. Chim. Acta.* **539**: 107-116.

- Tsai HA, Syu MJ (2011) Preparation of imprinted poly(tetraethoxysilanol) sol–gel for the specific uptake of creatinine. *Chem. Engineer. J.* **168**: 1369-1376.
- Tsikas D, Wolf A, Frolich JC (2004) Simplified HPLC method for urinary and circulating creatinine. *Clin. Chem.* **50**: 201-203.
- Tsuchida T, Yoda K (1983) Multi-enzyme membrane electrodes for determination of creatinine and creatine in serum. *Clin. Chem.* **29**: 51-55.
- Tsuru D, Oka I, Yoshimoto T (1976) Creatinine decomposing enzymes in *Pseudomonas putida*. *Agric. Biol. Chem.* **40**: 1011-1018.
- Urdike SJ, Hicks GP (1967) The enzyme electrode. *Nature* **214**: 986-988.
- Vaddirajua S, Tomazos I, Burgess DJ, Jain FC, Papadimitrakopoulou F (2010) Emerging synergy between nanotechnology and implantable biosensors: A review. *Biosens Bioelectronics.* **25**: 1553-1565.
- Van-Landuyt K, Thienpoint L, Leenheer A, Stockl D (1994) Determination of creatinine in human serum using isocratic HPLC with an aluminium oxide stationary phase. *J. Chromatogr. Sci.* **32**: 294-297.
- Van-Nierkerk BDH, Reid JT, Bensadoun A, Paladines OL (1963) Urinary creatinine as an index of body composition. *J. Nutr.* **79**: 463-473.
- Van-Pilsum JF, Martin RP, Kito E, Hess J (1956) Determination of creatine, creatinine, arginine, guanidinoacetic acid, guanidine and methyl guanidine in biological fluids. *J. Biol. Chem.* **222**: 225-236.
- Van-Stekelenburg GJ, Valk C, Van-de-Kamp JS (1978) A new automated method for the determination of true creatinine in serum by means of the centrifugal analyser, based on Slot's principle, with special reference to low substrate concentration. *Clin. Chim. Acta.* **89**: 79-91.
- Verhelst J, Berwaerts J, Marescau B, Abs R, Neels H, Mahler C, De-Deyn PP (1997) Serum creatine, creatinine, and other guanidino compounds in patients with thyroid dysfunction. *Metabolism* **46**: 1063-1067.
- Viraraghavan S, Blass KG (1990) Effect of glucose upon alkaline picrate: A Jaffe interference. *J. Clin. Chem. Clin. Biochem.* **28**: 95-105.
- Vuorinen P, Harmoinen APT, Jokela H (1987) Elimination of interference from bilirubin in the kinetic Jaffe reaction for determining creatinine in serum. *Ann. Clin. Biochem.* **22**: 212-215.

- Walsh DA, Dempsey E (2002) Comparison of electrochemical, electrophoretic and spectrophotometric methods for creatinine determination in biological fluids. *Anal. Chim. Acta.* **459**: 187-198.
- Wang J, Chatrathi MP (2003) Microfabricated electrophoresis chip for bioassay of renal markers, *Anal. Chem.* **75**: 525-529.
- Wang YZ, Joo J, Hsu CH, Epstein AJ (1995) Charge transport of camphor sulfonic acid-doped polyaniline and poly(*o*-toluidine) fibers: role of processing. *Synth. Met.* **68**: 207-211.
- Weber JA, Zanten AP (1991) Interferences in current methods for measurements of creatinine. *Clin. Chem.* **37**: 695-700.
- Wei C, Srivastava D, Cho K (2002) Thermal expansion and diffusion coefficients of carbon nanotube-polymer composites. *Nano Lett.* **2**: 647-650.
- Wei Y, Hariharan R, Patel SA (1990a) Chemical and electrochemical co-polymerization of aniline with alkyl ring substituted anilines. *Macromolecules* **23**: 758-764.
- Wei Y, Jang GW, Chan CC, Hsuen KF, Hariharan R, Patel SA, Whitecar CK (1990b) Polymerization of aniline in the presence of aromatic additives. *J. Phys. Chem.* **94**: 7716-7721.
- Wei Y, Tang X, Sun Y, Focke WW (1989) A study of the mechanism of aniline polymerization. *J. Polym. Sci.* **27**: 2385-2396.
- Welch MJ, Cohen A, Hertz HS, Ng KJ, Schaffer R, Van-der-Lijn P, White VE (1986) Determination of serum creatinine by isotope dilution mass spectrometry as a candidate definitive method. *Anal. Chem.* **58**: 1681-1685.
- Werner ER, Fuchs D, Hausen A, Reibenegger G, Wachter H (1987) Simultaneous determination of neopterin and creatinine in serum with solid-phase extraction and on line elution liquid chromatography. *Clin. Chem.* **33**: 2028.
- Whelton A, Watson AJ, Rock RC (1994) In: Tietz textbook of clinical chemistry, edited by C.A Burtis and ER Ashwood (WB Saunders Company, London), p. 1528.
- White-Stevens R (1982) Interference by ascorbic acid in test systems involving peroxidase. 1. Reversible indicators and effects of copper, iron and mercury. *Clin. Chem.* **28**: 578-588.
- Wilson DW, Plass ED (1917) Creatine and creatinine in whole blood and plasma. *J. Biol. Chem.* **29**: 413-423.

- Winqvist F, Lundstrom I (1986) Determination of creatinine by an ammonia-sensitive semiconductor structure and immobilized enzymes. *Anal. Chem.* **58**: 145-148.
- Wisser H, Knoll E (1987) Enzymatisches Verfahren der Kreatininbestimmung. *Internist* **28**: 123-127.
- Witte D, Brown L, Feld R (1978) Effect of bilirubin on detection of hydrogen peroxide by use of peroxidase. *Clin. Chem.* **24**: 1778-1782.
- Wizeman WJ, Kofimas P (2001) Molecularly imprinted polymer hydrogels displaying isomerically resolved glucose binding. *Biomater.* **22**: 1485-1491.
- Wong FCM, Ahmed M, Heng LY, Peng LB (2006) An optical biosensor for dichlovo using stacked sol-gel films containing acetylcholinesterase and a lipophilic chromoionophore. *Talanta* **69**: 888-893.
- Wyss M, Kaddurah-Daouk R (2000) Creatine and creatinine metabolism. *Physiol. Rev.* **80**: 1107-1213.
- Xue GP, Fishlock RC, Snoswell AM (1988) Determination of creatinine in whole blood, plasma, and urine by high-performance liquid chromatography. *Anal. Biochem.* **171**: 135-140.
- Yadav S, Devi R, Bhar P, Singhla S, Pundir CS (2012) A creatinine biosensor based on iron oxide nanoparticles/chitosan-g-polyaniline composite film electrodeposited on Pt electrode. *Enz. Microb. Technol.* **50**: 247-254.
- Yadav S, Devi R, Kumar A, Pundir CS (2011b) Tri-enzyme functionalized ZnO-NPs/CHIT/c-MWCNT/PANI composite film for amperometric determination of creatinine. *Biosens. Bioelectrons.* **28**: 64-70.
- Yadav S, Devi R, Kumari S, Yadav S, Pundir CS (2011a) An amperometric oxalate biosensor based on sorghum oxalate oxidase bound carboxylated multiwalled carbon nanotubes-polyaniline composite film. *J. Biotechnol.* **151**: 212-217.
- Yamato H, Ohwa M, Wemet W (1995) A polypyrrole/three-enzyme electrode for creatinine detection. *Anal. Chem.* **67**: 2776-2780.
- Yan SL, Lin PZ, Hsiao MW (1999) Separation of urea, uric acid, creatine and creatinine by micellar electrokinetic capillary chromatography with sodium cholate. *J. Chromatogr. Sci.* **37**: 45-50.
- Yang YD (1998) Simultaneous determination of creatine, uric acid, creatinine and hippuric acid in urine by high performance liquid chromatography. *Biomed. Chromatogr.* **12**: 47-49.

- Yao T, Kotegawa K (2002) Simultaneous flow-injection assay of creatinine and creatine in serum by the combined use of a 16-way switching valve, some specific enzyme reactors and a highly selective hydrogen peroxide electrode. *Anal. Chim. Acta.* **462**: 283-291.
- Yasuda M, Sugahara K, Zhang J, Ageta T, Nakayama K, Shuin T, Kodama H (1997) Simultaneous determination of creatinine, creatine, and guanidinoacetic acid in human serum and urine using liquid chromatography–atmospheric pressure chemical ionization mass spectrometry. *Anal. Biochem.* **253**: 231-235.
- Yatzidis H (1974) New method for direct determination of “True” creatinine. *Clin. Chem.* **20**: 1131-1134.
- Yokoyama Y, Horikoshi S, Takahashi T, Sato H (2000) Low-capacity cation-exchange chromatography of ultraviolet-absorbing urinary basic metabolites using a reversed-phase column coated with hexadecylsulfonate. *J. Chromatogr. A.* **886**: 297-302.
- Yokoyama Y, Ozaki O, Sato H (1996) Separation and determination of amino acids, creatinine, bioactive amines and nucleic acid bases by dual-mode gradient ion-pair chromatography. *J. Chromatogr. A.* **739**: 333-342.
- Yokoyama Y, Tsuchiya M, Sato H, Kakinuma H (1992) Determination of creatinine and ultraviolet-absorbing amino acids and organic acids in urine by reversed-phase high-performance liquid chromatography. *J. Chromatogr.* **583**: 1-10.
- Yokoyama Y, Tsuji S, Sato H (2005a) Simultaneous determination of creatinine, creatine, and UV-absorbing amino acids using dual-mode gradient low-capacity cation-exchange chromatography. *J. Chromatogr. A.* **1085**: 110-116.
- Yokoyama Y, Yamasaki K, Sato H (2005b) Simultaneous determination of urinary creatinine and UV-absorbing amino acids using a novel low-capacity cation-exchange chromatography for the screening of inborn errors of metabolism. *J. Chromatogr. B: Analyt. Technol. Biomed. Life Sci.* **816**: 333-338.
- Yokozawa T, Fujitsuka N, Oura H, Ienaga K, Nakamura K (1997) In vivo effect of hydroxyl radical scavenger on methylguanidine production from creatinine. *Nephron* **75**: 103.
- Yoshimoto T, Oka I, Tsuru D (1976) Creatine amidinohydrolase of *Pseudomonas putida*: crystallization and some properties. *Arch. Biochem. Biophys.* **177**: 508-515.
- Zhang Y, Tadigadapa S (2004) Calorimetric biosensors with integrated microfluidic channels. *Biosens. Bioelectron.* **19**: 1733-1743.

- Zhao J, Chen H, Ni P, Xu B, Luo X, Zhan Y, Gao P, Zhu D (2011) Simultaneous determination of urinary tryptophan, tryptophan-related metabolites and creatinine by high performance liquid chromatography with ultraviolet and fluorimetric detection. *J. Chromatogr. B Analyt. Technol. Biomed. Life Sci.* **879**: 2720-2725.
- Zhiri A, Houot O, Wellman-Bednawska M, Siest G (1985) Simultaneous determination of uric acid and creatinine in plasma by reversed-phase liquid chromatography. *Clin. Chem.* **31**: 109-112.
- Zinellu A, Carru C, Usai MF, Sotgia S, Deiana L (2004) Creatinine determination in serum by capillary electrophoresis. *Electrophoresis* **25**: 1096-1101.
- Zinellu A, Sotgia S, Zinellu E, Chessa R, Deiana L, Carru C (2006) Assay for the simultaneous determination of guanidinoacetic acid, creatinine, and creatine in plasma and urine by capillary electrophoresis UV detection. *J. Sep. Sci.* **29**: 704-708.