


REFERENCES

- Aleksëev, V.M. (1962). "On the teratology of the trematodes. (Russian Text). Zool. Zhurnal., 41: 453-454.
- Alvarado, R. (1951). El tegumento, la musculature y el parenquima de Fasciola hepatica. Trab. Inst. Cienc. nat. Acosta, 3: 1-90.
- Bácq, Z.M. and Oury, A. (1937). Note sur la repartition de la cholinesterase chez les etres vivants. Acad. Roy. Belg. Bull. Classe des Sciences, 23: 891-893.
- B D H. (1972). Biological Stains and Staining Methods. BDH Chemical Ltd. Poole England. 5th ed. 1972.
- Becejac, S. and Kravavica, S. (1964). On the presence and localisation of alkaline phosphatase activity in the liver fluke (Fasciola hepatica L.). Vet. Arch. 34: 56-59.
- Bennett, C.E. (1975a). Surface features sensory structures and movement of the newly excysted juvenile Fasciola hepatica L. J. Parasitol., 61: 886-891.
- Bennett, C.E. (1975b). Fasciola hepatica : development of caecal epithelium during migration in mouse. Expl. Parasitol., 37: 426-441.

- Bennett, C.E. (1975c). Scanning electron microscopy of Fasciola hepatica L. During growth and maturation in the mouse. J. Parasitol., 61: 892-898.
- Bennett, C.E. and Threadgold, L.T. (1973). Electron microscope studies of Fasciola hepatica XIII. Fine structure of newly excysted juvenile. Expl. Parasitol., 34: 85-99.
- Bennett, C.E. and Threadgold, L.T. (1975). Fasciola hepatica: Development of tegument during migration in mouse. Expl. Parasitol., 38: 38-55.
- Bern, H.A. and Hagadorn, I.R. (1965). Neurosecretion, Vol. 1. In: T.H. Bullock and G.A. Horridge, eds., p. 353-429. San Francisco and London : W.H. Freeman & Co.
- Berthier, L. (1954). Contribution a l' etude du tegument de Fasciola hepatica L. Arch. Zool. exp. gen., 91: 89-102.
- Bettendorf, H. (1897). Uber Muskulatur und Sinneszellen der Trematoden. Zool. Jahrb. Abt. Anat., 10: 307-358.
- Bhalerao, G.D. (1935). Helminth. parasites of the domesticated animals in India. Sci. Monogr. Goun. Agric. Res. India, No. 6, 365 pp.
- Bjorkman, N. and Thorsell, W. (1962). The fine morphology of the mitochondria from the parenchymal cells in the liver-fluke (Fasciola hepatica L.). Exp. Cell. Res., 27:342-346.

- Bjorkman, N. and Thorsell, W. (1964). On the fine structure and resorptive function of the cuticle of the liver fluke, Fasciola hepatica L. Exp. Cell Res., 33: 319-329.
- Bjorkman, N., Thorsell, W. and Lienert, E. (1963). Studies on the action of some enzymes on the cuticle of Fasciola hepatica L. Experientia., 19: 3-5.
- Bugge, G. (1929). Beginnt das Excretionsgefasssystem der Fasciola hepatica mit terminalorganen? Berl. tierarztl. Wschr., 45: 557-561.
- Burton, P.R. (1960). Gametogenesis and fertilization in the frog lung fluke, Haematoloechus medioplexus Stafford (Trematoda:Plagiorchiidae). J. Morphol., 107: 93-121.
- Burton, P.R. (1963). A histochemical study of vitelline cells, egg capsules and Mehlis' gland in the frog lung fluke, Haematoloechus medioplexus. J. exp. Zool., 154 : 247-258.
- Chance, M.R.A. and Mansour, T. (1953). A contribution to the pharmacology of movement of the liver fluke. Brit. J. Pharm., 8: 134-138.
- *Clegg, J.A. (1957). Studies on the maintenance of Fasciola hepatica in vitro. Ph.D., thesis, University of London.

- *Cobbold, T.S. (1855). Description of a new trematode (Fasciola gigantica). Edinb. N. Phil. J. n. s., 2 : 262-266.
- Dawes, B. (1940). Notes on the formation of the egg capsules in the monogenetic trematode, Hexacotyl extensicauda Dawes, 1940. Parasitology, 32: 287-295.
- Dawes, B. (1946). The Trematoda. 1st edit, Cambridge University Press. 644 pp.
- Dawes, B. (1954). Maintenance in vitro of Fasciola hepatica. Nature, 174 : 654-655.
- Dawes, B. (1962). A histological study of the caecal epithelium of Fasciola hepatica L. Parasitology, 52 : 483-493.
- Dawes, B. (1963a). The migration of juvenile forms of Fasciola hepatica L. through the wall of the intestines in the mouse, with some observations on food and feeding. Parasitology, 53: 109-122.
- Dawes, B. (1963b). Hyperplasia of the bile duct in fascioliasis and it's relation to the problem of maturation in the liver-fluke. Parasitology, 53: 123-134.
- Dawes, B. (1963c). Some observations of Fasciola hepatica L. during feeding operations, in the hepatic parenchyma of the mouse, with notes on the nature of liver damage in the host. Parasitology, 53 : 135-144.



- Dissamarn, R. (1955). La douve du foie, du boeuf et du buffle en Thaïlande. Bull. Off. int. Epiz., 43: 435-438.
- Dixon, K.E. and Mercer, E.H. (1965). The fine structure of the nervous system of the cercaria of the liver fluke Fasciola hepatica L. J. Parasitol., 51: 962-976.
- Fraipont, J. (1880). Recherches sur l'appareil excréteur des Trematodes et des Cestodes. I. Arch. Biol., Gand., 1 : 415-456.
- Gallagher, S.S.E. and Threadgold, L.T. (1967). Electron microscope studies of Fasciola hepatica II. The interrelationship of the parenchyma with other organ systems. Parasitology, 57: 627-632.
- Gallego-Berenguer, J. (1952). Use caso de teratologia en trematodes. Atrofia unilateral total de la glandula vitelogenica on Fasciola hepatica. Rev. iber. Parasit., 12: 65-67.
- Goil, M.M. (1958a). Fat metabolism in trematode parasites. Z. Parasitkde., 18: 320-323.
- Goil, M.M. (1958b). Protein metabolism in trematode parasites. J. Helminth., 32: 119-124.
- Goil, M.M. (1961). Physiological studies on trematodes; Fasciola gigantica; carbohydrate metabolism. Parasitology, 51: 335-337.

- Gomori, C. (1952). Microscopic Histochemistry. University of Chicago Press, Chicago. 273 pp.
- Govaert, J. (1953a). Deoxyribonucleic acid content of the germinal vesicle of the ovocyte in Fasciola hepatica. Nature, 172 : 302-303.
- Govaert, J. (1953b). Sur la teneur en acide desoxyribonucleique du noyae des cellules vitellogenes chez Fasciola hepatica. C. R. Soc. Biol., Paris, 147 : 1494-1496.
- Govaert, J. (1954). La teneur en acide desoxyribonucleique des noyaux des elements de la lignee spermatique chez Fasciola hepatica. Biol. Jahrb., 21 : 202-209.
- Govaert, J. (1955). Etude quantitative de L'acide desoxy-pentose-nucleique loss de la maturation et de la fecondation de l'eeuf chez Fasciola hepatica. C. R. Soc. Biol., Paris, 149 : 1066-1069.
- Govaert, J. (1960). Etude cytologique et cytochimique des cellules de la lignee germinative chez Fasciola hepatica. Expl. Parasitol., 9 : 141-158.
- Grasso, M. (1967a). Prime indagini sulla presenza di cellule neurosecretrici in Fasciola hepatica. R. C. Acad. Naz. Lincei., 42 : 85-87.

- Grasso, M. (1967b). Prime indagini sulla presenza di cellule neurosecretrici in Fasciola hepatica. R.C. Acad. Naz. Lincei., 42 : 903-905.
- Grasso, M. (1972). Ultrastructural studies in neurosecretion and gamete ripening in platyhelminthes. Gen. comp. Endocr., 18 : 3.
- Gresson, R.A.R. (1957). Spermateleosis in Fasciola hepatica. Quart. J. micr. Sci., 98 : 493-498.
- Gresson, R.A.R. and Threadgold, L.T. (1959). A light and electron microscope study of the epithelial cells of the gut of Fasciola hepatica L. J. biophys. biochem. Cytol., 6 : 157-162.
- Gresson, R.A.R. and Threadgold, L.T. (1964). The large neurons and interstitial material of Fasciola hepatica L. Proc. roy. Soc. Edinb., 42 : 261-266.
- Halton, D.W. (1966). Occurrence of microvilli-like structures in the gut of digenetic trematodes. Experientia, 22: 828-829.
- Halton, D.W. (1967a). Studies on phosphatase activity in Trematoda. J. Parasitol., 53: 46-54.
- Halton, D.W. (1967b). Histochemical studies of carboxylic esterase activity in Fasciola hepatica. J. Parasitol., 53 :1210-1216.

- Hanna, R.E.B. (1975). Fasciola hepatica: An electron microscope autoradiographic study of protein synthesis and secretion in gut cells in tissue slices. Expl. Parasitol., 38 : 167-180.
- Havet, J. (1900). Contribution a l'etude du systeme nerveux des Trematodes (Distomum hepaticum). La Cellule, 17: 353-381.
- Healy, G.R. (1955). Anomalies in Fasciola hepatica. J. Parasitol., 41 : 35.
- Hockley, D.J. (1973). Ultrastructure of the tegument of Schistosoma. In "Advances in Parasitology" (Ben Dawes, ed). Academic Press, New York and London, 11: 233-305.
- Hyman, L.H. (1951). The Invertebrates: Platyhelminthes and Rhynchocoela. The acoelomate Bilateria. VII + 550 pp. New York. Mc Graw Hill 1951.
- (1970)
- Irwin, S.W.B. and Threadgold, L.T. Electron-microscope studies on Fasciola hepatica. VIII. The development of the Vitelline cells. Expl. Parasitol., 28 : 399-411.
- Jackson, H.G. (1921). A revision of the genus Fasciola, with particular reference to F. gigantica (Cobbold) and F. nyanzi (Leiper). Parasitology, 13 : 48-56.

- Kalyankar, S.D. and Kankal, N.C. (1980). Histology and histochemical observations on the neurosecretory cells in Proalrioides tropidonotis (Trematoda:Digenea). (Abstract). Indian J. Parasitol., 3 (Suppl), B34: 21.
- Kawana, H. (1940). Study on the development of the excretory system of Fasciola hepatica L. with special reference to its first intermediate host in Central China. J. Shanghai Sci. Inst., 5: 13-34.
- Kendall, S.B. (1954). Fascioliasis in Pakistan. Ann. trop. Med. Parasit., 48: 307-313.
- Kibakin, V.V. (1961). The epizootology of fascioliasis in sheep in the Turkmen S.S.R. Conference (5th) on Problems in Parasitology. Alma Ata, 1959, 389-393.
- Kishore, S. and Shyamasundari, K. (1980). Nervous system and neurosecretory cells in a trematode Encyclometra colubriamurorum (Digenea:Plagiorchidae) (Abstract). Indian J. Parasitol., 3 (Suppl.), B36: 22.
- Kümmel, G. (1959). Das Terminalorgan der Protonephridien, Feinstruktur und Deutung der Funktion. Z. Naturf., 13 B: 677-679.
- Kümmel, G. (1960). Feinstruktur der Wimperflamme in der Protonephridien. Protoplasma, 51: 371-376.

Lal, M.B. and Shrivastava, S.C. (1960). Some histochemical observations on the cuticle of Fasciola indica Varma 1953. Experientia, Basle., 16 : 185-186.

*Lang, A. (1880). Untersuchungen zur vergleichenden Anatomie und Histologie des Nervensystems der Plathelminthen. II. Über das Nervensystem der Trematoden. Mitt. Zool. Sta. Neapel, 2 : 28-52.

Lee, D.L. (1966). The structure and composition of the helminth cuticle. In "Advances in Parasitology" (Ben Dawes, ed.), Academic Press, New York and London, 5 : 187-284.

Lee, D.L. (1972). The structure of the helminth cuticle. In "Advances in Parasitology". (Ben Dawes ed)., Academic Press, New York and London, 10 : 347-379.

LeFlore, W.B. and Smith, B.F. (1976). The histochemical localization of esterases in whole mounts of Cysticercus fasciolaris. Trans. Amer. Microsc. Soc., 95: 73-79.

*Leuckart, R. (1881). Zur Entwicklungsgeschichte des Leberegels. Zool. Anz., 99: 641-646.

Leuckart, R. (1886). Die Parasiten des Menschen und die von ihnen herrührenden Krankheiten. Ein Hand und Lehrbuch für Naturforscher und Aerzte. 2. Aufl., V 1,3: 855-1000 pp. Leipzig. u. Heidelberg.

- Lillie, R.D. (1965). Histopathologic Technic and Practical Histochemistry, 3rd (ed.). Mc Graw-Hill, New York, pp. 715.
- Looss, A. (1885). Beitrage zur Kenntnis der Trematoden. Z. Wiss. Zool., 41 : 390-446.
- Lutz, P.L. & Siddiqi, A.H. (1967). Composition of haemoglobin of Fasciola gigantica (Trematoda:Digenea) and it's host. Expl. Parasitol., 20 : 83-87.
- Lyons, K.M. (1977). Epidermal adaptations of Parasitic Platyhelminths. In "Comparative Biology of Skin" (Spearman, eds.). Symp. Zool. Soc. London., 39: 97-144.
- Macy, R.W. (1950). Anomalies in the trematodes, Prosthogonimus macrorchis with special reference to rudimentary vitellaria and effect on egg shell formation. Trans. Amer. microsc. Soc., 69: 47-49.
- Madhavi, R. and Rao, H.K. (1972). A specimen of Astiotrema monticellii Stossich, 1904, with anomalous female reproductive system. J. Parasitol., 58: 1012-1014.
- Manter, H.W. (1931). Some abnormalities of Trematodes. J. Parasitol., 13: 199-202.
- Monne, L. (1959). On the external cuticles of various helminths and their role in the host-parasite relationship. Ark. Zool. Stockholm, 12: 343-358.

- Müller, W. (1923). Die Nahrung von Fasciola hepatica und ihre Verdauung. Zool. Anz., 57 : 273-281.
- Nacheva, L.V. (1977). Cytological and histochemical study of the integument and intestine of some digeneans. (Russian text). Trud. Buryat. Inst. Estestv. Nauk. (Faun, Ekol. Issledov. V. Zab.).
- Ortner-Schoenbach, P. (1913). Zur Morphologie des Glykogens bei trematoden und Cestoden. Arch. Zellforsch., 11: 413-349.
- Pantelouris, E.M. (1964). Localization of glycogen in Fasciola hepatica L. and an effect of insulin. J. Helminth., 38 : 283-286.
- Pantelouris, E.M. (1965). The Common Liver Fluke. Pergamon Press, London, 259 pp.
- Pantelouris, E.M. and Gresson, R.A.R. (1960). Autoradiographic studies on Fasciola hepatica L. Parasitology, 50:165-169.
- Pantelouris, E.M. and Hale, P. (1962). Iron and Vitamin C in Fasciola hepatica L. Res. vet. Sci., 3 : 300-303.
- Pantelouris, E.M. and Threadgold, L.T. (1963). The excretory system of the adult Fasciola hepatica L. La Cellule, 64 : 63-67.

- Patnaik, M.M. (1971). Studies on geographical distribution and ecology of Fasciola gigantica with special reference to its intermediate host. Lymnaea auricularia var. rufescens in Orissa. Indian J. Helminth., 23 : 115-134.
- Pearse, A.G.E. (1960). Histochemistry: Theoretical and Applied. 2nd ed. J & A. Churchill. 998 pp.
- Prenant, M. (1922). Recherches sur la parenchyme des helminthes, essai d'histologie comparee. Arch. Morphol., 5 : 1-474.
- Price, E.W. (1930). Some malformations in trematodes (Abstract of report before Helminth Soc. Washington, Sept. 19, 1929). J. Parasitol., 16 : 161.
- Ramisz, A. (1967). Studies on the nervous system of nematodes and cestodes by means of histochemical method for active acetylcholinesterase. III. Nematodes of the genus Capillaria and cestodes of the genus Dilepis and Choanotaenia. Acta. Parasitol. Polon., 14: 365-385.
- Rao, K.H. (1959a). Observations on the Mehlis' gland complex in the liver fluke, Fasciola hepatica. J. Parasitol., 45 : 347-351.
- Rao, K.H. (1959b). Histochemistry of Mehlis' gland and egg-shell formation in the liver fluke. Experientia, 15: 464.

- Rao, K.H. (1960). The problem of Mehlis' gland. Parasitology, 50 : 349-350.
- Rao, K.H. and Madhavi, R. (1962). Mehlis' gland complex in the cattle liver fluke Fasciola gigantica Cobbold. Curr. Sci. Bangalore., 31 : 101-102.
- Robinson, G. and Threadgold, L.T. (1975). Electron microscope studies of Fasciola hepatica XII. The fine structure of the gastrodermis. Expl. Parasitol., 37: 20-36.
- Rohde, K. (1965). The cell types found in the pharynx of Polystomoides malayi Rohe, 1963. Med. J. Malaya, 20:55.
- Roy, S.C. (1954). Liver fluke disease amongst cattle in the district of Darjeeling with it's main source of infection. Proc. Indian. Sc. Congr., 41st (1954), Part III: 224-225.
- Sarwar, M.M. (1957). Fasciola indica Varma, a synonym of F. gigantica Cobbold. Biologia, Lahore, 3 : 168-175.
- Schardein, J.L. and Waitz, J.A. (1965). Histochemical studies of esterases in the cuticle and nerve cords of four cyclophyllidean cestodes. J. Parasitol., 51: 366-363.
- Schubmann, W. (1905). Uber die Eibildung und Embryonalentwicklung von Fasciola hepatica L. Zool. Jb. Anat., 21 : 571-606.

- Sekardi, L. and Ehrlich, I. (1962). Acetilkolinesteraza velikog metilja (Fasciola hepatica L.). Biol. Glasnik., 15: 229-233.
- Senft, A.W. (1959). Electron microscope study of the integument, flame cells and gut of the human parasite, Schistosoma mansoni. Biol. Bull. Woods Hole, 117-387.
- Senft, A.W., Philpott, D.E. and Pelofsky, A.H. (1961). Electron microscope observations of the integument, flame cells, and gut of Schistosoma mansoni. J. Parasitol., 47: 217-229.
- Shield, J. (1969). Dipylidium caninum, Echinococcus granulosus and Hydatigera taeniaeformis : histochemical identification of cholinesterases. Expl. Parasitol., 25: 217-231.
- Shyamasundari, K. and Rao, K.H. (1975). The structure and cytochemistry of the neurosecretory cells of Fasciola gigantica Cobbold and Fasciola hepatica L. Z. parasitkde. 47: 103-109.
- Siddiqi, A.H. and Lutz, P.L. (1966). Osmotic and ionic regulation in Fasciola gigantica (Trematoda : Digenea). Expl. Parasitol., 19: 348-357.

- Smyth, J.D. (1951). Egg-shell formation in Trematodes and Cestodes as demonstrated by the methyl or malachite green techniques. Nature (Lond.), 168: 322-323.
- Smyth, J.D. and Clegg, J.A. (1959). Egg shell formation in trematodes and cestodes. Expl. Parasitol., 8: 286-323.
- Sommer, F. (1880). Die Anatomie des Leberegels Distomum hepaticum L. Z. Wiss. Zool., 34: 539-640.
- Stein, P.C. and Lumsden, R.D. (1973). Schistosoma mansoni: topochemical features of cercariae, schistosomulae and adults. Expl. Parasitol., 35 : 499-514.
- Stephenson, W. (1947a). Physiological and histochemical observations on the adult liver fluke, Fasciola hepatica L. Survival in vitro. Parasitology, 38: 116-122.
- Stephenson, W. (1947b). Physiological and histochemical observations on the adult liver fluke, Fasciola hepatica L. II. Feeding. Parasitology, 38 : 123-127.
- Stephenson, W. (1947c). Physiological and histochemical observations on the adult liver fluke, Fasciola hepatica L. III. Egg-shell formation. Parasitology, 38 : 128-139.

- Stephenson, W. (1947d). Physiological and histochemical observations on the adult liver fluke, Fasciola hepatica L. IV. The excretory system. Parasitology, 38 : 140-144.
- *Stossich, M. (1892). I distomi degli ucelli. Lavoromonografico. Boll. Soc. Adriat. Sc. Nat. Trieste., 13: 143-196.
- Tay, J. and Biagi, F.F. (1968). Ultrastructura de F. hepatica I. Cuticula, glandulas vitelogen as, Ciegos intestinales y Cellulas en flama. La prensa med. Mex. 33: 397-405.
- Thapar, G.S. and Tandon, R.S. (1952). On the life history of liver-fluke, Fasciola gigantica Cobbold, 1885, in India. Indian J. Helm., 4 : 1-36.
- *Thomas, A.P. (1881). Reports of experiments on the development of the liver-fluke (Fasciola hepatica). J. Roy Agric. Soc. England, 17 : 1-29.
- *Thomas, A.P. (1882). The rot in sheep, or the life history of the liver-fluke. Nature, 26 : 606-608.
- Thomas, A.P. (1883a). The life history of the liver fluke. Quart. J. Micr. Sci., 23 : 99-133.
- *Thomas, A.P. (1883b). The natural history of the liver fluke and the prevention of rot. J. Roy Agric. Soc. (England), 19 : 276-305.

- Thorsell, W. and Bjorkman, N. (1965). Morphological and biochemical studies on absorption and secretion in the alimentary tract of Fasciola hepatica L. J. Parasitol., 51 : 217-223.
- Threadgold, L.T. (1963a). The ultrastructure of the 'cuticle' of Fasciola hepatica. Expl. Cell Res., 30: 240-242.
- Threadgold, L.T. (1963b). The tegument and associated structures of Fasciola hepatica. Q. Jl. microsc. Sci., 104: 505-512.
- Threadgold, L.T. (1967). Electron microscope studies of Fasciola hepatica. III. Further observations on the tegument and associated structures. Parasitology, 57: 633-637.
- Threadgold, L.T. (1968). Electron microscope studies of Fasciola hepatica. IV. The ultrastructural localization of phosphatases. Expl. Parasitol., 23: 264-276.
- Threadgold, L.T. (1975a). Electron microscope studies of Fasciola hepatica. IV. The fine structure of the prostate gland. Expl. Parasitol., 37: 117-124.
- Threadgold, L.T. (1975b). Fasciola hepatica: the ultrastructure of the epithelium of the seminal vesicle, the ejaculatory duct and the cirrus. Parasitology, 71: 437-444.
- Threadgold, L.T. and Gallagher, S.S. (1966). Electron microscope studies of Fasciola hepatica, I. The ultrastructure and Interrelationship of the parenchymal cells. Parasitology, 56: 299-304.

- Ude, J. (1962). Neurosekretorische Zellen im Cerebralganglion von Dicrocoelium lanceatum St. u. M. (Trematoda-Digenea). Zool. Anz., 169: 455-457.
- Varma, A.K. (1953). On Fasciola indica n. sp. with some observations on F. hepatica and F. gigantica. J. Helminth., 23 : 185-198.
- Venkataramakrishna, G., Kunde, S.H. and Simha, S.S. (1980). Studies on the neurosecretory cells of Fischoederius cobboldi (Trematoda:Digenea) (Abstract). Indian J. Parasitol. 3 (Suppl), B 61 : 30.
- von Brand, T. and Mercado, T.T. (1961). Histochemical glycogen studies on Fasciola hepatica. J. Parasitol., 47:459-463.
- von Brand, T. and Weinland, E. (1924). Über tropfenformige Ausscheidungen bei Fasciola hepatica (Distomum hepaticum). Z. Wiss. Biol., C, 2: 209-214.
- Watanabe, S. (1958). General review on Fascioliasis in Japan. J. Jap. Vet. med. Ass., 11: 293-299.
- Watanabe, S. and Iwata, S. (1956). The morphological studies on the genus Fasciola in Japan. II. (Proc. 41 Med. Japan. Soc. Vet. Sc., Tokyo, Apr. 5-6). Japan. J. Vet. Sc., Tokyo, 8 (Suppl) : 62-63.

- Watanabe, S. and Iwata, S. (1958). Morphological studies of the genus Fasciola in Japan, with particular reference to F. gigantica Cobbold. (Proc. 27 Ann. Meet. Japan Soc. Parasitol). Kiseichugaku Zasshi, 7: 281.
- Watanabe, S. and Ueno, H. (1959). Morphological studies of the genus Fasciola in Japan, with particular reference to F. hepatica and F. gigantica (Proc. 28 Ann. Meet. Japan. Soc. Parasitol., Apr. 6-7). Kiseichugaku Zasshi, 6: 368-369.
- Watanabe, S. and Ueno, H. (1960a). A revision of the genus Fasciola, with particular reference to Japanese species. Kiseichugaku Zasshi, 9 : 379.
- Watanabe, S. and Ueno, H. (1960b). Morphological studies on the genus Fasciola with particular reference to Japanese species. Japan. J. Vet. Sc. Tokyo, 22: 472.
- *Weinland, E. and von Brand, T. (1926). Beobachtungen au Fasciola hepatica. Z. vergl. Physiol., 4: 212-285.
- Willey, C.H. (1930). Studies on the lymph system of digenetic trematodes. J. Morphol., 50: 1-38.
- Wilson, V.C.L.C. and Schiller, E.L. (1969). The neuroanatomy of Hymenolepis diminuta and H. nana. J. Parasitol., 55 : 261-270.

*Wisniewski, L.W. (1930). Rodraj anatomiezne, histogenetzne, systematyezne i biologiezne. Das Genus Archigelès R. Leuck. Eine studie zur Biologie (German text). Med. et. Nat., 2 : 1-160.

Wootton, R.M. and Sogandáres-Bernal, F. (1963). A report on the occurrence of microvillus-like structures in the caeca of certain trematodes (Paramphistomatidae). Parasitology, 53: 157-161.

Yamao, Y. and Saito, A. (1952). Histochemical studies on endoparasites. IV. Distribution of glyceromonophosphatases in the tissues of the liver fluke, Fasciola hepatica L. Zikken Seibutsugaku Ho., 2: 153-158.

Yosufzai, H.K. (1952a). Cytological studies on the spermatogenesis of Fasciola hepatica L. Cellule, 55: 5-19.

Yosufzai, H.K. (1952b). Female reproductive system and egg-shell formation in Fasciola hepatica L. Nature, 169: 549.

Yosufzai, H.K. (1953a). Cytological studies on the oogenesis of Fasciola hepatica L. Cellule, 55 : 23-42.

Yosufzai, H.K. (1953b). Shell gland and egg-shell formation in Fasciola hepatica L. Parasitology, 43 : 88-93.

* References not seen in original.